화학특강 **(2010년 2학기)**


제2장 DNA, 단백질 그리고 Biotechnology

황현석

강원대학교 화학과

2010년 9월 27일 월요일

DNA Origami




Rothemund, Nature **440**, 297 (2006)


Structures of Deoxyribonucleic Acid (DNA) and Ribonucleic Acid (RNA)

- Structure of Nucleotide

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



- Deoxyribose and ribose




Chang, 25.4


KNU

Structures of Deoxyribonucleic Acid (DNA) and Ribonucleic Acid (RNA)


- Purine bases





- Pyrimidine bases



- DNA double helix








Chang, 25.4

KNU

History of DNA

- Chargaff's rules in the 1940's
  1. The amount of adenine (a purine) is equal to that of thymine (a pyrimidine); that is A = T, or A/T = 1.
  2. The amount of guanine (a purine) is equal to that of cytosine (a pyrimidine); that is G = C, or G/C = 1.
  3. The total number of purine bases is equal to the total number of pyrimidine bases; that is, A + G = T + C .

- X-ray diffraction photograph by R. Franklin





Chang, 25.4

History of DNA

- Nature paper by Watson & Crick




MOLECULAR STRUCTURE OF NUCLEIC ACIDS

A Structure for Deoxyribose Nucleic Acid

Watson & Crick, Nature 171, 737 (1953)

- Nature paper by Wilkins




Molecular Structure of Deoxypentose Nucleic Acids

Wilkins, Stoke, and Wilson, Nature 171, 738 (1953)

©1953 Nature Publishing Group

History of DNA

- Nature paper by Rosalind Franklin



740 NATURE April 25, 1953 Vol. 171

We wish to thank Prof. J. T. Randall for encouragement; Profs. E. Chargaff, R. Signer, J. A. V. Butler and Drs. J. D. Watson, J. D. Smith, L. Hamilton, J. C. White and G. R. Wyatt for supplying material without which this work would have been impossible; also Drs. J. D. Watson and Mr. F. H. C. Crick for stimulation, and our colleagues R. E. Franklin, R. G. Gosling, G. L. Brown and W. E. Seeds for discussion. One of us (H. R. W.) wishes to acknowledge the award of a University of Wales Fellowship.

M. H. F. Wilkins
Medical Research Council Biophysics Research Unit,

A. R. Stokes
H. R. Wilson
Wheatstone Physics Laboratory, King's College, London.
April 2.

Sodium deoxyribose nucleate from calf thymus. Structure B

Molecular Configuration in Sodium Thymonucleate

Sodium thymonucleate fibres give two distinct types of X-ray diagram. The first corresponds to a crystalline form, structure A, obtained at about 75 per cent relative humidity; a study of this is described in detail elsewhere[3]. At higher humidities a different structure, structure B, showing a lower degree of order, appears and persists over a wide range of ambient humidity. The change from A to B is reversible. The water content of structure B fibres which undergo this reversible change may vary from 40–50 per cent to several hundred per cent of the dry weight. Moreover, some fibres never show structure A, and in these structure B can be obtained with an even lower water content.

The X-ray diagram of structure B (see photograph) shows in striking manner the features characteristic of helical structures, first worked out in this laboratory by Stokes (unpublished) and by Crick, Cochran and Vand[4]. Stokes and Wilkins were the first to propose such structures for nucleic acid as a result of direct studies of nucleic acid fibres, although a helical structure had been previously suggested by Furberg (thesis, London, 1949) on the basis of X-ray studies of nucleosides and nucleotides.

While the X-ray evidence cannot, at present, be taken as direct proof that the structure is helical, other considerations discussed below make the existence of a helical structure highly probable.

Structure B is derived from the crystalline structure A when the sodium thymonucleate fibres take up quantities of water in excess of about 40 per cent of their weight. The change is accompanied by an increase of about 30 per cent in the length of the fibre, and by a substantial re-arrangement of the molecule. It therefore seems reasonable to suppose that in structure B the structural units of sodium thymonucleate (molecules or groups of molecules) are relatively free from the influence of neighbouring molecules, each unit being shielded by a sheath of water. Each unit is then free to take up its least-energy configuration independently of its neighbours and, in view of the nature of the long-chain molecules involved, it is highly likely that the general form will be helical[5]. If we adopt the hypothesis of a helical structure, it is immediately possible, from the X-ray diagram of structure B, to make certain deductions as to the nature and dimensions of the helix.

The innermost maxima on the first, second, third and fifth layer lines lie approximately on straight lines radiating from the origin. For a smooth single-strand helix the structure factor on the nth layer line is given by:

$$F_n = J_n(2\pi rR)\exp i n(\psi + \tfrac{1}{2}\pi),$$

where $J_n(u)$ is the nth-order Bessel function of u, r is the radius of the helix, and R and ψ are the radial and azimuthal co-ordinates in reciprocal space[6]; this expression leads to an approximately linear array of intensity maxima of the type observed, corresponding to the first maxima in the functions J_1, J_2, J_3, etc.

If, instead of a smooth helix, we consider a series of residues equally spaced along the helix, the transform in the general case treated by Crick, Cochran and Vand is more complicated. But if there is a whole number, m, of residues per turn, the form of the transform is as for a smooth helix with the addition, only, of the same pattern repeated with its origin at heights $m\sigma^*$, $2m\sigma^*$ . . . etc. (σ is the fibre-axis period).

In the present case the fibre-axis period is 34 A. and the very strong reflexion at 3·4 A. lies on the tenth layer line. Moreover, lines of maxima radiating from the 3·4-A. reflexion as from the origin are visible on the fifth and lower layer lines, having a J_4 maximum coincident with that of the origin series on the fifth layer line. (The strong outer streaks which apparently radiate from the 3·4-A. maximum are not, however, so easily explained.) This suggests strongly that there are exactly 10 residues per turn of the helix. If this is so, then from a measurement of R_0 the position of the first maximum on the nth layer line (for n ≠ 0–C), the radius of the helix, can be obtained. In the present instance, measurements of R_1, R_2, R_3 and R_4 all lead to values of r of about 10 A.

©1953 Nature Publishing Group

Franklin & Gobling, Nature 171, 740 (1953)

The Nobel Prize in Physiology or Medicine 1962

"for their discoveries concerning the molecular structure of nucleic acids and its significance for information transfer in living material"



Francis Harry Compton Crick

1/3 of the prize

United Kingdom

MRC Laboratory of Molecular Biology
Cambridge, United Kingdom



James Dewey Watson

1/3 of the prize

USA

Harvard University
Cambridge, MA, USA





Maurice Hugh Frederick Wilkins

1/3 of the prize

United Kingdom and New Zealand


London University
London, United Kingdom














Amino Acids

- **Amino acids**: the basic structural units of proteins. An amino acid is a compound that contains at least one amino group (-NH2) and at least one carboxyl group (-COOH)



Chang, 25.4

TABLE 25.1 The 20 Amino Acids Essential to Living Organisms*


Name	Abbreviation	Structure
Alanine	Ala	
Arginine	Arg	
Asparagine	Asn	
Aspartic acid	Asp	
Cysteine	Cys	
Glutamic acid	Glu	
Glutamine	Gln	
Glycine	Gly	
Histidine	His	
Isoleucine	Ile	


(Continued)

*The shaded portion is the R group of the amino acid.

Proteins

- **Proteins**: A polymer of amino acids through peptide bonds.






Chang, 25.4


Condensation reaction

Protein Structure


- α helix



- β sheet



- Primary, secondary, and tertiary structures



Chang, 25.4


KNU

Polymerase Chain Reaction (PCR)

- **PCR**
  - A technique to amplify a piece of DNA
  - DNA template (target), two primers, DNA polymerase (Taq polymerase), deoxinucleoside triphosphates.

- Procedure
  - Initialization: heating to 94-96°C
  - Denaturation: yielding ssDNAs
  - Annealing: lowering to 50-65°C and annealing primers to ssDNAs
  - Elongation: inducing DNA polymerases to synthesize a new DNA strand

- Applications of PCR
  - Isolation of genomic DNA
  - DNA sequencing
  - Forensic analysis
  - Disease diagnosis



KNU

Polymerase Chain Reaction (PCR)




The Nobel Prize in Chemistry 1993

"for contributions to the developments of methods within DNA-based chemistry"

"for his invention of the polymerase chain reaction (PCR) method"


"for his fundamental contributions to the establishment of oligonucleotide-based, site-directed mutagenesis and its development for protein studies"


Kary B. Mullis



Michael Smith


KNU

What are ion channels?



Voet and Voet, Biochemistry (3rd. Ed.)





KcsA ion channel
Science **280,** 69 (1998)




KNU

Chemistry

The Nobel Prize in Chemistry 2003

"for discoveries concerning channels in cell membranes"

"for the discovery of water channels"



Peter Agre

"for structural and mechanistic studies of ion channels"





Roderick MacKinnon

KNU


DNA translocation

- Alpha hemolysin: an ion channel






- Alpha hemolysin inside a lipid bilayer




KNU

DNA translocation





Miller et al., PRL **86**, 3435 (2001)




Schulten et al., PNAS **102**, 12377 (2005)

DNA Transfection








Lipid phases in lipid-DNA complex (=lipoplex)

- Lamellar(L_α^c) and hexagonal(H_{II}^c) phases.



(Science, **281**, 78 (1998))

Beer-Lambert Law

- Derivation of Beer-Lambert Law

$$dI = -\kappa[J]Idl \Rightarrow \frac{dI}{I} = -\kappa[J]dl$$

$$\int_{I_0}^{I} \frac{dI}{I} = -\kappa[J]\int_0^l dl \Rightarrow \ln\frac{I}{I_0} = -\kappa[J]l$$

$$\ln\frac{I}{I_0} = -\kappa[J]l \Rightarrow \log\frac{I}{I_0} = -\log e\kappa[J]l = -\varepsilon[J]l$$

- Absorbance

$$\boxed{A = \log\frac{I_0}{I} = \varepsilon[J]l}$$

Atkins, Physical Chemistry (8th ed.) 432-433




Fig. 13.4 The integrated absorption coefficient of a transition is the area under a plot of the molar absorption coefficient against the wavenumber of the incident radiation.

Colors and Complementary Colors

The colors of the visible light


color	wavelength interval	frequency interval
red	700–635 nm	430–480 THz
orange	635–590 nm	480–510 THz
yellow	590–560 nm	510–540 THz
green	560–490 nm	540–610 THz
blue	490–450 nm	610–670 THz
violet	450–400 nm	670–750 THz

Complementary colors



Hyperchromic effect of DNA

- DNA melting curve





Relative absorbance: the ratio of the absorbance at the indicated temperature to that at 25°C.

- Extinction coefficients of ss- and dsDNA at 260 nm
  1. For ssDNA: 0.027 $(\mu g/ml)^{-1}cm^{-1}$
  2. For dsDNA: 0.020 $(\mu g/ml)^{-1}cm^{-1}$

- Hyperchromic effect of DNA: when DNA denatures, its UV absorbance, which is almost entirely due to its aromatic bases, increases by ~40 % at all wavelengths.

Selective Colorimetric Detection of DNA






Mirkin et al., Nature **382**, 607 (1996)

KNU

Selective Colorimetric Detection of DNA


높은 온도 (58.5°C 이상)





n Polynucleotide targets



낮은 온도 (58.5°C 이하)








일반 DNA 이중나선의 상전이 그림


DNA 금나노입자의 상전이 그림

Mirkin et al., Science 277, 1078 (1997)

KNU


Selective Colorimetric Detection of DNA






Mirkin et al., Science 277, 1078 (1997)


Bio-Bar-Code-Based DNA Detection





Mirkin et al., JACS **126**, 5932 (2004)

Remote Electronic Control of DNA Hybridization through Nanoscale Antenna




Hamad-Schifferli et al., Nature **415**, 152 (2002)

KNU


Aptamer

Aptamer: 핵산, 단백질, 금속이온과 같이 생체 내의 주요물질에 높은 결합력과 선택성을 가진 단일사슬 핵산 (single-stranded DNA)이나 단백질로서, 단백질로 이루어진 항체와는 달리 화학적, 물리적, 생물학적 안정성이 크다. (aptamer means “to fit” in Greek)



KNU


Colorimetric Sensing of Adenosine Based on Aptamers




Nutiu and Li, JACS **125**, 4771 (2003)

KNU

Colorimetric Sensing of Cocaine Based on Aptamers Attached to Gold Nanoparticles






cocaine

DNA 금나노입자가 뭉쳐있을 때 → 700 nm 흡수, 파랑 또는 보라색


DNA 금나노입자가 떨어져 있을 때 → 522 nm 흡수, 붉은색




Liu and Lu, Angew. Chem. Int. Ed. **45**, 90 (2006)


KNU


Colorimetric detection of DNA sequences based on electrostatic interactions






citrate








Li & Rothberg, PNAS **101**, 14036 (2004)

Colorimetric Sensing of Cocaine Based on Aptamers and Gold Nanoparticles





Zhang et al., Small **4**, 1196 (2008)

IgNobel Prizes

IMPROBABLE RESEARCH

Research that makes people LAUGH and then THINK

Subscribe

Search go

- Home
- What is Improbable Research?
- Ig Nobel Prizes
  - Info
  - Ceremony & lectures
  - The winners
- Publications
  - Magazine (Annals of Improbable Research)
  - Newsletter (mini-AIR)
  - Newspaper column
  - Classics
- Improbable TV
- Improbable Research Events
- Subscribe
- Press Clips
- Luxuriant Hair Clubs for Scientists
- Store
- Contact Us



The Ig® Nobel Prizes

The **Twentieth 1st Annual Ig Nobel Prize ceremony** will happen on **Thursday, September 30, 2010**.
Tickets to the ceremony are now sold out. The event will be webcast live.

The Ig Nobel Prizes honor achievements that first make people **laugh**, and then make them **think**. The prizes are intended to celebrate the unusual, honor the imaginative — and spur people's interest in science, medicine, and technology.

> "Last, but not least, there are the Ig Nobel awards. These come with little cash, but much cachet, and reward those research projects that 'first make people laugh, and then make them think'" — *Nature*

- **The winners** (the complete list, from 1991 through the present)
- **Recent news** (the Ig blog)
- **The ceremonies** (all years, including 2009 — details, video, opera librettos)
- **Some thoughts** about the Igs, about some of the winners, about what they may have done, what the Igs are, what they aren't, and what it could all mean, maybe.



The 2009 Ig Nobel Prize winners, joined by nine amused Nobel laureates, take a bow as the ceremony concludes. Photo: Richard Baguley.

KNU

The Nobel Prize in Physics 2010
Andre Geim, Konstantin Novoselov

- The Nobel Prize in Physics 2010
- Andre Geim
- Konstantin Novoselov



Photo: Sergeom, Wikimedia Commons

Andre Geim



Photo: University of Manchester, UK


Konstantin Novoselov


The Nobel Prize in Physics 2010 was awarded jointly to Andre Geim and Konstantin Novoselov *"for groundbreaking experiments regarding the two-dimensional material graphene"*

KNU

Graphene

- graphene – 모든 탄소 동소체의 모체
- graphene에 대한 가까운 미래의 응용





Geim, Nature Mater., **6**, 183 (2007)

KNU

The Nobel Prize in Chemistry 2010
Richard F. Heck, Ei-ichi Negishi, Akira Suzuki

The Nobel Prize in Chemistry 2010
Richard F. Heck
Ei-ichi Negishi
Akira Suzuki



Photo: University of Delaware, USA

Richard F. Heck




Photo: Purdue University, USA

Ei-ichi Negishi



Photo: Hokkaido University, Japan

Akira Suzuki

The Nobel Prize in Chemistry 2010 was awarded jointly to Richard F. Heck, Ei-ichi Negishi and Akira Suzuki *"for palladium-catalyzed cross couplings in organic synthesis"*.

KNU