

Notes on RMSD inference

for Alexander, Fábio, Ruth

Luca  <pgl@portamana.org>

16 June 2021; updated 4 July 2021

1 Overview

We consider a quantity, the Root-Mean-Squared Distance, denoted by y , and a set of quantities called “features”, denoted by x . A value of RMSD y and values for the features x can be associated with each target-ligand pair, which I’ll call a “datapoint”.

Our problem is to infer y_0 given:

- x_0 for a new datapoint,
- known pairs $(\bar{y}, \bar{x}) := ((y_1, x_1), (y_2, x_1), \dots, (y_N, x_N))$ for other N datapoints,
- additional facts and hypotheses H .

Additional hypotheses are unavoidable since we are making an extrapolation.

Our uncertainty about y_0 is expressed by the probability (density)

$$p(y_0 | x_0, \bar{y}, \bar{x}, H), \quad (1)$$

which we want to quantify.

The first question we must ask in approaching this problem is: how much does our uncertainty about y_0 depend on the relative frequencies for previous datapoints (\bar{y}, \bar{x}) ? For example, if y_0 is hand-picked by someone, then knowledge of the frequencies for (\bar{y}, \bar{x}) is irrelevant. But even in such a case it might be that the reversed probability $p(x_0 | y_0, \bar{y}, \bar{x}, H)$ *does* depend on the frequencies of previous datapoints, and it can be used to calculate (1) as

$$p(y_0 | x_0, \bar{y}, \bar{x}, H) = \frac{p(x_0 | y_0, \bar{y}, \bar{x}, H) p(y_0 | H)}{\sum_{y_0} p(x_0 | y_0, \bar{y}, \bar{x}, H) p(y_0 | H)}. \quad (2)$$

The important difference between eqs (1) and (2) is that in the latter we must also quantify $p(y_0 | H)$ using extra-data facts or hypotheses.

In technical terms we are asking whether y_0 is *exchangeable* given x_0 , or vice versa. Using Fisherian (1956 §§ II.4, IV.1) parlance we are asking whether y_0 should be considered as belonging to a *subpopulation* determined by x_0 or vice versa. (Or neither, in which case our study would simply end here, so we won't consider this third possibility.) Another way of seeing our question is whether the “causal connections” mainly go in the direction $x_0 \rightsquigarrow y_0$ or $y_0 \rightsquigarrow x_0$ (cf. Pearl 1988 §§ 2.1.2, 2.2.5).

This question is discussed in a brilliant paper by Lindley & Novick (1981), who show with examples its fundamental importance for making correct inferences. Exchangeability is the basic fact or assumption upon which machine-learning algorithms and the calculations presented here are based. A short intuitive summary of this notion is given in the next section.

In this study we'll study both possibilities mentioned above. At the end we'll compare the results and discuss which assumption makes more sense in various circumstances, for example during testing.

2 Short explanation of exchangeability

Infinite exchangeability is the fact or assumption that the ordering of known datapoints is irrelevant for new inferences. It has the following important consequence. If we knew the infinite-limit long-run relative frequencies of the possible values of past and future datapoints, then the probability of observing a new value would be equal to the long-run relative frequency of that value, for symmetry reasons.

3 Direct case: methodology

For the direct case (1) we consider the RMSD y as a continuous variable, mapped to a log-scale to avoid dealing with finite ranges.

In the direct case the assumption is that y_0 is exchangeable given x_0 , but x_0 is not exchangeable given the known datapoints \bar{x} . This a sensible assumption – a fact indeed – if we are actually *choosing* the specific value of x_0 . The values (\bar{y}, \bar{x}) are exchangeable.

The calculation of $p(y_0 | x_0, \bar{y}, \bar{x}, H)$ is done via the calculation of the joint probability $p(y_0, x_0 | \bar{y}, \bar{x}, H)$ through

$$p(y_0 | x_0, \bar{y}, \bar{x}, H) = \frac{p(y_0, x_0 | \bar{y}, \bar{x}, H)}{p(x_0 | \bar{y}, \bar{x}, H)} \quad (3)$$

with the assumption that the marginal probability for x_0 is independent of the datapoints and equal to unity: $p(x_0 | \bar{y}, \bar{x}, H) = p(x_0 | H) = 1$, which is a sensible assumptions if we are actually *choosing* the specific value of x_0 , as we for example do in testing.

4 Reverse case: methodology

For the reverse case (2) we shall consider a binned RMSD, divided into the three categories “good”, “uncertain”, “bad”.

Bibliography

- (“de X” is listed under D, “van X” under V, and so on, regardless of national conventions.)
- Fisher, R. A. (1956): *Statistical Methods and Scientific Inference*. (Oliver and Boyd, Edinburgh). <https://archive.org/details/in.ernet.dli.2015.134555>.
- Lindley, D. V., Novick, M. R. (1981): *The role of exchangeability in inference*. Ann. Stat. **9**¹, 45–58. DOI:10.1214/aos/1176345331.
- Pearl, J. (1988): *Probabilistic Reasoning in Intelligent Systems: Networks of Plausible Inference*, rev. 2nd pr. (Kaufmann, San Francisco). DOI:10.1016/C2009-0-27609-4.