



**Non Ruminant Research Institute**  
**P.O. Box 25-20117, Tel: + 254 708620095/7, Naivasha**  
**E-mail: [kalro.poultry@kalro.org](mailto:kalro.poultry@kalro.org)**

**Hatching Date:** \_\_\_\_\_

**Batch No.** \_\_\_\_\_

### RECOMMENDED INDIGENOUS CHICKEN VACCINATION PROGRAM

AGE	VACCINE	MODE OF ADMINISTRATION	REMARKS
Day old	Mareks	Subcutaneous	Mainly for commercial hatcheries
	Newcastle+Infectious Bronchitis	Aerosol Spray	
DAY 14	Gumboro (1 <sup>st</sup> dose)	Drinking water	
Day 21	Newcastle disease (1 <sup>st</sup> dose)	Eye drop or Drinking water	
Day 24	Gumboro (2 <sup>nd</sup> dose)	Drinking water	
3 Weeks ( in hot spot areas)	Fowl pox	Wing web stab	
6 Weeks (Other areas)			
8 Weeks	Newcastle disease (2 <sup>nd</sup> dose)	Eye drop or Drinking water	
	Fowl typhoid	Intramuscular injection	
18 Weeks	Newcastle disease (3 <sup>rd</sup> dose at point of lay)	Eye drop or Drinking water	Repeat every 3 months
19 Weeks	De-worming	Drinking water	Repeat every 3 months

Notes:

- NEVER vaccinate sick chicken
- Consult your veterinary/livestock staff for detailed vaccination programs in your area

