

# ASTR615 HW#4

Group 4: ChongChong He & November 18, 2017

## Problem 2: Simulations of Globular Clusters

We perform a series of simulations of globular clusters.

### Mass distribution

We implement in all of our simulations the Kroupa IMF:

$$\phi(m) \propto \begin{cases} m^{-1.3} & (0.08M_{\odot} < m < 0.5M_{\odot}) \\ 0.5 m^{-2.3} & (0.5M_{\odot} < m < 100M_{\odot}) \end{cases} \quad (1)$$

after doing transformation we get

$$m = \begin{cases} -\frac{0.566179}{\sqrt[3]{1.7987 - x} (x^3 - 5.39611x^2 + 9.70599x - 5.81939)} & (0 < x < 0.760707) \\ \frac{0.166558}{(1.00024 - x)^{10/13}} & (0.760707 < x < 1) \end{cases} \quad (2)$$

where  $x$  is uniform random numbers between 0 and 1.

### Initial setup

The cluster is a specially uniformly distributed sphere with a radius of 1. The initial velocities are from a Gaussian distribution with a dispersion correspondent to a virial ratio of  $\alpha \sim 0.4$ . A virial ratio  $\alpha < 0.5$  implies the system is bounded. The velocity dispersion crossing time is  $t \approx 0.08$ , so we use a step size of 0.001, i.e. 80 steps per crossing time. The video of this specific setup is *simulations/cluster03.mp4*.

### Units and Time Scales

The units are:  $[T] = 2.5395 \times 10^{15} \text{ s} \approx 80 \text{ Myr}$ ,  $[L] = 3.08 \times 10^{18} \text{ cm} \approx 1 \text{ pc}$ ,  $[M] = 6.7925 \times 10^{31} \text{ g} \approx 0.03416M_{\odot}$ . In this setup the gravitational constant is unity.

Suppose a cluster of stars from the results of *RAMSES* simulation which has  $\sim 1000$  stars with Salpeter IMF in a box of length  $L = 50$ . If this stellar system is virialized,  $\sigma^2 = M_{\text{tot}}/R$ , from which we obtain  $\sigma \approx 27$ . Here I use a mean mass of  $0.638 M_{\odot} = 20$  (code unit) from Salpeter IMF and  $R = 25$ . Then, the crossing time of dispersion velocity is  $t_{\sigma} = L/\sigma \approx 2$ . If we choose a step size  $\Delta t = t_{\sigma}/50 = 0.04$ , i.e. 50 steps in a course time, then with 5000 steps we are able to simulate 100 course time, which is equal to 16 Gyr. This can be done with a PP n-body code on my laptop.

### Binary Problem

I am trying to conquer the problem of close encounters in our simulation. The basic idea is to check if two stars 1) are close enough to each other and 2)  $K + V < 0$ , every  $n$  steps. If yes, we just replace these two stars with one star at the center of mass. Binaries may further merge into tr binaries, and so on so forth.

## Realization of the Binary Problem

If two stars both with velocity  $\sigma$  are in virial equilibrium, i.e.  $\alpha = K/|W| = 0.5$ , the separation between them would be  $d_{\text{virial}} = m/2\sigma^2 = 0.016$ . However, the typical displacement of a particle in one step is  $d_{\text{step}} = \sigma\Delta t \approx 1$ , much greater than  $d_{\text{virial}}$ . Therefore our simulation is not able to identify binary stars. We need a step size  $\sim 1000$  times smaller to achieve the resolution of binary systems.

## Searching for close encounters

We consider two stars in the center-of-mass frame. We defined the following two parameters:

- The *close-encounter parameter*  $\alpha$  which confirms that they are close enough to each other so that we start to do further test:

$$d_{\text{close}} = |\mathbf{r}_1 - \mathbf{r}_2| < \alpha \cdot d_{\text{virial}}. \quad (3)$$

- The *escape parameter*  $\beta$  which confines the particles in a small region:

$$\frac{K}{|W|} < 1 - \beta^{-1}. \quad (4)$$

This relation gives the largest separation between the two particles at any time,

$$d_{\text{max}} = \beta d_{\text{close}}, \quad (5)$$

ignoring interactions with other particles<sup>1</sup>. When  $d_{\text{max}} \ll d_{\text{step}}$  the two particles may be considered as a binary.

The solution to the binary problem then becomes balancing the typical particle separation  $d_{\text{sepa}} = L/\sqrt[3]{N}$ , the one-step displacement  $d_{\text{step}} = \sigma\Delta t$ , the close-encounter parameter  $\alpha$ , and the escape parameter  $\beta$ .

With  $d_{\text{sepa}} \sim 5$ , we set  $\Delta t = 0.04$ ,  $\alpha = 5$  and  $\beta = 2$ , which imply  $d_{\text{step}} = 1$ ,  $d_{\text{close}} = 0.08$ , and  $d_{\text{max}} = 0.16$ .

---

<sup>1</sup>This relation is obtained by solving equation  $(1 - \beta^{-1})|V_0| + V_0 = 0 + V_1$  and  $V_0 \propto 1/d_0$ ,  $V_1 \propto 1/d_{\text{max}}$ .