Lake Erie Harmful Algal Bloom Bulletin

07 October, 2019, Bulletin 29

Analysis

Microcystis cyanobacteria in the western basin of Lake Erie continues to decrease in concentration and extent. Observed winds (10/3-6) caused mixing that may have reduced any remaining surface concentrations. Recent satellite imagery has been partially obscured by clouds, limiting analysis. Satellite imagery from yesterday (10/6) indicates some chlorophyll is still present within Maumee Bay, but is absent elsewhere within the western basin. The persistent cyanobacteria bloom in Sandusky Bay continues.

Forecasts

Forecast winds (5-12 kn) today through Thursday (10/7-10) will promote mixing of any remaining surface *Microcystis* concentrations. The water temperature has dropped below 68˚F (20˚C) and an increase in bloom concentrations is no longer possible. -Davis, Keeney

Additional Resources

To find a safe place for recreation, visit the Ohio DOH "BeachGuard" site: *http://publicapps.odh.ohio.gov/beachguardpublic/*
Ohio EPA's site on harmful algal blooms: *http://epa.ohio.gov/HAB-Algae*
NOAA's GLERL provides additional HAB data here: *http://www.glerl.noaa.gov/res/HABs_and_Hypoxia*

The images below are "GeoPDF". Please visit https://go.usa.gov/xReTC for instructions on viewing longitude and latitude.






Figure 1. Cyanobacterial Index from modified Copernicus Sentinel 3 data collected 06 October, 2019 at 12:08 EST. Grey indicates clouds or missing data. The estimated threshold for cyanobacteria detection is 20,000 cells/mL.



Wind speed and direction from Marblehead, OH. Blooms mix through the water column at wind speeds greater than 15 knots (or 7.7 m/s).



For more information and to subscribe, please visit the NOAA HAB Foreacst page: https://tidesandcurrents.noaa.gov/hab/lakeerie.html