


7.5 Bipartite Matching

Bipartite Matching

Bipartite matching.

- Input: undirected, **bipartite** graph $G = (L \cup R, E)$.
- $M \subseteq E$ is a **matching** if each node appears in at most one edge in M .
- Max matching: find a max cardinality matching.




max matching

1-1', 2-2', 3-3' 4-4'

Bipartite Matching

Max flow formulation.

- Create digraph $G' = (L \cup R \cup \{s, t\}, E')$.
- Direct all edges from L to R , and assign unit (or infinite) capacity.
- Add source s , and unit capacity edges from s to each node in L .
- Add sink t , and unit capacity edges from each node in R to t .




Bipartite Matching: Proof of Correctness

Theorem. Max cardinality matching in G = value of max flow in G' .

Pf. \leq

- Given max matching M of cardinality k .
- Consider flow f that sends 1 unit along each of k paths.
- f is a flow, and has value at least k . ▪





Bipartite Matching: Proof of Correctness

Theorem. Max cardinality matching in G = value of max flow in G' .

Pf. \geq

- Let f be a max flow in G' of value k .
- Integrality theorem $\Rightarrow k$ is integral and can assume f is 0-1.
- Consider M = set of edges from L to R with $f(e) = 1$.
 - each node in L and R participates in at most one edge in M
 - $|M| = k$: consider cut $(L \cup s, R \cup t)$ ▪



Bipartite Matching: Running Time

Which max flow algorithm to use for bipartite matching?

- Generic augmenting path: $O(m \text{ val}(f^*)) = O(mn)$.
- Capacity scaling: $O(m^2 \log C) = O(m^2)$.
- Shortest augmenting path: $O(m n^{1/2})$.

Which max flow algorithm to use for Non-bipartite matching?

- Structure of non-bipartite graphs is more complicated, but well-understood. [Tutte-Berge, Edmonds-Galai]
- Blossom algorithm: $O(n^4)$. [Edmonds 1965]
- Best known: $O(m n^{1/2})$. [Micali-Vazirani 1980]


7.6 Disjoint Paths

Edge Disjoint Paths

Disjoint path problem. Given a digraph $G = (V, E)$ and two nodes s and t , find the max number of edge-disjoint s - t paths.


Def. Two paths are **edge-disjoint** if they have no edge in common.

Ex: communication networks.



Edge Disjoint Paths

Max flow formulation: assign unit capacity to every edge.




Theorem. Max number edge-disjoint s-t paths equals max flow value.

Pf. \leq

- Suppose there are k edge-disjoint paths P_1, \dots, P_k .
- Set $f(e) = 1$ if e participates in some path P_i ; else set $f(e) = 0$.
- Since paths are edge-disjoint, f is a flow of value k . ▀

Edge Disjoint Paths

Max flow formulation: assign unit capacity to every edge.



Theorem. Max number edge-disjoint s-t paths equals max flow value.

Pf. \geq


- Suppose max flow value is k .
- Integrality theorem \Rightarrow there exists 0-1 flow f of value k .
- Consider edge (s, u) with $f(s, u) = 1$.
 - by conservation, there exists an edge (u, v) with $f(u, v) = 1$
 - continue until reach t , always choosing a new edge
- Produces k (not necessarily simple) edge-disjoint paths. ■

can eliminate cycles to get simple paths if desired

Network Connectivity

Network connectivity. Given a digraph $G = (V, E)$ and two nodes s and t , find min number of edges whose removal disconnects t from s .

Def. A set of edges $F \subseteq E$ **disconnects t from s** if all s - t paths uses at least one edge in F .





Edge Disjoint Paths and Network Connectivity

Theorem. [Menger 1927] The max number of edge-disjoint s - t paths is equal to the min number of edges whose removal disconnects t from s .

Pf. \leq

- Suppose the removal of $F \subseteq E$ disconnects t from s , and $|F| = k$.
- All s - t paths use at least one edge of F . Hence, the number of edge-disjoint paths is at most k . ■





Disjoint Paths and Network Connectivity

Theorem. [Menger 1927] The max number of edge-disjoint s - t paths is equal to the min number of edges whose removal disconnects t from s .

Pf. \geq

- Suppose max number of edge-disjoint paths is k .
- Then max flow value is k .
- Max-flow min-cut \Rightarrow cut (A, B) of capacity k .
- Let F be set of edges going from A to B .
- $|F| = k$ and disconnects t from s . ▪



7.12 Baseball Elimination



Acton Vale



Thetford Mines



Baseball Elimination

Team i	Wins w_i	Losses l_i	To play r_i	Against = r_{ij}			
							
AV 	83	71	8	-	1	6	1
SHA 	80	79	3	1	-	0	2
TM 	78	78	6	6	0	-	0
MTL 	77	82	3	1	2	0	-

Which teams have a chance of finishing the season with most wins?

- Montréal eliminated since it can finish with at most 80 wins, but Acton Vale already has 83.
- $w_i + r_i < w_j \Rightarrow$ team i eliminated.
- Only reason sports writers appear to be aware of.
- Sufficient, but not necessary!

Baseball Elimination

Team i	Wins w_i	Losses l_i	To play r_i	Against = r_{ij}			
							
AV 	83	71	8	-	1	6	1
SHA 	80	79	3	1	-	0	2
TM 	78	78	6	6	0	-	0
MTL 	77	82	3	1	2	0	-

Which teams have a chance of finishing the season with most wins?

- Shawinigan can win 83, but is still eliminated . . .
- If Acton Vale loses a game, then some other team wins one.

Remark. Answer depends not just on **how many** games already won and left to play, but also on **whom** they're against.

Baseball Elimination


Baseball elimination problem.

- Set of teams S .
- Distinguished team $x \in S$.
- Team x has won w_x games already.
- Teams x and y play each other r_{xy} additional times.
- Is there any outcome of the remaining games in which team x finishes with the most (or tied for the most) wins?

Baseball Elimination: Max Flow Formulation

Can team 3 finish with most wins?


- Assume team 3 wins all remaining games $\Rightarrow w_3 + r_3$ wins.
- Divide remaining games so that all teams have $\leq w_3 + r_3$ wins.













Baseball Elimination: Max Flow Formulation

Theorem. Team 3 is not eliminated iff max flow saturates all edges leaving source s .

- Integrality theorem \Rightarrow each remaining game between x and y added to number of wins for team x or team y .
- Capacity on (x, t) edges ensure no team wins too many games.



Baseball Elimination: Explanation for Sports Writers

Team i	Wins w_i	Losses l_i	To play r_i	Against = r_{ij}				
								
AV 	75	59	28	-	3	8	7	3
Sha 	71	63	28	3	-	2	7	4
TM 	69	66	27	8	2	-	0	0
She 	63	72	27	7	7	0	-	-
Mtl 	49	86	27	3	4	0	0	-

Which teams have a chance of finishing the season with most wins?

- Montréal could finish season with $49 + 27 = 76$ wins.

Certificate of elimination. $R = \{AV, Sha, TM, She\}$

- Have already won $w(R) = 278$ games.
- Must win at least $r(R) = 27$ more (loosing all non- R games).
- Average team in R wins at least $305/4 > 76$ games.

Baseball Elimination: Explanation for Sports Writers

Certificate of elimination.

$$T \subseteq S, \quad w(T) := \overbrace{\sum_{i \in T} w_i}^{\text{\# wins}}, \quad r(T) := \overbrace{\sum_{\{x,y\} \in T} r_{xy/2}}^{\text{\# remaining games}},$$

LB on avg # games won
If $\frac{w(T) + r(T)}{|T|} > w_z + r_z$ then z is **eliminated** (by subset T).

Theorem. [Hoffman-Rivlin 1967] Team z is eliminated iff there exists a subset T^* that eliminates z .

Proof idea. Let $T^* =$ team nodes in A (on source side) of min cut (A,B) .


Baseball Elimination: Explanation for Sports Writers

Pf of theorem.

- Use max flow formulation, and consider min cut (A, B) .
- Define T^* = team nodes in A (on source side of min cut).
- Observe $x-y \in A$ iff both $x \in T^*$ and $y \in T^*$.
- $r(S - \{z\}) > \text{cap}(A, B)$

$$\begin{aligned}
 &= \overbrace{r(S - \{z\}) - r(T^*)}^{\text{capacity of game edges leaving } A} + \overbrace{\sum_{x \in T^*} (w_z + r_z - w_x)}^{\text{capacity of team edges leaving } A} \\
 &= r(S - \{z\}) - r(T^*) - w(T^*) + |T^*|(w_z + r_z)
 \end{aligned}$$

- Rearranging terms: $w_z + r_z < \frac{w(T^*) + r(T^*)}{|T^*|}$ ■



7.10 Image Segmentation

Image Segmentation

Image segmentation.


- Central problem in image processing.
- Divide image into coherent regions.

Ex: Three people standing in front of complex background scene.
Identify each person as a coherent object.

Image Segmentation

Foreground / background segmentation.

- Label each pixel in picture as belonging to foreground or background.
- V = set of pixels, E = pairs of neighbouring pixels.
- $a_i \geq 0$ is likelihood pixel i in foreground.
- $b_i \geq 0$ is likelihood pixel i in background.
- $p_{ij} \geq 0$ is separation penalty for labelling one of i and j as foreground, and the other as background.



Goals.

- Accuracy: if $a_i > b_i$ in isolation, prefer to label i in foreground.
- Smoothness: if many neighbours of i are labeled foreground, we should be inclined to label i as foreground.
- Find partition (A, B) that maximizes:

$$\sum_{i \in A} a_i + \sum_{j \in B} b_j - \sum_{\substack{(i,j) \in E \\ |A \cap \{i,j\}| = 1}} p_{ij}$$

\nearrow
foreground

\nwarrow
background

Image Segmentation

Formulate as min cut problem.

- Maximization.
- No source or sink.
- Undirected graph.

Turn into minimization problem.

- Maximizing
$$\sum_{i \in A} a_i + \sum_{j \in B} b_j - \sum_{\substack{(i,j) \in E \\ |A \cap \{i,j\}| = 1}} p_{ij}$$

is equivalent to minimizing
$$\underbrace{\left(\sum_{i \in V} a_i + \sum_{j \in V} b_j \right)}_{\text{a constant}} - \sum_{i \in A} a_i - \sum_{j \in B} b_j + \sum_{\substack{(i,j) \in E \\ |A \cap \{i,j\}| = 1}} p_{ij}$$

- or alternatively
$$\sum_{j \in B} a_j + \sum_{i \in A} b_i + \sum_{\substack{(i,j) \in E \\ |A \cap \{i,j\}| = 1}} p_{ij}$$

Image Segmentation

Formulate as min cut problem.

- $G' = (V', E')$.
- Add source s to correspond to foreground; add sink t to correspond to background
- Use two anti-parallel edges instead of undirected edge.





Image Segmentation


Consider min cut (A, B) in G' .

- A = foreground.

$$cap(A, B) = \sum_{j \in B} a_j + \sum_{i \in A} b_i + \sum_{\substack{(i,j) \in E \\ i \in A, j \in B}} p_{ij}$$

if i and j on different sides,
 p_{ij} counted exactly once

- Precisely the quantity we want to minimize.



7.11 Project Selection

Project Selection

Projects with prerequisites.

can be positive or negative



- Set P of possible projects. Project v has associated revenue p_v .
 - some projects generate money: create interactive e-commerce interface, redesign web page
 - others cost money: upgrade computers, get site license
- Set of prerequisites E . If $(v, w) \in E$, can't do project v and unless also do project w .
- A subset of projects $A \subseteq P$ is **feasible** if the prerequisite of every project in A also belongs to A .


Project selection. Choose a feasible subset of projects to maximize revenue:

$$\sum_{v \in A} p_v$$


Project Selection: Prerequisite Graph

Prerequisite graph.

- Include an edge from v to w if can't do v without also doing w .
- $\{v, w, x\}$ is feasible subset of projects.
- $\{v, x\}$ is infeasible subset of projects.



feasible




infeasible

Project Selection: Min Cut Formulation

Min cut formulation.


- Assign capacity ∞ to all prerequisite edge.
- Add edge (s, v) with capacity p_v if $p_v > 0$.
- Add edge (v, t) with capacity $-p_v$ if $p_v < 0$.
- For notational convenience, define $p_s = p_t = 0$.



Project Selection: Min Cut Formulation

Claim. (A, B) is min cut iff $A - \{s\}$ is optimal set of projects.


- Infinite capacity edges ensure $A - \{s\}$ is feasible.
- Max revenue because:
$$\begin{aligned} \text{cap}(A, B) &= \sum_{v \in B: p_v > 0} p_v + \sum_{v \in A: p_v < 0} (-p_v) \\ &= \underbrace{\sum_{v: p_v > 0} p_v}_{\text{constant}} - \sum_{v \in A} p_v \end{aligned}$$



Example: Open Pit Mining

Open-pit mining. (studied since early 1960s)

- Blocks of earth are extracted from surface to retrieve ore.
- Each block v has net value $p_v = \text{value of ore} - \text{processing cost}$.
- Can't remove block v before w or x .



7.7 Extensions to Max Flow

Circulation with Demands

Circulation with demands.

- Directed graph $G = (V, E)$.
- Edge capacities $c(e)$, $e \in E$.
- Node supply and demands $d(v)$, $v \in V$.



demand if $d(v) > 0$; supply if $d(v) < 0$; transshipment if $d(v) = 0$

Def. A **circulation** is a function f that satisfies:

- For each $e \in E$: $0 \leq f(e) \leq c(e)$ (capacity)
- For each $v \in V$: $\sum_{e \text{ in to } v} f(e) - \sum_{e \text{ out of } v} f(e) = d(v)$ (conservation)


Circulation problem: given (G, c, d) , does there exist a circulation?

Circulation with Demands

Necessary condition: sum of supplies = sum of demands.


$$\sum_{v: d(v) > 0} d(v) = \sum_{v: d(v) < 0} -d(v) =: D$$

Pf. Sum conservation constraints for every demand node v .



Circulation with Demands


Max flow formulation.



Circulation with Demands

Max flow formulation.

- Add new source s and sink t .
- For each v with $d(v) < 0$, add edge (s, v) with capacity $-d(v)$.
- For each v with $d(v) > 0$, add edge (v, t) with capacity $d(v)$.
- Claim: G has circulation iff G' has max flow of value D .



Circulation with Demands

Integrality theorem. If all capacities and demands are integers, and there exists a circulation, then there exists one that is integer-valued.

Pf. Follows from max flow formulation and integrality theorem for max flow.

Characterization. Given (G, c, d) , there does **not** exist a circulation iff there exists a node partition (A, B) such that $\sum_{v \in B} d(v) > \text{cap}(A, B)$



Pf idea. Look at min cut in G' .

demand by nodes in B exceeds supply of nodes in B plus max capacity of edges going from A to B

Circulation with Demands and Lower Bounds

Feasible circulation.

- Directed graph $G = (V, E)$.
- Edge capacities $c(e)$ and lower bounds $\ell(e)$, $e \in E$.
- Node supply and demands $d(v)$, $v \in V$.

Def. A **circulation** is a function that satisfies:



- For each $e \in E$: $\ell(e) \leq f(e) \leq c(e)$ (capacity)
- For each $v \in V$: $\sum_{e \text{ in to } v} f(e) - \sum_{e \text{ out of } v} f(e) = d(v)$ (conservation)

Circulation problem with lower bounds. Given (G, ℓ, c, d) , does there exists a circulation?

Circulation with Demands and Lower Bounds

Idea. Model lower bounds with demands.

- Send $\ell(e)$ units of flow along edge e .
- Update demands of both endpoints.



Theorem. There exists a circulation in G iff there exists a circulation in G' . If all demands, capacities, and lower bounds in G are integers, then there is a circulation in G that is integer-valued.

Pf sketch. $f(e)$ is a circulation in G iff $f'(e) = f(e) - \ell(e)$ is a circulation in G' .

7.8 Survey Design

Survey Design

Survey design.

- Design survey asking n_c consumers about n_p products.
- Can only survey consumer i about a product j if he owns it.
- Ask consumer i between c_i and c'_i questions.
- Ask between p_j and p'_j consumers about product j .

Goal. Design a survey that meets these specs, if possible.

Survey Design

Algorithm. Formulate as a circulation problem with lower bounds.

- Include an edge (i, j) if customer i owns product j .
- Integer circulation \Leftrightarrow feasible survey design.
- (t, s) makes sure problem has a circulation.

