



International Journal of Homoeopathic Sciences

E-ISSN: 2616-4493
P-ISSN: 2616-4485
www.homoeopathicjournal.com
IJHS 2025; 9(2): 71-73
Received: 23-01-2025
Accepted: 29-02-2025

Dr. Navroop Bhatia
PG Scholar, Department of
Materia Medica, Dr. MPK
Homoeopathic Medical
College, Hospital & Research
Centre, Jaipur, Rajasthan,
India

Dr. Sachin Kumar Yadav
PG Scholar, Department of
Materia Medica, Dr. MPK
Homoeopathic Medical
College, Hospital & Research
Centre, Jaipur, Rajasthan,
India

Dr. Suraj Bhan Singh Rathore
PG Scholar, Department of
Materia Medica, Dr. MPK
Homoeopathic Medical
College, Hospital & Research
Centre, Jaipur, Rajasthan,
India

Dr. Sonia Tuteja
Professor, Dr. MPK
Homoeopathic Medical
College, Hospital & Research
Centre, Jaipur, Rajasthan,
India

Corresponding Author:
Dr. Navroop Bhatia
PG Scholar, Department of
Materia Medica, Dr. MPK
Homoeopathic Medical
College, Hospital & Research
Centre, Jaipur, Rajasthan,
India

LM potency explained: The future of homeopathic healing

Dr. Navroop Bhatia, Dr. Sachin Kumar Yadav, Dr. Suraj Bhan Singh Rathore and Dr. Sonia Tuteja

DOI: <https://www.doi.org/10.33545/26164485.2025.v9.i2.B.1477>

Abstract

LM potency also known as fifty-millesimal potency or Q potency. Dr. Samuel Hahnemann aimed to achieve a faster, gentler, and less aggravating method of cure, leading to his development of the potentization process. Initially, he introduced the centesimal scale, but he was dissatisfied with its results due to frequent aggravations and slower recovery. LM potencies, being Hahnemann's final refinement, are highly gentle, effective in deep pathological cases, and suitable for hypersensitive patients. They allow for frequent repetition without severe aggravations, significantly reducing the treatment duration. Their flexibility and precision make them an essential tool in homeopathic practice, offering a more refined and efficient approach to healing.

Keywords: LM potency, fifty millesimal potency, Q potency, deep pathological cases

Introduction

Hahnemann introduced the fifty-millesimal potencies, which are referred to by different names across various regions and even within the same country. The terminology used for these potencies varies, with the most commonly recognized names being "LM potency" and "Q potency". Similarly, the symbols used to represent these potencies also differ based on location and practice^[1].

The term "LM potency" is widely used in the Indian subcontinent and Latin America, gaining popularity through the work of Rudolf Flury from Switzerland. It derives from Roman numerals, where 'L' represents 50 and 'M' represents 1000. However, this name is technically inaccurate, as the combination "LM" numerically signifies 950 rather than 50,000^[1].

Q potency The letter 'Q' represents "quinquaginta milia" in Latin, meaning 50,000 ("quinquaginta" for fifty and "milia" for thousand), making it the technically accurate term for fifty-millesimal potencies. However, in the Indian subcontinent, 'Q' is mistakenly used to denote mother tinctures, whereas the correct symbol for mother tinctures is 'ϕ' (phi)^[1].

Dr. Hahnemann aimed to accelerate the healing process in a gentler manner with minimal aggravation and no harm to the patient. Through his experiments, he developed the potentization process and initially introduced the centesimal scale. However, he found its results unsatisfactory, as it often led to stronger aggravations compared to the LM scale. He also observed that the LM scale facilitated a much faster recovery. Unfortunately, due to the delayed publication of the 6th edition of the Organon, many homeopathic practitioners remained unaware of LM potencies. By that time, most homeopaths had already adopted centesimal potencies, as they were yielding effective results^[2].

LM potency is the refined result of Hahnemann's extensive experience and deep understanding, as documented in the 6th edition of the Organon. These potencies are remarkably gentle, causing minimal to no violent aggravations. They are particularly effective in treating deep pathological conditions, often reducing the treatment duration by half or even less. Additionally, LM potencies are well-suited for hypersensitive patients, allowing for safe and effective treatment. The selection of potency is straightforward, and frequent repetition can be done when necessary without adverse effects^[3].

The origins of LM potency: Hahnemann's final innovation ^[4].

Samuel Hahnemann's development of LM (Q) potencies remained largely hidden for years due to secrecy and controversy. Following his death, his widow, Mélanie d'Hervilly, tightly controlled his manuscripts and withheld the release of the 6th edition of the Organon, which first introduced LM potencies. She set an unreasonably high price for the manuscript, preventing its publication during her lifetime. In 1856, a German journal revealed hints about Hahnemann's new potentization method, which infuriated Mélanie. Despite efforts by Clemens von Bönninghausen, one of Hahnemann's closest associates, she refused to release the manuscript. The 6th edition of the Organon was eventually published in 1920, but by then, homeopathy was already well-established with centesimal (C) potencies. It was only in 1950, through the work of Dr. Charles Pahud

and Dr. Pierre Schmidt, that LM potencies gained recognition and active exploration. In his final years, Hahnemann extensively used LM potencies, prescribing them in at least 1,836 cases. Initially, he experimented with lower LM potencies (Q1-Q10), comparing their effects with Centesimal potencies before fully adopting the 50-millesimal scale. Later, researchers such as Dr. Rudolf Flury, Ubiratan Adler, and Dr. RP Patel conducted studies on LM potencies, emphasizing their flexibility, potency, and minimal aggravations when used correctly. Unlike Centesimal potencies, LM potencies require frequent repetition and precise adjustment of succussions. Some experts caution against administering high LM potencies in dry doses, as they may cause strong aggravations.

Today, LM potencies are recognized as a gentler yet potent alternative to centesimal potencies, offering greater control and individualized dosing for chronic and sensitive patients.

Table 1: Understanding LM potency: How it's made ^[5].

	Remedy in Plant Form	Liquid Form	Mineral / Solid Form
1c	Triturate the entire plant	Begin with the liquid substance such as sepia ink, venoms, petroleum, etc. Add it to lactose in a ratio of 1 part liquid to 100 parts lactose for example, 1 ml of liquid to 99 g of lactose, then triturate.	Begin with the powdered mineral. Mix 1 part of it with 99 parts of lactose and triturate the mixture thoroughly.
2c	Triturate 1 part of 1c preparation with 99 parts of lactose.		
3c	Triturate 1 part of 2c preparation with 99 parts of lactose.		
LM1	Take 1 grain (0.062 grams) of the 3C potency and add 500 drops of a solution made from 1 part alcohol and 4 parts distilled water (i.e., 100 drops of alcohol and 400 drops of water). Without succussing this mixture, take one drop and add it to 2 ml of alcohol. Succuss the mixture 100 times. This creates your LM1 stock solution. Then, place one drop from the LM1 stock onto 500 poppy seed-sized pellets (#10 pellets). These pellets become your LM1 dose, which you'll use to prepare the patient's remedy bottle.		
LM2	Dissolve one LM1 pellet in a single drop of water. Then add 99 drops of alcohol to the solution and succuss it 100 times. This creates your LM2 stock solution. Place one drop from the LM2 stock onto 500 #10 pellets. These pellets are your LM2 dose.		
LM3	Dissolve one LM2 pellet in a drop of water. Add 99 drops of alcohol to the mixture and succuss it 100 times. This forms your LM3 stock solution. Then, dispense one drop from the LM3 stock onto 500 #10 pellets. These are your LM3 pellets.		

How to administer medicine in LM potency ^[6].

- Thoroughly cleanse a 4-ounce phial and use a fresh cork.
- Fill three-fourths of the bottle with distilled or purified drinking water.
- Add 15-20 drops of rectified spirit to the solution to ensure preservation.
- Crush one No. 10 globule of 0/1, 0/2, or 0/3 (whichever medicine is selected) with 2-3 grains of milk sugar, then add the mixture to the bottle.
- Divide the phial into seven doses. The medicinal solution is now prepared for dispensing.
- Before consumption, each of the seven doses should be succussed 8, 10, or 12 times, depending on the patient's susceptibility (8 times for highly sensitive patients and 12 times for those least sensitive).
- After succussion, place one dose of the medicine into a glass containing approximately 4 ounces of pure water, and stir thoroughly using a teaspoon.
- From the prepared solution, instruct the patient to take only 1 to 2 teaspoons of the dose and discard the remaining liquid. Repeat this method for each subsequent dose.

If any aggravation occurs after taking the remedy as described, discontinue further doses. The symptoms will ease on their own shortly. Once the aggravation has settled,

the same remedy can be taken again, but this time diluted further — using a second, third, or even fourth glass prepared in the same manner. This approach helps to manage and reduce the aggravation.

Dosage

- Only one single, simple remedy should be given at a time
- Each dose should be kept as minimal as possible, except in the following situations:
 1. A recent outbreak of skin itch.
 2. An untreated chancre on the genitals, lips, or other areas.
 3. The presence of fig warts.

In these three situations, the recommended remedy should be administered in larger doses and at higher potencies on a daily basis.

- The remedy should be administered in multiple doses rather than just one. A single No. 10 pellet infused with the medicine only comes into contact with a limited number of nerve fibers. However, when the pellet is crushed, blended with a bottle of lactose powder (sugar of milk), and then dissolved in water, it becomes a much more potent preparation.
- Remedies prepared on the 50 millesimal scale should be administered exclusively in liquid form

Potency

- As stated in § 270 of the 6th edition of the Organon, Dr. Hahnemann advises using potencies ranging from 0/1 to 0/30. Most patients experience significant healing within the 0/10 range, making higher potencies generally unnecessary.
- Begin treatment with the lowest level of dynamization (such as 0/1 to 0/3) and gradually move on to higher potencies over time.

Time for Administration of Remedy

- In cases of fever, the best time to give the remedy is immediately after the fever paroxysm has ended.
- Many acute and chronic illnesses have specific times during their course when symptoms worsen. Medication should not be given during these aggravation periods. For example, in syphilitic patients, symptoms typically intensify at night, so it's best to avoid administering the remedy during nighttime hours.
- Many remedies have a distinct time when they cause medicinal aggravation. This period is not suitable for administering those medicines. For e.g.,
- Arsenicum album tends to aggravate between 12 a.m. and 2 a.m., as well as from 12 p.m. to 2 p.m.
- Kali group remedies show aggravation typically between 3 a.m. and 5 a.m.
- Lycopodium symptoms worsen from 4 p.m. to 8 p.m.
- Natrium muriaticum aggravation occurs around 10 a.m. to 11 a.m.
- Sulphur usually aggravates between 10 a.m. and 2 a.m.
- When treating menstrual disorders, the most effective time to prescribe a remedy is after the menstrual period has ended.

Discussion

Despite being Hahnemann's preferred system, fifty millesimal potencies have been slowly accepted in general homoeopathic practice. The delay of almost eighty years before publication of the 6th edition in 1920 and the subsequent controversy over its authenticity certainly contributed to the situation ^[7]. LM potencies hold great potential in modern homeopathic practice, offering flexibility, precision, and reduced risk of aggravations when administered correctly. With further study and training, they can become a valuable tool for improving treatment outcomes.

Conclusion

LM potencies represent a refined and powerful advancement in homeopathy, offering greater flexibility, gentleness, and control in treatment. Despite initial resistance, they are now gaining wider acceptance. With further research and education, LM potencies have the potential to revolutionize homeopathic practice, making treatments more effective and adaptable to individual patient needs.

References

1. Pareek A, Pareek N, Nayak C. Fifty millesimal potencies—an overview. *Reverse Repertory*. [date unknown];29.
2. Sumanth BL. Effect of LM potency in homoeopathic general practice [dissertation]. Bengaluru: Rajiv Gandhi University of Health Sciences (India); [year unknown].

3. Hussain DT. Hidden treasure of Organon-LM potency. *Allg Homöopath Ztg*. 2017 Mar;262(2):CM18-03.
4. Bala R, Srivastava A. A review on history of LM potency: Tracing its roots in the past. *Int J Hum Sci*. 2019;3(3):26-31.
5. De Schepper L. *Hahnemannian textbook of classical homoeopathy for the professional*. 4th impression. New Delhi: B Jain Publishers; 2010. p. 88-91.
6. Banerjee DD. *Textbook of homoeopathic pharmacy*. New Delhi: B Jain Publishers; 2002.
7. Baker D. Use of fifty millesimal potencies in contemporary homoeopathy. *J Aust Tradit Med Soc*. 2003 Jun 1;9(2):[page numbers not provided].

How to Cite This Article

Bhatia N, Yadav SK, Rathore SBS, Tuteja S. LM potency explained: The future of homeopathic healing. *International Journal of Homoeopathic Sciences*. 2025; 9(2): 71-73.

Creative Commons (CC) License

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International (CC BY-NC-SA 4.0) License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.