
### Tell us whether you accept cookies

We use <u>cookies to collect information</u> about how you use GOV.UK. We use this information to make the website work as well as possible and improve government services.

Accept all cookies

Set cookie preferences



Home > COVID-19: infection prevention and control (IPC)



#### Guidance

# Explanation of the updates to infection prevention and control guidance

Updated 4 April 2020

#### Contents

- 1. Process for updating the guidance (as published on 2 April 2020)
- 2. Main changes to the guidance

# 1. Process for updating the guidance (as published on 2 April 2020)

Public Health England (PHE) has worked with NHS England and NHS Improvement, and the Devolved Administrations to review the UK's infection prevention and control recommendations for COVID-19.

PHE also consulted with the Royal College of Nursing, other Royal Colleges and professional societies.

Having assessed the available evidence and feedback received from guidance users, we have updated sections to improve the guidance and ensure that we continue to make recommendations that will help prevent the spread of COVID-

# 2. Main changes to the guidance

- three new tables have been added to clearly explain the PPE required for different common clinical scenarios one for hospitals, one for primary care, outpatient and community care and one for ambulance and first responders
- an additional, fourth table describes when to use PPE for all patient encounters (not just patients with suspected or confirmed COVID-19) at a time when there is sustained community transmission of COVID-19, as is currently occurring in the UK, and the likelihood of any patient having coronavirus infection is raised
- the guidance explains that in some circumstances PPE can be worn for an entire session (such as a ward round) and does not need to be changed between each patient
- patient contact is now defined as being within 2 metres (rather than within 1 metre) of a patient, which is more precautionary and is consistent with the distancing recommendations used elsewhere
- hand-washing advice has been updated to include washing of forearms, when forearms have been exposed or may
  have been exposed to respiratory droplets or other body fluids
- FFP2 and N95 respirators may be used for some aerosol-generating procedures if FFP3 respirators are not available. FFP3 respirators offer a slightly higher level of protection than FFP2 respirators
- advice on re-usable PPE components, including the need to refer to manufacturer's guidance on decontamination
- · recommendations about the use of facemasks by patients
- · general formatting improvements to make it easier to access different sections of the guidance

Is this page
useful? Yes No

Is there anything wrong with this
page?

#### **Coronavirus (COVID-19)**

Coronavirus (COVID-19): what you need to do

## **Transition period**

Transition period: check how to get ready

#### Services and information

**Benefits** 

Births, deaths, marriages and care

Business and self-employed

Childcare and parenting

Citizenship and living in the UK

Crime, justice and the law

Disabled people

**Driving and transport** 

Education and learning

**Employing people** 

Environment and countryside

Housing and local services

Money and tax

Passports, travel and living abroad

Visas and immigration

Working, jobs and pensions

## **Departments and policy**

How government works

**Departments** 

Worldwide

Services

Guidance and regulation

News and communications

Research and statistics

Policy papers and consultations

Transparency and freedom of information releases

<u>Help Privacy Cookies Contact Accessibility statement Terms and conditions Rhestr o Wasanaethau Cymraeg</u>

Built by the <u>Government Digital Service</u>

# **OGL**

All content is available under the Open Government Licence v3.0, except where otherwise stated



© Crown copyright