Last Day: Matrix Multiplication

$$\begin{bmatrix} 2 & 3 \\ \hline 1 & 5 \end{bmatrix} \begin{bmatrix} -1 \\ 2 \end{bmatrix} = \begin{bmatrix} 2(-1) + 3(2) \\ \hline 1(-1) + 5(2) \end{bmatrix} = \begin{bmatrix} 4 \\ 9 \end{bmatrix}$$

square 2 2x1 2x1 col. vector = 3 => result is outer dimensions: 2x1

match

$$\begin{bmatrix} 2 & 3 \\ 1 & 5 \end{bmatrix} \begin{bmatrix} -1 \\ 2 \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \end{bmatrix} (-1) + \begin{bmatrix} 3 \\ 5 \end{bmatrix} (2)$$
as a combination of columns.

$$= \left[2(-1) + 3(2) \right] = \left[4 \right],$$

Orda #1

Let's make an order motrix for ports!

3 Part 1 5 Part 2

Let's make a list of cost of parts:

Now: Multiply

$$Am. \begin{cases} 2 & 3 \\ 6 & 4 \end{cases}$$

$$Am. \begin{cases} 2 & 3 \\ 1 & 4 \end{cases} \begin{cases} 3 & p_1 \\ 5 & p_2 \end{cases} = \begin{cases} 2(3) + 3(5) \\ 1(3) + 4(5) \end{cases} = \begin{cases} 21 \\ 23 \end{cases} Eb.$$

C N

(cost) (# orders) (comount paid)

ic (CN = P)

What if Ordin size unknown & Cost & Paid amount one known? Can we find N?

 $\begin{bmatrix} 2 & 3 \\ 1 & 4 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 21 \\ 23 \end{bmatrix}$ (3) (3) (4) (5) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7) (7

This works for any linear system! 2x + fy - 2 = 2 x + 7y + 2 = 3-3 x + 2 x = 5 $\begin{bmatrix} 1 & 7 & 1 \\ 1 & 7 & 1 \\ -3 & 2 & 0 \end{bmatrix} \begin{bmatrix} \chi \\ \chi \\ \chi \end{bmatrix} = \begin{bmatrix} 2 & 7 \\ 3 & 5 \end{bmatrix}$ coefficient matrix Constant > Variables

Goal: Invent division! => First, invent exponents,

A o = ? For all A, AAO = A = A

nxn sized

nxn sized

nxn hxn
hxn "Identity matrix" (acts like 1 but widen!)

$$I_{3} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \quad I_{2} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$I_{h} = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \end{bmatrix} \quad A = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \end{bmatrix} \quad A = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \end{bmatrix} \quad A = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \end{bmatrix} \quad A = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \end{bmatrix} \quad A = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \end{bmatrix} \quad A = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \end{bmatrix} \quad A = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \end{bmatrix} \quad A = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \end{bmatrix} \quad A = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \end{bmatrix} \quad A = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \end{bmatrix} \quad A = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \end{bmatrix} \quad A = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1$$

Is A unique for a given A? Say BA = AB = I >> Bis an invale of A & CA = AC = I => C is anothy invoce of A then B = BI = BAC = IC = C $\Rightarrow B = C$ => invare is unique (if exist!) A -1 = inverce of A", defined as A -1 H = AA-1 = I If it exists => "A is invertible" or (non-singular

If it exists => "A is invertible" or (non-singular

If it does not exist => "A is not invertible

or Singular

Proportion of Inverse

$$(A^{-1})^{-1} = A$$

Proof
$$(kA)^{-1} \cdot kA = I = kA \cdot (kA)^{-1}$$

doe, $\frac{1}{k}A^{-1} \cdot f.i$?

Cleck $(\frac{1}{k}A^{-1})(kA) = \frac{k}{k}A^{-1}A = 1I = I$
 $(kA)(\frac{1}{k}A^{-1}) = \frac{h}{k}AA^{-1} = I$

Cleck
$$\left(\frac{1}{k}A^{-1}\right)(kA) = \frac{k}{k}A^{-1}A = 4I = I$$

 $(kA)(\frac{1}{k}A^{-1}) = \frac{k}{k}AA^{-1} = I$

4)
$$(A^{+})^{-1} = (A^{-1})^{+}$$

5) $(A + B)^{-1} = Horror!$

Sample Problem If A is an nxn matrix
find an expression for A-1 if

Solution

$$3A^{3} + 2A^{2} + 7A = -2I$$

 $-\frac{3}{5}A^{3} + A^{2} - \frac{1}{5}A = I$

Why the I' above?

Consider: Say
$$A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$$

Then $A^2 + 2A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$

Then $A^2 + 2A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 2 & 4 \end{bmatrix} + 2 \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$

If we don't use the I here, this object is not a matrix!

$$\begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} + 2 \quad \text{is undefined}$$

Then $A^2 + 2A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} = \begin{bmatrix} 1 & 2 \\ 2 & 4 \end{bmatrix} + 2 \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$