MATHEMATICS 1LS3 TEST 1

Day Class
Duration of Examination: 60 minutes
McMaster University, 7 October 2015

CI1

E. Clements, G. Dragomir, M. Lovrić

First name (PLEASE PRINT):	
Family name (PLEASE PRINT):	
Student No.:	

THIS TEST HAS 8 PAGES AND 7 QUESTIONS. YOU ARE RESPONSIBLE FOR ENSURING THAT YOUR COPY OF THE PAPER IS COMPLETE.

Total number of points is 40. Marks are indicated next to the problem number. Any non-graphing calculator is allowed.

USE PEN TO WRITE YOUR TEST. IF YOU USE A PENCIL YOUR TEST WILL NOT BE ACCEPTED FOR REMARKING (IF NEEDED).

You must show work to receive full credit.

Problem	Points	Mark
1	4	
2	6	
3	8	
4	4	
5	6	
6	5	
7	7	
TOTAL	40	

1. Multiple choice questions: circle ONE answer. No justification is needed.

- (a)[2] Identify all correct statements about the function $f(x) = \frac{x^2 1}{x^2 + 1}$.
 - (I) f(x) is continuous at x = 0.
 - (II) x = -1 is a vertical asymptote of the graph of f(x).
 - (III) $\mathbf{x} = 1$ is a horizontal asymptote of the graph of f(x)
 - (A) none
- (B) I only
- (C) II Only
- (D) III only

- (E) I and II
- (F) I and III
- (G) II and III
- (H) ALL

(b)[2] The average half-life of acetaminophen (active ingredient in tylenol) is 2.5 hours. Assume that a patient is given a dose of 1000 mg of acetaminophen. Identify all correct statements.

- (I) After 5 hours, 250 mg of acetaminophen is left unabsorbed in patient's body.
- (II) After 2 hours, 450 mg of acetaminophen is left unabsorbed in patient's body.
- (III) After 10 hours, less than 100 mg of acetaminophen is left unabsorbed in patient's body.
 - (A) none
- (B) I only
- (C) II Only
- (D) III only

- (E) I and II
- (F) I and III
- (G) II and III
- (H) ALL

2. Identify each statement as true or false, or yes or no (circle your choice). You do not need to justify your answer.

(a)[2] The fact that $\lim_{x\to 2} f(x) = 3$ implies that f(x) is continuous at x=2.

TRUE

OR

FALSE

(b)[2] The formula H = Mx + 4, where M is a constant, represents a proportional relationship between H and x.

TRUE

OR

FALSE

(c)[2] The average density of a human jaw bone is 1.07 oz/in^3 , which is equivalent to approximately 1.85 g/cm^3 . [Conversion factors: 1 oz = 28.35 g, 1 in = 2.54 cm]

TRUE

OR

FALSE

Questions 3-7: You must show work to receive full credit.

3. Consider the function

$$f(x) = \begin{cases} \frac{x^3 - x}{x - 1} & \text{if } x < 1\\ 2x & \text{if } x \ge 1 \end{cases}$$

(a)[4] Find $\lim_{x\to 1} f(x)$.

(b)[2] Is f(x) continuous at x = 1? Explain why or why not.

(c)[2] Is f(x) continuous at x = -1? Explain why or why not.

Name:		
Student No.:		

4. The following excerpt is taken from *The laminar cortex model: a new continuum cortex model incorporating laminar architecture.* J. Du, V. Vegh, and D.C. Reutens. PLoS Computational Biology. 8.10 (Oct. 2012).

the average of membrane potentials of neurons in the element, that is

$$V = \frac{N_{\rm e}V_{\rm e} + N_{\rm i}V_{\rm i}}{N_{\rm e} + N_{\rm i}}$$

where N_e , N_i are the numbers of excitatory and inhibitory neurons and V_e and V_i are the (average) membrane potentials of excitatory and inhibitory neuron populations respectively.

View V as a function of N_i .

By reading the text, we learn that V is the average of membrane potentials and N_i is the number of inhibitory neurons.

(a)[1] State (in one sentence) what question is answered by finding the inverse function of V.

(b)[3] Find a formula for the inverse function of V.

Name:	
Student No.:	

5. (a)[3] Consider the formula for human population growth

$$P(t) = 4.43 \left(\frac{\pi}{2} + \arctan \frac{t - 2007}{42}\right)$$

where t is a calendar year and P(t) is in billions. Find the range of P(t). Based on it, state the maximum world population predicted by this model.

(b)[3] A population of river sharks (freshwater sharks) in New Zealand changes periodically with a period of 12 months. In January, it reaches a maximum of 12,600, and in July it reaches a minimum of 5,800. By selecting an appropriate trigonometric function, find a formula which describes how the population of river sharks changes with time.

6. The survival rate (i.e., percent) S(D) of clonogenic cells (cancer cells) is modelled by


$$S(D) = e^{-0.6D}$$

where $D \ge 0$ represents the applied radiation dose (measured in grays, Gy).

(a)[1] Assume that the dose D=5 Gy is applied to a cancer. What percent of cancer cells is going to survive this treatment? Nearest integer;

(b)[2] Sketch the semilog graph (use ln) of the survival rate. Land cores,

(c)[2] The linear graph below is a semilog graph of a function. Find an explicit formula for that function (i.e., write it in the form $y = \ldots$).



7. (a)[3] Identify the domain of the function $f(x) = \ln(e^x - 1)$.

(b)[2] Find $\lim_{x\to\pi/2^+} \tan x$

(c)[2] Find $\lim_{x\to\infty} e^{-x^2}$