

Small sample correction methods for cluster-robust variance estimators and hypothesis  
tests

James E. Pustejovsky

The University of Texas at Austin

Elizabeth Tipton

Columbia University

Author Note

Abstract

*Keywords:*

Small sample correction methods for cluster-robust variance estimators and hypothesis tests

Cluster-robust variance estimators (CRVE) and hypothesis tests based upon such estimators are ubiquitous in applied econometric work. Nearly every respectable paper in the past 15 years uses cluster-robust variance estimators because to do otherwise would be to risk being seen as insufficiently rigorous (or worse, anti-conservative....ughh....how gauche!).

There's been a lot of fretting recently that even CRVE may actually not be rigorous enough. Cite the following people so as not to get their ire up:

- Brewer, Crossley, and Joyce (2013)
- Cameron, Gelbach, and Miller (2008)
- Cameron and Miller (2015)
- Carter, Schnepel, and Steigerwald (2013)
- Ibragimov and Müller (2010)
- Imbens and Kolesar (2012)
- Kezdi (2004)
- McCaffrey, Bell, and Botts (2001)
- McCaffrey and Bell (2006)
- Webb and MacKinnon (2013)
- Kline and Santos (2012)

### **Econometric framework**

We will consider linear regression models in which the errors within a cluster have an unknown variance structure. The model is

$$\mathbf{Y}_j = \mathbf{X}_j\boldsymbol{\beta} + \mathbf{e}_j, \tag{1}$$

for  $j = 1, \dots, m$ , where  $\mathbf{Y}_j$  is  $n_j \times 1$ ,  $\mathbf{X}_j$  is an  $n_j \times p$  matrix of regressors for cluster  $j$ ,  $\boldsymbol{\beta}$  is a  $p \times 1$  vector, and  $\mathbf{e}_j$  is an  $n_j \times 1$  vector of errors. Assume that  $E(\mathbf{e}_j | \mathbf{X}_j) = \mathbf{0}$  and  $\text{Var}(\mathbf{e}_j | \mathbf{X}_j) = \boldsymbol{\Sigma}_j$ , for  $j = 1, \dots, m$ , where  $\boldsymbol{\Sigma}_1, \dots, \boldsymbol{\Sigma}_m$  may be unknown. Let  $\boldsymbol{\Sigma} = \bigoplus_{j=1}^m \boldsymbol{\Sigma}_j$ .

The vector of regression coefficients is estimated by weighted least squares (WLS).

Given a set of  $m$  weighting matrices  $\mathbf{W}_1, \dots, \mathbf{W}_m$ , the WLS estimator is

$$\hat{\boldsymbol{\beta}} = \mathbf{M} \sum_{j=1}^m \mathbf{X}_j' \mathbf{W}_j \mathbf{Y}_j, \quad (2)$$

where  $\mathbf{M} = \left( \sum_{j=1}^m \mathbf{X}_j' \mathbf{W}_j \mathbf{X}_j \right)^{-1}$ .

Common choices for weighting include the unweighted case, in which  $\mathbf{W}_j$  is an identity matrix of dimension  $n_j \times n_j$ , and inverse-variance weighting under a working model. In the latter case, the errors are assumed to follow some known structure,  $\text{Var}(\mathbf{e}_j | \mathbf{X}_j) = \boldsymbol{\Phi}_j$ , where  $\boldsymbol{\Phi}_j$  is either known or a function of a low-dimensional parameter. The weighting matrices are then taken to be  $\mathbf{W}_j = \boldsymbol{\Phi}_j^{-1}$ , possibly based on estimates of the variance parameters.

The WLS estimator also encompasses the estimator proposed by Ibragimov and Müller (2010) for clustered data. Assuming that  $\mathbf{X}_j$  has rank  $p$ , their proposed approach involves estimating  $\boldsymbol{\beta}$  separately within each cluster and taking the simple average of these estimates. The resulting average is equivalent to the WLS estimator with weights

$$\mathbf{W}_j = \mathbf{X}_j \left( \mathbf{X}_j' \mathbf{X}_j \right)^{-2} \mathbf{X}_j.$$

**Cluster-robust variance estimators**

**Considerations with panel models**

**Hypothesis testing**

**Single-constraint tests**

**Multiple-constraint tests**

**Examples**

**Simulation evidence**

**Discussion**

## References

- Brewer, M., Crossley, T. F., & Joyce, R. (2013). *Inference with difference-in-differences revisited* (No. 7742). Bonn, Germany.
- Cameron, A. C., Gelbach, J. B., & Miller, D. (2008). Bootstrap-based improvements for inference with clustered errors. *The Review of Economics and Statistics*, 90(3), 414–427.
- Cameron, A. C., & Miller, D. L. (2015). *A practitioner's guide to cluster-robust inference*.
- Carter, A. V., Schnepel, K. T., & Steigerwald, D. G. (2013). Asymptotic Behavior of a t Test Robust to Cluster Heterogeneity. , 1–32.
- Ibragimov, R., & Müller, U. K. (2010). t-Statistic based correlation and heterogeneity robust inference. *Journal of Business & Economic Statistics*, 28(4), 453–468. doi: 10.1198/jbes.2009.08046
- Imbens, G. W., & Kolesar, M. (2012). *Robust standard errors in small samples: some practical advice*. Cambridge, MA. Retrieved from <http://www.nber.org/papers/w18478>
- Kezdi, G. (2004). *Robust standard error estimation in fixed-effects panel models*. Retrieved from <http://papers.ssrn.com/sol3/Delivery.cfm?abstractid=596988>
- Kline, P., & Santos, A. (2012). A score based approach to wild bootstrap inference. *Journal of Econometric Methods*, 1(1). doi: 10.1515/2156-6674.1006
- McCaffrey, D. F., & Bell, R. M. (2006). Improved hypothesis testing for coefficients in generalized estimating equations with small samples of clusters. *Statistics in medicine*, 25(23), 4081–98. doi: 10.1002/sim.2502
- McCaffrey, D. F., Bell, R. M., & Botts, C. H. (2001). Generalizations of biased reduced linearization. In *Proceedings of the annual meeting of the american statistical association*.
- Webb, M., & MacKinnon, J. G. (2013). *Wild Bootstrap Inference for Wildly Different Cluster Sizes* (No. 1314). Kingston, Ontario, Canada.