1. Introduction

A scalar records one measurement. A vector records a magnitude and direction, so it has one axis, where an axis means an independent mode of access. A matrix has two axes, and many scientific and engineering settings require tensors with three or more axes. These we call *higher valence* tensors. Tensors encapsulate all of these abstractions in one unifying concept. For instance, a matrix may be called a 2-tensor, a cube of numbers a 3-tensor, and so on. This tensor abstraction is used in various disciplines to record information for some fixed reference frame. Some complementary perspectives studying tensors include [Bro97] [KB09] [Lan12] [RS18] [DLDMV00] [Tuc66]. Below we give two real world examples.

The first is the stress tensor in mechanics. For an object such as a steel beam supporting traffic, we can track how the internal force depends on the orientation of a surface element inside the material. At each point, with respect to a coordinate system, we have a 3×3 Cauchy stress tensor, where diagonal entries capture normal compression and tension, while off-diagonal entries capture shear [GFA10]. For-axis tensors are needed to model constitutive laws, for instance mapping a strain tensor to a stress tensor, since each of the two output indices can depend on each of the two input indices.

The second comes from recommender systems, of which the Netflix tensor is the most famous. The starting point is to store a user $\times$ movie matrix of ratings. The factorization of this matrix was central to recommenders competing for the Netflix Prize [KBV09]. But recommendation systems may need more axes, for example, by considering preferences over time, and considering multiple devices. This necessitates higher valence tensors, for instance with axes user $\times$ movie $\times$ time. Tensor factorization methods explicitly model and work with these higher valence tensors [Kar+10].

1.1. Main Results.

We briefly describe the two main problems tackled by my dissertation.

The first is on computing of algebraic invariants of tensors. They have been key to discovering basis-independent cluster patterns in tensors [BKW24], decomposing p-groups [Wil09], computing endomorphisms of modules [BL08], and advancing isomorphism testing [IQ19], [BMW17] [BW12] [BMW22].

Throughout, $\mathbb{F}$ will denote a field. Suppose we are given a 3-tensor $T \in \mathbb{F}^{m\times n\times r}$ which represents trilinear measurements in a fixed basis. Let the (i,j,k)th component be denoted $T[i,j,k]$, and let $T_1,\dots,T_r \in \mathbb{F}^{m\times n}$ be slices using the Matlab/Julia inspired slice notation, where $T_i := T[:,:,i]$. For $Z = (z_{ji}) \in \mathbb{F}^{r\times r}$, define T^Z as the tensor with slices

$$(T^Z)_i := \sum_{j=1}^{r} z_{ji} T_j.$$

Equations for three algebraic invariants (adjoint, centroid, and derivation) we study are defined as

$$\begin{aligned}
&\mathrm{Adj}(T) := \{(X,Y) \in \mathbb{F}^{m\times m} \times \mathbb{F}^{n\times n} : (\forall i),\ XT_i = T_iY\ \},\\
&\mathrm{Cen}(T) := \{(X,Y,Z) \in \mathbb{F}^{m\times m} \times \mathbb{F}^{n\times n} \times \mathbb{F}^{r\times r} : (\forall i),\ XT_i = T_iY = (T^Z)_i\},\\
&\mathrm{Der}(T) := \{(X,Y,Z) \in \mathbb{F}^{m\times m} \times \mathbb{F}^{n\times n} \times \mathbb{F}^{r\times r} : (\forall i),\ XT_i + T_iY = (T^Z)_i\}.
\end{aligned} \tag{1.1}$$

Each corresponds to a linear system. Our first main result, Theorem 3.33, shows that for tensors $T \in \mathbb{F}^{n\times n\times n}$ satisfying a regularity condition, the derivation algebra of T can be computed in $O(n^{4.5})$ time. This is an $n^{0.5}$ overhead over the $O(n^4)$ cost of deterministic verification.

The second result is about understanding structure when we combine two tensors into one via multiplication. Multiplication is a natural way to inductively build up data from smaller pieces. For instance, if A and B are $\mathbb{F}$-algebras, then $A \otimes B$ is again a $\mathbb{F}$-algebra. An example of this is the extensions of scalars construction $A \otimes L$ given a $\mathbb{F}$-algebra A and a field extension $L/\mathbb{F}$.

We define a product of tensors that serves the same role by combining interactions while preserving multilinearity. Our definition is most naturally stated for bilinear maps $b : U \times V \rightarrowtail W$.

After choosing bases of U, V, and W, the map b is encoded by its structure constants, and hence by a 3-tensor. We define the product bilinear map

$$s \otimes t : (\mathbb{F}^m \otimes \mathbb{F}^p) \times (\mathbb{F}^n \otimes \mathbb{F}^q) \rightarrowtail \mathbb{F}^l \otimes \mathbb{F}^r$$

on pure tensors by $(s \otimes t)(u \otimes x,\ v \otimes y) := s(u, v) \otimes t(x, y)$, and extended bilinearly.

Example 1.1. Consider the multiplication tensors of $\mathbb{F}[x]/(x^2 + 1)$ and the multiplication tensor of $\mathbb{F}[y]/(y^2 + 5y + 10)$. Each algebra is given a $2 \times 2 \times 2$ multiplication table, and their combined multiplication is as shown below.

	1	x
1	1	x
x	x	-1

$\otimes$

	1	y
1	1	y
y	y	$-5y - 10$

$=$

	1	x	y	xy
1	1	x	y	xy
x	x	-1	xy	$-y$
y	y	xy	$-5y - 10$	$-5xy - 10x$
xy	xy	$-y$	$-5xy - 10x$	$5y + 10$

The output is the product bilinear map on the right. It is a $4 \times 4 \times 4$ tensor, with four slices indexed by the basis $\{1, x, y, xy\}$.

Our second main result, Theorem 5.1, is a theorem that states given bilinear maps s and t, we can understand $\mathrm{Der}(s \otimes t)$ by individually understanding the adjoint, centroid, and derivation algebras of s and t. This is important, because the bilinear map $s \otimes t$ preserves multilinearity at the cost of multiplicative dimension growth. For instance, given $s : \mathbb{F}^m \times \mathbb{F}^n \rightarrowtail \mathbb{F}^l$ and $t : \mathbb{F}^p \times \mathbb{F}^q \rightarrowtail \mathbb{F}^r$, the product $s \otimes t : (\mathbb{F}^m \otimes \mathbb{F}^p) \times (\mathbb{F}^n \otimes \mathbb{F}^q) \rightarrowtail \mathbb{F}^l \otimes \mathbb{F}^r$ requires $(mp)(nq)(lr)$ coordinates to describe. Yet our theorem allows for the individual understanding of the algebraic invariants of s (only mnl coordinates) and t (only pqr coordinates) to together give understanding of $\mathrm{Der}(s \otimes t)$.

2. Preliminaries and Prior Work

Given $\mathbb{F}$-vector spaces U, V and W, a function $f : U \times V \to W$ is additive if $f(u + u', v) = f(u, v) + f(u', v)$, and $f(u, v + v') = f(u, v) + f(u, v')$ for all u, u' in U and v, v' in V. A function f is $\mathbb{F}$-bilinear if f is additive and satisfies $f(cu, v) = f(u, cv) = cf(u, v)$ for all $c \in \mathbb{F}$, $u \in U$, and $v \in V$. We write $f : U \times V \rightarrowtail W$ ($\rightarrowtail$ for bilinear). When context is clear, we elide the prefix $\mathbb{F}$.

Example 2.1. Examples of bilinear maps include:

(1) If A is an $\mathbb{F}$-algebra, then multiplication map $\mu : A \times A \rightarrowtail A$ is bilinear.
(2) For positive integers m, n, p, the rectangular matrix multiplication map $\mathbb{F}^{m\times n} \times \mathbb{F}^{n\times p} \rightarrowtail \mathbb{F}^{m\times p}$ given by $(X, Y) \mapsto XY$ is bilinear.
(3) For $U = \mathbb{F}^2$ and $V = \mathbb{F}^3$, any bilinear form $b : U \times V \rightarrowtail \mathbb{F}$ can be represented by a matrix $B \in \mathbb{F}^{2\times 3}$ via $b(u, v) = u^T Bv$.

The above extends to explain $\mathbb{F}$-trilinear and $\mathbb{F}$-multilinear in general. Given a bilinear map $b : U \times V \rightarrowtail W$, fix bases for U, V, and W. Then evaluating b on a basis of U, V, and dual basis of W, we may identify b with a trilinear form in $\mathbb{F}^m \times \mathbb{F}^n \times \mathbb{F}^l \rightarrowtail \mathbb{F}$ where $m = \dim U$, $n = \dim V$, $l = \dim W$. This trilinear form is identified as a 3-tensor (as an element of $\mathbb{F}^{m\times n\times l}$), and thus throughout this dissertation, $\mathbb{F}$-bilinear, trilinear, and multilinear maps are also called *tensors*.

For vector spaces U and V, the *tensor product* of U and V is the vector space $U \otimes V$ with the canonical map $\varphi : U \times V \rightarrowtail U \otimes V$, such that for every bilinear map f with domain $U \times V$, there is a unique induced linear map $\hat{f}$ satisfying $f = \hat{f} \circ \varphi$. For vectors $s \in U$ and $t \in V$, the notation $s \otimes t$ denotes the image of (s, t) under φ. The elements $s \otimes t$ are called pure tensors.

Next we describe subspaces of a tensor product space. For subsets $A \subseteq U$ and $B \subseteq V$, define the subspace

$$A \otimes B := \mathrm{span}\{a \otimes b : a \in A,\ b \in B\} \leq U \otimes V.$$

If $A \leq U$ and $B \leq V$ are subspaces, then there is also the *external* tensor product space $(A \otimes B, \varphi : A \times B \rightarrowtail A \otimes B)$. For $\iota_A : A \hookrightarrow U$ and $\iota_B : B \hookrightarrow V$ inclusion maps, the induced map $\iota_A \otimes \iota_B : A \otimes B \to U \otimes V$ is injective, and we identify the external tensor product with its image, which is precisely the span of $\{a \otimes b : a \in A, b \in B\}$. This clarifies the notation $A \otimes B$ when A and B are subspaces of ambient vector spaces.

Lemma 2.2. Let U, V be vector spaces, and $A \leq U$ and $X \leq V$ be subspaces. Then $A \otimes X = (U \otimes X) \cap (A \otimes V)$ as subspaces of $U \otimes V$.

Proof. Let $\iota_A : A \hookrightarrow U$ and $\iota_X : X \hookrightarrow V$ be the inclusion maps. The intersection of $U \otimes X$ and $A \otimes V$ in $U \otimes V$ is the pullback of the two inclusion maps. Hence it suffices to show the following square is also a pullback:

$$\begin{tikzcd}
A \otimes X \arrow[r, "\iota_A \otimes \mathrm{id}_X"] \arrow[d, "\mathrm{id}_A \otimes \iota_X"'] & U \otimes X \arrow[d, "\mathrm{id}_U \otimes \iota_X"] \\
A \otimes V \arrow[r, "\iota_A \otimes \mathrm{id}_V"'] & U \otimes V
\end{tikzcd}$$

Let T be a vector space with maps $q_1 : T \to A \otimes V$ and $q_2 : T \to U \otimes X$ such that $(\iota_A \otimes \mathrm{id}_V) \circ q_1 = (\mathrm{id}_U \otimes \iota_X) \circ q_2$. We must construct the unique u making both triangles commute in the below diagram.

$$\begin{tikzcd}
T \arrow[dr, dashed, "u"] \arrow[drr, bend left, "q_2"] \arrow[ddr, bend right, "q_1"'] & & \\
& A \otimes X \arrow[r, "\iota_A \otimes \mathrm{id}_X"] \arrow[d, "\mathrm{id}_A \otimes \iota_X"'] & U \otimes X \arrow[d, "\mathrm{id}_U \otimes \iota_X"] \\
& A \otimes V \arrow[r, "\iota_A \otimes \mathrm{id}_V"'] & U \otimes V
\end{tikzcd}$$

This is done by showing $\mathrm{Im}(q_1) \leq \mathrm{Im}(\mathrm{id}_A \otimes \iota_X)$. Since $\mathrm{id}_A \otimes \iota_X$ is injective, any such factorization is unique. Since tensoring over a field preserves exact sequences, tensoring $0 \to X \xrightarrow{\iota_X} V \xrightarrow{q} V/X \to 0$ with A gives

$$0 \to A \otimes X \xrightarrow{\mathrm{id}_A \otimes \iota_X} A \otimes V \xrightarrow{\mathrm{id}_A \otimes q} A \otimes (V/X) \to 0.$$

So we must show $\mathrm{Im}(q_1) \leq \mathrm{Ker}(\mathrm{id}_A \otimes q)$, or $(\mathrm{id}_A \otimes q) \circ q_1 = 0$. This fact follows by the below commutative square.

$$\begin{tikzcd}
A \otimes V \arrow[r, "\iota_A \otimes \mathrm{id}_V"] \arrow[d, "\mathrm{id}_A \otimes q"'] & U \otimes V \arrow[d, "\mathrm{id}_U \otimes q"] \\
A \otimes (V/X) \arrow[r, "\iota_A \otimes \mathrm{id}_{V/X}"'] & U \otimes (V/X)
\end{tikzcd}$$

Indeed, as elements in $\mathrm{Hom}(A \otimes V, U \otimes (V/X))$,

$$(\mathrm{id}_U \otimes q) \circ (\iota_A \otimes \mathrm{id}_V) = (\iota_A \otimes \mathrm{id}_{V/X}) \circ (\mathrm{id}_A \otimes q), \tag{2.1}$$

because both sides send the pure tensor $a \otimes v$ to $\iota_A(a) \otimes q(v)$.

Applying $\mathrm{id}_U \otimes q$ to $(\iota_A \otimes \mathrm{id}_V) \circ q_1 = (\mathrm{id}_U \otimes \iota_X) \circ q_2$, the right-hand side vanishes because $(\mathrm{id}_U \otimes q) \circ (\mathrm{id}_U \otimes \iota_X) = \mathrm{id}_U \otimes (q \circ \iota_X) = 0$. Hence $(\mathrm{id}_U \otimes q) \circ (\iota_A \otimes \mathrm{id}_V) \circ q_1 = 0$, and by (2.1), $(\iota_A \otimes \mathrm{id}_{V/X}) \circ (\mathrm{id}_A \otimes q) \circ q_1 = 0$. Since $\iota_A \otimes \mathrm{id}_{V/X}$ is injective, $(\mathrm{id}_A \otimes q) \circ q_1 = 0$ as needed. Thus we find the unique u such that $q_1 = (\mathrm{id}_A \otimes \iota_X) \circ u$.

Finally, for u to make the upper triangule commute, we must show $q_2 = (\iota_A \otimes \mathrm{id}_X) \circ u$. We have

$$\begin{aligned}
(\mathrm{id}_U \otimes \iota_X) \circ (\iota_A \otimes \mathrm{id}_X) \circ u &= (\iota_A \otimes \mathrm{id}_V) \circ (\mathrm{id}_A \otimes \iota_X) \circ u \\
&= (\iota_A \otimes \mathrm{id}_V) \circ q_1 \\
&= (\mathrm{id}_U \otimes \iota_X) \circ q_2,
\end{aligned}$$

Therefore $(\mathrm{id}_U \otimes \iota_X) \circ \big((\iota_A \otimes \mathrm{id}_X) \circ u - q_2\big) = 0$. As $\mathrm{id}_U \otimes \iota_X$ is injective, we get $q_2 = (\iota_A \otimes \mathrm{id}_X) \circ u$ as needed. $\square$

2.1. **Algebras and Actions on bilinear maps.** Existing work such as [Jac37], [Mya90], [BW14], and [Wil16] highlights the role of the three algebras we saw in Equation (1.1). In this subsection, we expand on that and describe some operations and properties that we need for the future sections.

As before U, V, and W are finite-dimensional $\mathbb{F}$-vector spaces. Let $\mathrm{End}(U) = \mathrm{Hom}(U, U)$ be the $\mathbb{F}$-linear endomorphisms of a vector space U, and let $b : U \times V \rightarrowtail W$ be a bilinear map. We will also use $\mathrm{Bil}(U, V; W)$ for the $\mathbb{F}$-vector space of bilinear maps $U \times V \rightarrowtail W$. For $\alpha \in \mathrm{End}(U)$, $\beta \in \mathrm{End}(V)$, and $\gamma \in \mathrm{End}(W)$, define the bilinear maps arising from operators acting on axes as

$$\begin{aligned} b \bullet_2 \alpha &: U \times V \rightarrowtail W, & (u, v) &\mapsto b(\alpha u, v), \\ b \bullet_1 \beta &: U \times V \rightarrowtail W, & (u, v) &\mapsto b(u, \beta v), \\ b \bullet_0 \gamma &: U \times V \rightarrowtail W, & (u, v) &\mapsto \gamma(b(u, v)). \end{aligned}$$

Definition 2.3. The left, middle, and right nuclei of b are respectively

$$\begin{aligned} \mathcal{L}(b) &\coloneqq \{(\alpha, \gamma) \in \mathrm{End}(U) \times \mathrm{End}(W) : b \bullet_2 \alpha = b \bullet_0 \gamma\}, \\ \mathcal{M}(b) &\coloneqq \{(\alpha, \beta) \in \mathrm{End}(U)^{\mathrm{op}} \times \mathrm{End}(V) : b \bullet_2 \alpha = b \bullet_1 \beta\}, \\ \mathcal{R}(b) &\coloneqq \{(\beta, \gamma) \in \mathrm{End}(V) \times \mathrm{End}(W) : b \bullet_1 \beta = b \bullet_0 \gamma\}. \end{aligned}$$

Here $\mathrm{End}(U)^{\mathrm{op}}$ is the opposite algebra of $\mathrm{End}(U)$, meaning composition order is reversed. The reason for using $\mathrm{End}(U)^{\mathrm{op}}$ in $\mathcal{M}(b)$ is that if $(\alpha, \beta), (\alpha', \beta') \in \mathcal{M}(b)$, then $b((\alpha'\alpha)u, v) = b(\alpha u, \beta' v) = b(u, (\beta\beta')v)$, so the product is $(\alpha, \beta)(\alpha', \beta') = (\alpha'\alpha, \beta\beta')$. Notice $\mathcal{M}(b)$ is the coordinate free description of the adjoint algebra in Equation (1.1).

The centroid of b is

$$\mathcal{C}(b) \coloneqq \{(\alpha, \beta, \gamma) \in \mathrm{End}(U) \times \mathrm{End}(V) \times \mathrm{End}(W) : b \bullet_2 \alpha = b \bullet_1 \beta = b \bullet_0 \gamma\},$$

and the derivation algebra of b is

$$\mathrm{Der}(b) \coloneqq \{(\alpha, \beta, \gamma) \in \mathrm{End}(U) \times \mathrm{End}(V) \times \mathrm{End}(W) : b \bullet_2 \alpha + b \bullet_1 \beta = b \bullet_0 \gamma\}.$$

Remark 2.4. Each of $\mathcal{L}(b)$, $\mathcal{M}(b)$, $\mathcal{R}(b)$, $\mathcal{C}(b)$, $\mathrm{Der}(b)$ is an $\mathbb{F}$-vector subspace of the corresponding product of endomorphism spaces. Furthermore, $\mathcal{L}(b)$, $\mathcal{M}(b)$, $\mathcal{R}(b)$, and $\mathcal{C}(b)$ are associative algebras, with $\mathcal{C}(b)$ also commutative when b is fully nondegenerate, while $\mathrm{Der}(b)$ is a Lie algebra. [BMW20, Section 1.2].

2.2. **Radicals and non-degeneracy of bilinear maps.** Let $b : U \times V \rightarrowtail W$ be a bilinear map. For $u \in U$ and $v \in V$, define

$$b(u, V) \coloneqq \{b(u, v') : v' \in V\}, \quad b(U, v) \coloneqq \{b(u', v) : u' \in U\}, \quad b(U, V) \coloneqq \mathrm{span}\{b(u', v') : u' \in U,\ v' \in V\}.$$

The *left radical*, *right radical*, and *coradical* of a bilinear map b are respectively

$$\mathrm{Rad}_L(b) \coloneqq \{u \in U : b(u, V) = 0\}, \quad \mathrm{Rad}_R(b) \coloneqq \{v \in V : b(U, v) = 0\}, \quad \mathrm{Corad}(b) \coloneqq W/b(U, V).$$

We call b *fully nondegenerate* if $\mathrm{Rad}_L(b) = \mathrm{Rad}_R(b) = 0$ and $\mathrm{Corad}(b) = 0$. Now define the *axis actions*

$$\begin{aligned} \Lambda_2^b &: \mathrm{End}(U) \to \mathrm{Bil}(U, V; W), \quad \alpha \mapsto b \bullet_2 \alpha, \\ \Lambda_1^b &: \mathrm{End}(V) \to \mathrm{Bil}(U, V; W), \quad \beta \mapsto b \bullet_1 \beta, \\ \Lambda_0^b &: \mathrm{End}(W)^{\mathrm{op}} \to \mathrm{Bil}(U, V; W), \quad \gamma \mapsto b \bullet_0 \gamma. \end{aligned}$$

The superscript is suppressed when context is clear.

Lemma 2.5. If $b : U \times V \rightarrowtail W$ is fully nondegenerate, then Λ_2, Λ_1, and Λ_0 are injective.

Proof. If $\Lambda_2(\alpha) = 0$, then $b(\alpha u, v) = 0$ for all $u \in U$ and $v \in V$, so $\alpha u \in \mathrm{Rad}_L(b)$ for every u. Since $\mathrm{Rad}_L(b) = 0$, we get $\alpha = 0$. The proof for Λ_1 is analogous. For Λ_0, if $\Lambda_0(\gamma) = 0$, then $\gamma(b(u, v)) = 0$ for all u, v, so γ vanishes on $b(U, V) = W$. Hence $\gamma = 0$. □

Fact 2.6. [Wil15, Section 8.3] For any bilinear map $b : U \times V \rightarrowtail W$, there is a canonical induced fully nondegenerate bilinear map

$$b/\sqrt{b} : \big(U/\operatorname{Rad}_L(b)\big) \times \big(V/\operatorname{Rad}_R(b)\big) \rightarrowtail b(U,V),$$

defined by $\big(b/\sqrt{b}\big)\big(u + \operatorname{Rad}_L(b), v + \operatorname{Rad}_R(b)\big) := b(u,v)$.

The algebraic invariants of $b/\sqrt{b}$ gives us an understanding of the invariants of b through the exact sequence

$$0 \to \operatorname{Hom}(U, \operatorname{Rad}_L(b)) \times \operatorname{Hom}(\operatorname{Corad}(b), W) \to \mathcal{L}(b) \xrightarrow{\rho_L} \mathcal{L}(b/\sqrt{b}) \to 0,$$

where $\rho_L(\alpha,\gamma) := (\bar{\alpha}, \gamma|_{b(U,V)})$ and $\bar{\alpha}$ is induced by α on $U/\operatorname{Rad}_L(b)$. We justify this exact sequence. First, for well-definedness of ρ_L, we prove $(\alpha,\gamma) \in \mathcal{L}(b)$ satisfies $\alpha(\operatorname{Rad}_L(b)) \leq \operatorname{Rad}_L(b)$, and $\gamma\big(b(U,V)\big) \leq b(U,V)$. Let $u \in \operatorname{Rad}_L(b)$. Then $b(u,v) = 0$ for all v, so $\gamma(b(u,v)) = \gamma(0) = 0$. But $\gamma(b(u,v)) = b(\alpha u, v)$ as well, so $\alpha u \in \operatorname{Rad}_L(b)$. The proof for $\gamma(b(U,V)) \leq b(U,V)$ is analogous.

To see the image of ρ_L lies in $\mathcal{L}(b/\sqrt{b})$, we check for all $u \in U$, $v \in V$,

$$\begin{aligned}\big(b/\sqrt{b}\big)\big(\bar{\alpha}(u + \operatorname{Rad}_L(b)), v + \operatorname{Rad}_R(b)\big) &= b(\alpha u, v) = \gamma(b(u,v)) \\ &= \gamma|_{b(U,V)}\Big(\big(b/\sqrt{b}\big)\big(u + \operatorname{Rad}_L(b), v + \operatorname{Rad}_R(b)\big)\Big),\end{aligned}$$

so $(\bar{\alpha}, \gamma|_{b(U,V)}) \in \mathcal{L}(b/\sqrt{b})$. We now show ρ_L is surjective. Given $(\bar{\alpha}, \gamma_0) \in \mathcal{L}(b/\sqrt{b})$, choose the lift $\alpha \in \operatorname{End}(U)$ of $\bar{\alpha}$ with $\alpha|_{\operatorname{Rad}_L(b)} = 0$, and choose $\gamma \in \operatorname{End}(W)$ with $\gamma|_{b(U,V)} = \gamma_0$ and $\gamma = 0$ on a chosen complement of $b(U,V)$. Then for all $u \in U$ and $v \in V$,

$$\begin{aligned}b(\alpha u, v) &= \big(b/\sqrt{b}\big)\big(\bar{\alpha}(u + \operatorname{Rad}_L(b)), v + \operatorname{Rad}_R(b)\big) \\ &= \gamma_0\Big(\big(b/\sqrt{b}\big)\big(u + \operatorname{Rad}_L(b), v + \operatorname{Rad}_R(b)\big)\Big) = \gamma(b(u,v)).\end{aligned}$$

Hence $(\alpha,\gamma) \in \mathcal{L}(b)$ as needed. The kernel of ρ_L are all pairs of endomorphisms where the first component acts like zero modulo $\operatorname{Rad}_L(b)$ (hence $\alpha(U) \subseteq \operatorname{Rad}_L(b)$) and the second component is zero restricted to $b(U,V)$. Thus,

$$\operatorname{Ker}(\rho_L) = \operatorname{Hom}(U, \operatorname{Rad}_L(b)) \times \{\gamma \in \operatorname{End}(W) : \gamma|_{b(U,V)} = 0\} \cong \operatorname{Hom}(U, \operatorname{Rad}_L(b)) \times \operatorname{Hom}(\operatorname{Corad}(b), W).$$

2.3. **Product of tuples of endomorphisms.** For $k \in \{0,1,2\}$, let U_k and V_k be finite-dimensional $\mathbb{F}$-vector spaces. We use the Kronecker isomorphisms $\kappa_k : \operatorname{End}(U_k) \otimes \operatorname{End}(V_k) \to \operatorname{End}(U_k \otimes V_k)$, defined by $\kappa_k(\alpha \otimes f)(u \otimes v) := \alpha u \otimes f v$ for all $u \in U_k$ and $v \in V_k$. The following is notation for combining tuples of endomorphisms.

Definition 2.7. Let K be finite. For each $k \in K$, let U_k, V_k be finite-dimensional $\mathbb{F}$-vector spaces. Define

$$\boxtimes : \Big(\prod_{k\in K} \operatorname{End}(U_k)\Big) \times \Big(\prod_{k\in K} \operatorname{End}(V_k)\Big) \to \prod_{k\in K} \operatorname{End}(U_k \otimes V_k)$$

by $((\alpha_k)_{k\in K}, (f_k)_{k\in K}) \longmapsto (\kappa_k(\alpha_k \otimes f_k))_{k\in K} =: (\alpha_k)_{k\in K} \boxtimes (f_k)_{k\in K}$.

For subspaces $A \leq \prod_{k\in K} \operatorname{End}(U_k)$ and $B \leq \prod_{k\in K} \operatorname{End}(V_k)$, define

$$A \boxtimes B := \operatorname{span}\{a \boxtimes b : a \in A,\ b \in B\} \leq \prod_{k\in K} \operatorname{End}(U_k \otimes V_k).$$

2.4. **Product of bilinear maps.** As described in Section 1, there is a need to combine two bilinear maps into one using a product construction.

Definition 2.8. Fix two bilinear maps $s : U_2 \times U_1 \rightarrowtail U_0$, and $t : V_2 \times V_1 \rightarrowtail V_0$. The *tensor product bilinear map*

$$s \otimes t : (U_2 \otimes V_2) \times (U_1 \otimes V_1) \rightarrowtail U_0 \otimes V_0$$

is the unique bilinear map defined in pure tensors as $(s \otimes t)((u_2 \otimes v_2), (u_1 \otimes v_1)) = s(u_2, u_1) \otimes t(v_2, v_1)$ and extended bilinearly.

Lemma 2.9. If s and t are fully nondegenerate bilinear maps, then $s \otimes t$ is fully nondegenerate.

Proof. For nondegeneracy of $U_2 \otimes V_2$, let $x = \sum_j u_{2,j} \otimes v_{2,j} \in U_2 \otimes V_2$ and assume $(s \otimes t)(x, (u_1 \otimes v_1)) = 0$ for all $u_1 \in U_1$ and $v_1 \in V_1$. We need to show $x = 0$. Define the curried functions $\lambda_s : U_2 \to \mathrm{Hom}(U_1, U_0)$ by $\lambda_s(u_2)(u_1) := s(u_2, u_1)$ and $\lambda_t : V_2 \to \mathrm{Hom}(V_1, V_0)$ by $\lambda_t(v_2)(v_1) := t(v_2, v_1)$. Identify $\mathrm{Hom}(U_1, U_0) \otimes \mathrm{Hom}(V_1, V_0)$ with $\mathrm{Hom}(U_1 \otimes V_1,\ U_0 \otimes V_0)$ via $(f \otimes g)(u_1 \otimes v_1) = f(u_1) \otimes g(v_1)$. Under this identification, the assumption gives $(\lambda_s \otimes \lambda_t)(x) = 0$. Since $\mathrm{Ker}(\lambda_s) = \mathrm{Rad}_L(s) = 0$ and $\mathrm{Ker}(\lambda_t) = \mathrm{Rad}_L(t) = 0$, both maps are injective, so $\lambda_s \otimes \lambda_t$ is injective and $x = 0$. The argument for $U_1 \otimes V_1$ is analogous.

For the coradical, write $u_0 = \sum_a s(u_{2,a}, u_{1,a})$ and $v_0 = \sum_b t(v_{2,b}, v_{1,b})$. Then

$$\begin{aligned} u_0 \otimes v_0 &= \left(\sum_a s(u_{2,a}, u_{1,a}) \right) \otimes \left(\sum_b t(v_{2,b}, v_{1,b}) \right) \\ &= \sum_{a,b} (s \otimes t)((u_{2,a} \otimes v_{2,b}), (u_{1,a} \otimes v_{1,b})), \end{aligned}$$

so $(s \otimes t)(U_2 \otimes V_2, U_1 \otimes V_1) = U_0 \otimes V_0$. This concludes $\mathrm{Corad}(s \otimes t) = 0$. □

Remark 2.10. If s and t are fully nondegenerate, then the axis action maps $\Lambda_2^{s \otimes t}$, $\Lambda_1^{s \otimes t}$, and $\Lambda_0^{s \otimes t}$ are injective by Lemmas 2.5 and 2.9.

Next we come to a key correspondence that allows us to analyze axis actions on tensor product bilinear maps by *linear algebra* techniques. Recall for X, Y vector spaces, and $(X \otimes Y, \varphi : X \times Y \rightarrowtail X \otimes Y)$ the tensor product space of X and Y, the element $\varphi(x, y) \in X \otimes Y$ is the pure tensor often written as $x \otimes y$. As the notation $x \otimes y$ can be ambiguous when x and y are themselves bilinear maps, when wanting to refer to the pure tensor element in the tensor product vector space, we write $\varphi(x, y)$.

Let $\mathcal{U} := \mathrm{Bil}(U_2, U_1; U_0)$, $\mathcal{V} := \mathrm{Bil}(V_2, V_1; V_0)$, and define the bilinear map

$$\Theta : \mathcal{U} \times \mathcal{V} \to \mathrm{Bil}(U_2 \otimes V_2, U_1 \otimes V_1; U_0 \otimes V_0), \quad \Theta(s, t) := s \otimes t \text{ as the tensor product bilinear map.}$$

The universal property of the tensor product space $\mathcal{U} \otimes \mathcal{V}$ gives a unique induced linear map

$$\widetilde{\Theta} : \mathcal{U} \otimes \mathcal{V} \to \mathrm{Bil}(U_2 \otimes V_2, U_1 \otimes V_1; U_0 \otimes V_0).$$

Write $u_k := \dim(U_k)$ and $v_k := \dim(V_k)$ for all $k \in \{0, 1, 2\}$. Then

$$\dim(\mathcal{U} \otimes \mathcal{V}) = \dim(\mathcal{U}) \dim(\mathcal{V}) = (u_2 u_1 u_0)(v_2 v_1 v_0) = (u_2 v_2)(u_1 v_1)(u_0 v_0) = \dim(\mathrm{Bil}(U_2 \otimes V_2, U_1 \otimes V_1; U_0 \otimes V_0))$$

Since the domain and codomain of $\widetilde{\Theta}$ have the same dimension, it suffices to show the map is injective to conclude that it is an isomorphism. Choose bases $e_{k,i}$ of U_k and $f_{k,j}$ of V_k, and let $s_{ij}^r \in \mathcal{U}$ and $t_{pq}^\ell \in \mathcal{V}$ be the bilinear maps sending $(e_{2,i}, e_{1,j})$ to $e_{0,r}$ and $(f_{2,p}, f_{1,q})$ to $f_{0,\ell}$, respectively, and all other basis pairs to 0. These form bases of $\mathcal{U}$ and $\mathcal{V}$. If $x \in \mathrm{Ker}(\widetilde{\Theta})$ is written in this basis, then evaluating $\widetilde{\Theta}(x)$ at $(e_{2,i} \otimes f_{2,p},\ e_{1,j} \otimes f_{1,q})$ produces a linear combination of $e_{0,r} \otimes f_{0,\ell}$ whose coefficients are exactly those of the terms $\varphi(s_{ij}^r, t_{pq}^\ell)$ in a basis expansion of x. Hence every coefficient of x is 0, concluding $\widetilde{\Theta}$ is injective.

We shall need the notation $\Xi := \widetilde{\Theta}^{-1}$ below. Also recall the Kronecker isomorphism $\kappa_k : \mathrm{End}(U_k) \otimes \mathrm{End}(V_k) \to \mathrm{End}(U_k \otimes V_k)$ given by $\kappa_k(\alpha \otimes f)(u \otimes v) = \alpha u \otimes f v$.

Lemma 2.11. Following the notation above, let $s \in \mathcal{U}$, $t \in \mathcal{V}$ be bilinear maps, and $(\mathcal{U} \otimes \mathcal{V}, \varphi : \mathcal{U} \times \mathcal{V} \rightarrowtail \mathcal{U} \otimes \mathcal{V})$ the tensor product space of $\mathcal{U}$ with $\mathcal{V}$. For all $k \in \{0, 1, 2\}$, $\alpha \in \mathrm{End}(U_k)$, and $f \in \mathrm{End}(V_k)$,

$$\Xi\big((s \otimes t) \bullet_k \kappa_k(\alpha \otimes f)\big) = \varphi(s \bullet_k \alpha, t \bullet_k f).$$

Proof. By definition of $\widetilde{\Theta}$, we have $\widetilde{\Theta}\big(\varphi(s \bullet_k \alpha, t \bullet_k f)\big) = \Theta(s \bullet_k \alpha, t \bullet_k f)$. Evaluating both the bilinear map $\Theta(s \bullet_k \alpha, t \bullet_k f)$ and $(s \otimes t) \bullet_k \kappa_k(\alpha \otimes f)$ on pure tensors shows they agree, so applying Ξ gives the claim. □

As a consequence, for $k \in \{0,1,2\}$ and $x \in \mathrm{End}(U_k \otimes V_k)$ with decomposition $x = \sum_j \kappa_k(\alpha_j \otimes f_j)$, linearity gives

$$\Xi\big((s \otimes t) \bullet_k x\big) = \sum_j \Xi\big((s \otimes t) \bullet_k \kappa_k(\alpha_j \otimes f_j)\big) = \sum_j \varphi(s \bullet_k \alpha_j, t \bullet_k f_j). \tag{2.2}$$

2.5. **Prior work.** Existing research places importance on the algebras coming from Equation (1.1). Myasnikov, Wilson, and others in [Mya90], [Wil16], [Wil12], [MM10], [Wil09] prove for the bilinear map b that the algebras $\mathrm{Adj}(b)$ and $\mathrm{Cen}(b)$ control direct sum decompositions and automorphisms of b, using them to prove properties for the originating algebraic structures. The most famous special case for the adjoint equation is the theory of Kronecker modules, which in our terminology corresponds to tensors in $\mathbb{F}^{m \times n \times 2}$ [Kro90]. Recent work [BKW24] shows that derivations also control the original tensor's null patterns, limiting the subspaces on which the tensor is non-zero.

There has also been prior work on understanding derivations of tensors coming from the multiplication tensors of algebras. Leger and Luks study generalized derivations, quasiderivations, and quasicentroids as extensions of algebra derivations [LL00], and Novotný and Hřivnák study (α, β, γ)-derivations together with associated invariant functions [NH08]. Closer to our results, Brešar studies derivations of tensor products of nonassociative algebras [Bre17]. Benkart and Osborn study derivations and automorphisms of nonassociative matrix algebras [BO81], and Azam studies derivations of tensor products of algebras [Aza08].

3. Algorithms for adjoint and derivation equations

The descriptions of the adjoint, centroid, and derivation algebras in Equation (1.1) are linear in the unknown matrix entries. So each admit a solution as a large system of linear equations. In collaboration with Joshua Maglione and James B. Wilson, we obtain an asymptotically faster algorithm for computing the adjoint algebra in certain scenarios, inspired by analogous results for matrices known as the Bartels-Stewart algorithm [BS72]. In this chapter, I state and prove results from this joint work using a novel approach. I also state and prove my own original result, which concerns the derivation algebra, following a similar approach.

Theorem 3.21. For cubic $(n \times n \times n)$ tensors, under the regularity conditions (established in Definition 3.32), SimultaneousSylvesterSystem is solved in the same cost as the full deterministic verification. This is $O(n^4)$ cost, as compared to the naive $O(n^7)$ cost by solving as a linear system of equations.

Theorem 3.33. For cubic $(n \times n \times n)$ tensors, under the regularity conditions (established in Definition 3.38), DerivationSystem is solved in cost $O(n^{4.5})$. This is an additional $n^{1/2}$ overhead compared to the deterministic verification cost of $O(n^4)$, as compared to the naive $O(n^7)$ cost by solving as a linear system of equations.

Our strategy in this section is to reduce each large problem to a set of smaller linear algebra problems. Their solutions are affine subspaces, which we define below.

3.1. **Preliminaries.** An *affine space* is a pair (A, U) where A is a non-empty set and U is a vector space acting on A by a fixed-point free and transitive action. Write the action as $+ : A \times U \to A$, $(a, u) \mapsto a + u$. We call U the *direction space* of A. When U is clear from context, we also say "A is an affine space".

Lemma 3.1. Let (A, U) be an affine space. For any $a, b \in A$, there is a unique $u \in U$ such that $a + u = b$. We denote this vector by $b - a$.

Proof. Existence follows from transitivity. If $a + u = b = a + v$, then $a + (u - v) = a$. Fixed-point freeness gives $u - v = 0$, so $u = v$. □

Let (A, U) be a d-dimensional affine space. An *affine frame* is a tuple $(a_0, \ldots, a_d)$ in A such that $\{a_1 - a_0, \ldots, a_d - a_0\}$ is a basis of U. Every $a \in A$ has unique coordinates $\lambda = (\lambda_1, \ldots, \lambda_d) \in \mathbb{F}^d$ with $a = a_0 + \sum_{i=1}^{d} \lambda_i (a_i - a_0)$.

Convention 3.2. For computation, we encode an affine space by an affine frame.

Let (A, U) and (B, V) be affine spaces. A map $f : A \to B$ is an *affine map* if there exists a linear map $L : U \to V$ such that $f(a + u) = f(a) + L(u)$ for all $a \in A$ and $u \in U$. We call L the underlying linear map of f. Below we list some properties of affine spaces and maps that we will use in following sections without further comment.

(1) Every vector space V is also an affine space with direction space V.
(2) If A, B are vector spaces and $f : A \to B$ is a linear map, then f is also an affine map.
(3) If (A, U) and (B, V) are affine spaces, then $(A \times B, U \times V)$ is an affine space with action $(a, b) + (u, v) := (a + u, b + v)$.
(4) An affine map $f : A \times B \to C$ has an underlying linear map $L : U \times V \to W$ with $f\big((a, b) + (u, v)\big) = f(a, b) + L(u, v)$.
(5) Let U and V be vector spaces, and let $L : U \to V$ be linear. If (A, W) is an affine subspace of U, then $L|_A : A \to V$ is affine, with underlying linear map $L|_W$.

Example 3.3. Let $A = \mathbb{R}^3$, $a_0 = (1, 2, 3)$, and $W = \{(x, y, 0) : x, y \in \mathbb{R}\} \leq \mathbb{R}^3$. Then $a_0 + W$ is the plane $z = 3$, an affine subspace of $(\mathbb{R}^3, \mathbb{R}^3)$.

As an example of (5) above, if $L : \mathbb{R}^3 \to \mathbb{R}$ is the linear map $L(x, y, z) = x + y + z$, then $L|_{a_0+W}$ is the affine map $(x, y, 3) \mapsto x + y + 3$, with underlying linear map $L|_W : (x, y, 0) \mapsto x + y$.

Finally, we describe how we analyze algorithms and our notation for tensors.

Convention 3.4. For functions $f, g : \mathbb{N} \to \mathbb{N}$, we write $f \in O(g)$ if there exist constants $C, n_0 \in \mathbb{N}$ such that $f(n) \leq C\, g(n)$ for all $n \geq n_0$.

Convention 3.5. We always use the standard basis for coordinate vector spaces $\mathbb{F}^n$. For $u = (u_1, \ldots, u_n)$ and $v = (v_1, \ldots, v_n)$, we use the bilinear pairing $\langle u, v \rangle := \sum_{i=1}^{n} u_i v_i$.

Convention 3.6. We analyze algorithms by their arithmetic complexity, meaning the number of field operations over $\mathbb{F}$ (addition, subtraction, multiplication, and division).

Definition 3.7. Let $T \in \mathbb{F}^{a \times b \times c}$ be a tensor, let $I \subseteq [a]$, $J \subseteq [b]$, and let $k \in [c]$. We use the slice notation

$$T[I, :, k] \in \mathbb{F}^{|I| \times b}, \qquad T[:, J, k] \in \mathbb{F}^{a \times |J|}, \qquad T[I, J, k] \in \mathbb{F}^{|I| \times |J|}.$$

For $M \in \mathbb{F}^{a \times b}$ a matrix, we write

$$M[I, :] \in \mathbb{F}^{|I| \times b}, \qquad M[:, J] \in \mathbb{F}^{a \times |J|}.$$

The indexing of sliced tensors and matrices is by the fixed ordering of the slicing sets. For instance, for $I = \{i_1 < \cdots < i_t\} \subset [a]$, the restricted matrix $X[I, :] \in \mathbb{F}^{t \times r}$ has $X[I, :][j, k] = X[i_j, k]$. We use this convention for all restricted tensor and matrix indexing.

Example 3.8. Let $T \in \mathbb{F}^{2 \times 2 \times 2}$ be defined by $T[:, :, 1] = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$, $T[:, :, 2] = \begin{bmatrix} 5 & 6 \\ 7 & 8 \end{bmatrix}$,. We have $T[\{1\}, :, 1] = \begin{bmatrix} 1 & 2 \end{bmatrix}$, and $T[:, \{2\}, 1] = \begin{bmatrix} 2 \\ 4 \end{bmatrix}$. In the matrix case, let $M = \begin{bmatrix} 9 & 10 & 11 \\ 12 & 13 & 14 \end{bmatrix} \in \mathbb{F}^{2 \times 3}$. Then $M[:, \{2, 3\}] = \begin{bmatrix} 10 & 11 \\ 13 & 14 \end{bmatrix} \in \mathbb{F}^{2 \times 2}$.

Definition 3.9. We define vectorization for matrices and tensors. For $M = (m_{ij}) \in \mathbb{F}^{a \times b}$, $\mathrm{vec}(M) \in \mathbb{F}^{ab}$ for $(i, j) \in [a] \times [b]$ by

$$\mathrm{vec}(M)_{\,i+(j-1)a} = m_{ij}.$$

For $T = (t_{ijk}) \in \mathbb{F}^{a \times b \times c}$, define $\mathrm{vec}(T) \in \mathbb{F}^{abc}$ for $(i, j, k) \in [a] \times [b] \times [c]$ by

$$\mathrm{vec}(T)_{\,i+(j-1)a+(k-1)ab} = t_{ijk}.$$

Example 3.10. For

$$M = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix} \in \mathbb{F}^{2\times 3}, \qquad T \in \mathbb{F}^{2\times 2\times 2}, \quad T[:,:,1] = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}, \quad T[:,:,2] = \begin{bmatrix} 5 & 6 \\ 7 & 8 \end{bmatrix},$$

we have $\mathrm{vec}(M) = (1,4,2,5,3,6)$, and $\mathrm{vec}(T) = (1,3,2,4,5,7,6,8)$.

3.2. Linear Algebra subroutines.

Lemma 3.11. Let (A,U) and (B,V) be affine spaces, let $f : A \to B$ be affine with underlying linear map $L : U \to V$, and let (S,W) be an affine subspace of B. Then the preimage, $f^{-1}(S) \subseteq A$, is either empty or an affine subspace of A.

Proof. If $f^{-1}(S) = \varnothing$, there is nothing to prove. Assume $f^{-1}(S) \neq \varnothing$, choose $a_0 \in f^{-1}(S)$, and set $U_0 := L^{-1}(W) \leq U$. For $u \in U$,

$$f(a_0 + u) = f(a_0) + L(u).$$

Since $f(a_0) \in S$ and S has direction space W,

$$f(a_0 + u) \in S \iff L(u) \in W \iff u \in U_0.$$

Hence $f^{-1}(S) = a_0 + U_0$, so $f^{-1}(S)$ is an affine subspace. □

Problem LinSolve($\mathbb{F}$).

Given: $\mathbb{F}$-vector spaces U, V, a linear map $L : U \to V$, and $v \in V$.
Return: A computational encoding of the solution set $\{u \in U : L(u) = v\}$.

Proposition 3.12. LinSolve($\mathbb{F}$) returns either an empty set or an affine subspace of U.

Proof. The map L is affine and $\{v\}$ is an affine subspace of V with direction space $\{0\}$, so the solution set is $L^{-1}(\{v\})$. Applying Lemma 3.11 concludes the proposition. □

Proposition 3.13. A coordinatized instance of LinSolve($\mathbb{F}$), with

$$U = \mathbb{F}^{n\times \ell}, \qquad V = \mathbb{F}^{m\times \ell}, \qquad L(X) = MX, \qquad M \in \mathbb{F}^{m\times n},$$

is solvable in cost $\mathtt{T}_{p1}(m,n,\ell) \in O(\min\{m,n\}^2 \max\{m,n\} + mn\ell)$.

This is the standard dense Gaussian-elimination bound for one shared coefficient matrix and ℓ right-hand sides. Next is the workhorse for our solution strategy, allowing us to lift subproblem solutions to solutions of the whole problem.

Problem LinearEqualsAffine($\mathbb{F}$).

Given: $\mathbb{F}$-vector spaces U, W, a linear map $L : U \to W$, an affine space (Θ, V), and an affine map $N : \Theta \to W$.
Return: A computational encoding of the solution set $\{(u,\theta) \in U \times \Theta : L(u) = N(\theta)\}$.

Proposition 3.14. The solution set of LinearEqualsAffine($\mathbb{F}$) is either empty or an affine subspace of $U \times \Theta$.

Proof. Define $F : U \times \Theta \to W$ by $F(u,\theta) := L(u) - N(\theta)$. Let $M : V \to W$ be the underlying linear map of N. Then for $(u', v) \in U \times V$,

$$F\big((u,\theta) + (u',v)\big) = L(u+u') - N(\theta + v) = F(u,\theta) + \big(L(u') - M(v)\big),$$

so F is affine with underlying linear map $(u',v) \mapsto L(u') - M(v)$. The solution set is $F^{-1}(\{0\})$, which is empty or an affine subspace by Lemma 3.11. □

Proposition 3.15. Let $\rho : W \to W/\operatorname{im}(L)$ be the quotient map and define the *feasible parameters*

$$\Theta_{\text{feas}} := \{\theta \in \Theta : \rho(N(\theta)) = 0\}.$$

Then (u,θ) satisfies $L(u) = N(\theta)$ if and only if $\theta \in \Theta_{\text{feas}}$ and $u \in u_0(\theta) + \operatorname{Ker}(L)$ for some fixed $u_0(\theta)$ such that $L(u_0(\theta)) = N(\theta)$.

Proof. For fixed θ, solvability is $N(\theta) \in \operatorname{im}(L)$, which is exactly $\rho(N(\theta)) = 0$. If $u_0(\theta)$ is one solution, then for any other u, $L(u) = N(\theta)$ if and only if $L(u - u_0(\theta)) = 0$ if and only if $u - u_0(\theta) \in \operatorname{Ker}(L)$. □

Proposition 3.16. Given a coordinatized instance of LinearEqualsAffine($\mathbb{F}$), with

$$U = \mathbb{F}^{n\times\ell}, \qquad W = \mathbb{F}^{m\times\ell}, \qquad L(X) = MX, \qquad M \in \mathbb{F}^{m\times n},$$

and with affine-frame coordinates $\theta \in \mathbb{F}^p$ for Θ and $N(\theta) = N_0 + \sum_{i=1}^p \theta_i N_i$, having $N_i \in W$, where

$$\kappa := \dim \operatorname{Ker}(M^T), \qquad q := \dim \Theta_{\text{feas}} \text{ if } \Theta_{\text{feas}} \neq \varnothing, \text{ and } 0 \text{ otherwise.}$$

The problem is solvable in cost

$$\mathtt{T}_{p2}(m,n,\ell,p,q,\kappa) \in O\Big(\underbrace{mn^2 + p\kappa m\ell + (\kappa\ell)p^2}_{\text{feasible parameters}} + \underbrace{mn^2 + n^2\ell}_{\text{kernel of } L} + \underbrace{(q+1)p\,m\ell + mn^2 + (q+1)mn\ell}_{\text{affine frame solve}}\Big).$$

Proof. The term is the cost to solve for feasible parameters. Compute a basis $w_1, \ldots, w_\kappa$ of $\operatorname{Ker}(M^\mathsf{T})$ in $O(mn^2)$. Since $\operatorname{Ker}(M^\mathsf{T}) = \operatorname{im}(M)^\perp$ under the standard pairing,

$$x \in \operatorname{im}(M) \iff \langle w_i, x\rangle = 0 \text{ for all } i \in [\kappa].$$

Let C be the matrix with rows $w_1, \ldots, w_\kappa$. Applying columnwise to $N(\theta)$ means $\rho(N(\theta)) = 0$ exactly when $CN(\theta) = 0$. Recall $N(\theta)$ is coordinatized as $N(\theta) = N_0 + \sum_{i=1}^p \theta_i N_i$. Thus we solve the inhomogeneous system $A_0 + \sum_{i=1}^p \theta_i A_i = 0$ with $A_i := CN_i$. Hence the three costs in the feasible-parameters term are $O(mn^2)$ to compute w_i, $O(p\kappa m\ell)$ to form A_i, and $O((\kappa\ell)p^2)$ to solve the $(\kappa\ell) \times p$ system.

The second term is to compute a basis for the kernel of L. Compute a basis $u_1, \ldots, u_\nu$ of $\operatorname{Ker}(M)$ in $O(mn^2)$, where $\nu := \dim \operatorname{Ker}(M)$. Since $L(X) = MX$ acts columnwise, a basis for $\operatorname{Ker}(L)$ is given by the matrices $u_i e_j^\mathsf{T}$ for $i \in [\nu]$ and $j \in [\ell]$. This costs $O(\nu n\ell) \subseteq O(n^2\ell)$.

The third term is to solve on an affine frame of the feasible parameters. We evaluate N on this affine frame of $(q+1)$ points in $O((q+1)p\,m\ell)$, row-reduce M once in $O(mn^2)$, and solve the $(q+1)$ right-hand sides with that reduction in $O((q+1)mn\ell)$. □

Example 3.17. In the following example, we have $m = 3, n = 2, p = 2$, and $\ell = 2$. And set

$$M = \begin{bmatrix} 1 & 2 \\ 0 & 1 \\ 0 & 0 \end{bmatrix} \in \mathbb{F}^{3\times 2}, \qquad N : \mathbb{F}^2 \to \mathbb{F}^{3\times 2}, \qquad N(\theta_1,\theta_2) = N_0 + \theta_1 N_1 + \theta_2 N_2,$$

$$N_0 = \begin{bmatrix} 2 & 3 \\ 1 & 4 \\ -1 & -2 \end{bmatrix}, \qquad N_1 = \begin{bmatrix} 1 & 0 \\ -1 & 0 \\ 1 & 2 \end{bmatrix}, \qquad N_2 = \begin{bmatrix} 0 & -1 \\ 0 & 2 \\ 1 & 2 \end{bmatrix}.$$

In the three-part process

(1) **Feasible parameters.** Here $\operatorname{im}(M) = \{(u,v,0)^\mathsf{T} : u, v \in \mathbb{F}\}$, so feasibility is exactly vanishing of the third row of $N(\theta_1,\theta_2)$. This is accomplished by the condition $\theta_1 + \theta_2 = 1$. Hence $\Theta_{\text{feas}} = \{(\theta_1,\theta_2) \in \mathbb{F}^2 : \theta_1 + \theta_2 = 1\}$.

(2) **Kernel.** $\operatorname{Ker}(M) = \{0\}$, so $\operatorname{Ker}(L) = \{0\}$.

(3) **Solve on an affine frame of Θ_{feas}.** Take frame points $\theta^{(0)} = (1,0)$ and $\theta^{(1)} = (0,1)$. Solving $MX = N(\theta^{(0)})$ and $MX = N(\theta^{(1)})$ gives

$$X^{(0)} = \begin{bmatrix} 3 & -5 \\ 0 & 4 \end{bmatrix}, \qquad X^{(1)} = \begin{bmatrix} 0 & -10 \\ 1 & 6 \end{bmatrix}.$$

Therefore parametrizing $\theta \in \Theta_{\text{feas}}$ as $\theta = (1-\lambda, \lambda)$, we conclude $X(\theta) = X^{(0)} + \lambda\big(X^{(1)} - X^{(0)}\big)$.

Remark 3.18. If L has a left inverse G, then for each $\theta \in \Theta_{\text{feas}}$, $L(u) = N(\theta)$ has exactly one solution, $u = G(N(\theta))$.

The last linear algebra subroutine solves the compatibility criterion that arises when lifting subproblem solutions.

Problem AffinePullback($\mathbb{F}$).

Given: $\mathbb{F}$-vector spaces U, V, W and affine maps $f : U \to W$, $g : V \to W$.
Return: A computational encoding of $P := \{(u, v) \in U \times V : f(u) = g(v)\}$.

The set P is either empty or an affine subspace of $U \times V$, as $P = h^{-1}(\{0\})$ for the affine map $h(u, v) := f(u) - g(v)$. A verification of the affine nature of h: if L_f and L_g are the underlying linear maps of f and g, then $h((u, v) + (u', v')) = f(u + u') - g(v + v') = f(u) + L_f(u') - (g(v) + L_g(v')) = h(u, v) + (L_f(u') - L_g(v'))$.

Proposition 3.19. Given a coordinate instance of AffinePullback($\mathbb{F}$) with

$$U = \mathbb{F}^d, \quad V = \mathbb{F}^e, \quad W = \mathbb{F}^m, \quad f(u) = Au + \alpha, \quad g(v) = Bv + \beta,$$

given by $A \in \mathbb{F}^{m \times d}$ and $B \in \mathbb{F}^{m \times e}$, it is solvable in cost $\mathtt{T}_{p1}(m, d + e, 1)$.

Proof. The condition $f(u) = g(v)$ is the block linear system $\begin{bmatrix} A & -B \end{bmatrix} \begin{bmatrix} u & v \end{bmatrix}^\mathsf{T} = \beta - \alpha$, which is one instance of LinSolve($\mathbb{F}$). □

3.3. **Simultaneous Sylvester Equations for** $\mathrm{Adj}(T)$**.** We work towards Theorem 3.21 in this section. Throughout, a, b, c, r, s will be natural numbers, and we use the notation

$$\mathcal{X} := \mathbb{F}^{a \times r}, \qquad \mathcal{Y} := \mathbb{F}^{s \times b},$$

to denote vector spaces. This notation may have subscripts to indicate restricted solution spaces. As an example, for $I \subseteq [a]$, $\mathcal{X}_I$ is the space $\mathbb{F}^{|I| \times r}$.

Problem SimultaneousSylvesterSystem.

Given: Tensors $R \in \mathbb{F}^{r \times b \times c}$, $S \in \mathbb{F}^{a \times s \times c}$, and $T \in \mathbb{F}^{a \times b \times c}$.
Return: Either $\varnothing$, or the affine subspace

$$\{(X, Y) \in \mathcal{X} \times \mathcal{Y} : (\forall k \in [c])\ X\, R[:, :, k] + S[:, :, k]\, Y = T[:, :, k]\}\,.$$

This problem is named as a generalization to the single matrix Sylvester equation $XA + BX = C$ [Syl84]. It has a naive solution using LinSolve($\mathbb{F}$) with $m = abc$, $n = ar + bs$, and $\ell = 1$. If $R \in \mathbb{F}^{a \times b \times c}$ and we take $S = -R$ and $T = 0$, then solutions give the adjoint for the tensor R.

Example 3.20. Below is an example of laying out the equations for SimultaneousSylvesterSystem to solve using LinSolve($\mathbb{F}$). Take $a = 2$, $b = 2$, $c = 2$, $r = 3$, and $s = 4$, with slices

$$R[:, :, 1] = \begin{bmatrix} 1 & 5 \\ 0 & -1 \\ 2 & 11 \end{bmatrix}, \qquad R[:, :, 2] = \begin{bmatrix} 0 & 1 \\ 1 & 1 \\ -1 & 1 \end{bmatrix},$$

$$S[:, :, 1] = \begin{bmatrix} 1 & 0 & 3 & 0 \\ 1 & 1 & 10 & -1 \end{bmatrix}, \qquad S[:, :, 2] = \begin{bmatrix} 0 & 1 & 7 & -1 \\ 1 & -1 & -4 & 1 \end{bmatrix},$$

$$T[:, :, 1] = \begin{bmatrix} 1 & 19 \\ -11 & 4 \end{bmatrix}, \qquad T[:, :, 2] = \begin{bmatrix} -10 & 4 \\ 8 & 2 \end{bmatrix}.$$

The corresponding augmented linear system is shown in Figure 1.

x_{11}	x_{12}	x_{13}	x_{21}	x_{22}	x_{23}	y_{11}	y_{21}	y_{31}	y_{41}	y_{12}	y_{22}	y_{32}	y_{42}	t
1	0	2				1	0	3	0					1
0	1	−1				0	1	7	−1					−10
5	−1	11								1	0	3	0	19
1	1	1								0	1	7	−1	4
			1	0	2	1	1	10	−1					−11
			0	1	−1	1	−1	−4	1					8
			5	−1	11					1	1	10	−1	4
			1	1	1					1	−1	−4	1	2

FIGURE 1. An augmented matrix for the Simultaneous Sylvester System in Example 3.20.

We describe an alternative approach, culminating in the following.

Theorem 3.21. For cubic ($n \times n \times n$) tensors, under the regularity conditions (established in Definition 3.32), SimultaneousSylvesterSystem is solved in the same cost as the full deterministic verification. This is $O(n^4)$ cost, as compared to the naive $O(n^7)$ cost by solving as a linear system of equations.

Example 3.22. We use the following running example throughout Section 3.3. Work over $\mathbb{F} = \mathbb{Q}$, with $a = b = r = s = 2$ and $c = 2$. Define

$$R[:,:,1] = S[:,:,1] = T[:,:,1] = \mathrm{diag}(1,0),$$
$$R[:,:,2] = S[:,:,2] = \mathrm{diag}(0,1), \quad T[:,:,2] = \mathrm{diag}(0,2).$$

The full solution family is 2-dimensional, parametrized by $\alpha, \beta \in \mathbb{F}$ as $X = \mathrm{diag}(\alpha, \beta)$, and $Y = \mathrm{diag}(1-\alpha, 2-\beta)$.

Our strategy for SimultaneousSylvesterSystem is to solve smaller systems with a subset of the constraints, and then lift them. We start with the following.

Problem DoubleRestricted.

Given: Tensors $R \in \mathbb{F}^{r\times b\times c}$, $S \in \mathbb{F}^{a\times s\times c}$, $T \in \mathbb{F}^{a\times b\times c}$, and index sets $I \subseteq [a]$, $J \subseteq [b]$ with $|I| = a'$ and $|J| = b'$.

Return: Return $\varnothing$, or the affine subspace

$$\mathcal{S}_{\mathrm{DR}} := \{(X_I, Y_J) \in \mathcal{X}_I \times \mathcal{Y}_J : (\forall k \in [c])\ X_I R[:,J,k] + S[I,:,k] Y_J = T[I,J,k]\}.$$

To solve, vectorizing yields an instance of LinSolve($\mathbb{F}$) which costs $\mathtt{T}_{p1}(m = a'b'c,\ n = a'r + b's,\ \ell = 1)$. We refer to this cost as $\mathtt{T}_{\mathrm{DR}}(a,b,c,r,s,a',b')$, or when clear, just $\mathtt{T}_{\mathrm{DR}}$. By Proposition 3.12, the returned solution set is affine.

Example 3.23. In the running example, take $I = J = \{1\}$. Then $a' = b' = 1$, and

$$X_I = \begin{bmatrix} x_1 & x_2 \end{bmatrix} \in \mathbb{F}^{1\times 2}, \qquad Y_J = \begin{bmatrix} y_1 \\ y_2 \end{bmatrix} \in \mathbb{F}^{2\times 1}.$$

Here DoubleRestricted gives the constraint $x_1 + y_1 = 1$, with no other constraints on (x_2, y_2). Hence

$$\mathcal{S}_{\mathrm{DR}} = \left\{ \left(\begin{bmatrix} \alpha & \beta \end{bmatrix}, \begin{bmatrix} 1-\alpha \\ \gamma \end{bmatrix} \right) : \alpha, \beta, \gamma \in \mathbb{F} \right\}.$$

Problem RowRestricted.

Given: Tensors $R \in \mathbb{F}^{r\times b\times c}$, $S \in \mathbb{F}^{a\times s\times c}$, $T \in \mathbb{F}^{a\times b\times c}$, and index sets $I \subseteq [a]$, $J \subseteq [b]$ with $|I| = a'$ and $|J| = b'$.

Return: Either $\varnothing$, or the affine subspace

$$\mathcal{S}_{\mathrm{R}} := \{(X_I, Y) \in \mathcal{X}_I \times \mathcal{Y} : (\forall k \in [c])\ X_I R[:,:,k] + S[I,:,k] Y = T[I,:,k]\}.$$

Proposition 3.24. For fixed $I \subseteq [a]$ and $J \subseteq [b]$, RowRestricted is solved by solving DoubleRestricted and then LinearEqualsAffine($\mathbb{F}$), with cost

$$O\big(\mathtt{T}_{p1}(m = a'b'c,\ n = a'r + b's,\ \ell = 1) + d\,c\,a'r(b-b') + \mathtt{T}_{p2}(m = ca',\ n = s,\ \ell = b-b',\ p = d,\ q = q_R,\ \kappa = \kappa_R)\big).$$

where $a' = |I|$, $b' = |J|$, d is the dimension of the solution set of DoubleRestricted ($d = \dim \mathcal{S}_{\mathrm{DR}}$), and (q_R, κ_R) are the feasible parameter dimension and the left nullspace dimension parameter respectively in the LinearEqualsAffine($\mathbb{F}$) instance (these three parameters are all 0 if $\varnothing$ is returned)

Proof. **Algorithm. Step 1.** Solve DoubleRestricted. If it returns $\varnothing$, return $\varnothing$ and stop.

Step 2. Set $\widehat{J} := [b] \setminus J$ and define an affine map $N_R : \mathcal{S}_{\mathrm{DR}} \to \mathbb{F}^{(ca')\times(b-b')}$ as well as an element $M_R \in \mathbb{F}^{(ca')\times s}$ by

$$N_R(X_I, Y_J) := \begin{bmatrix} T[I,\widehat{J},1] - X_I R[:,\widehat{J},1] \\ \vdots \\ T[I,\widehat{J},c] - X_I R[:,\widehat{J},c] \end{bmatrix} \in \mathbb{F}^{(ca')\times(b-b')}, \qquad M_R := \begin{bmatrix} S[I,:,1] \\ \vdots \\ S[I,:,c] \end{bmatrix} \in \mathbb{F}^{(ca')\times s} \tag{3.1}$$

Apply LinearEqualsAffine($\mathbb{F}$) to

$$U = \mathcal{Y}_{\widehat{J}}, \quad W = \mathbb{F}^{(ca')\times(b-b')},$$
$$(L_R : U \to W, \quad \widehat{Y} \mapsto M_R\widehat{Y}), \quad \Theta = \mathcal{S}_{\mathrm{DR}}, \quad N = N_R.$$

This returns either $\varnothing$, in which case our algorithm stops, or an affine subspace $\mathcal{A}_R \subseteq \mathcal{S}_{\mathrm{DR}} \times \mathcal{Y}_{\widehat{J}}$ of pairs $((X_I, Y_J), \widehat{Y})$ satisfying $M_R\widehat{Y} = N_R(X_I, Y_J)$.

Step 3. For each $((X_I, Y_J), \widehat{Y}) \in \mathcal{A}_R$, form the unique $Y \in \mathcal{Y}$ such that $Y[:,J] = Y_J$ and $Y[:,\widehat{J}] = \widehat{Y}$. Return the affine subspace of $\mathcal{X}_I \times \mathcal{Y}$ obtained by sending each $((X_I, Y_J), \widehat{Y}) \in \mathcal{A}_R$ to (X_I, Y).

Timing. Step 1 is one call to DoubleRestricted, costing $\mathtt{T}_{\mathrm{DR}}(a,b,c,r,s,a',b')$. Step 2 has two contributions. Recall $\dim(\mathcal{S}_{\mathrm{DR}}) = d$, and we are given an affine frame $(X_{I,i}, Y_{J,i})_{i=0}^{d}$. Building the resulting coordinatized affine map N_R from Equation (3.1) costs $O\big(d\,c\,a'r(b-b')\big)$, because each of the d matrices has size $a' \times r$, each $R[:,\widehat{J},k]$ has size $r \times (b-b')$, and we compute c multiplications between them.

Invoking Proposition 3.16 on (M_R, N_R) adds $\mathtt{T}_{p2}(m = ca',\ n = s,\ \ell = b-b',\ p = d,\ q = q_R,\ \kappa = \kappa_R)$. Adding the three steps gives the desired cost.

Correctness. Step 1 gives exactly $\mathcal{S}_{\mathrm{DR}}$, i.e. all (X_I, Y_J) that satisfy the (I,J)-restricted equations. Step 2 gives exactly the elements $\widehat{Y}$ for which the combined Y_J and $\widehat{Y}$ satisfy the row restricted equation. Step 3 combines $(Y_J, \widehat{Y})$ into Y, so $\mathcal{S}_{\mathrm{R}}$ is exactly the set of row-restricted solutions. □

Example 3.25. Continue the running example with $I = J = \{1\}$, so $\widehat{J} = \{2\}$. From Example 3.23, write

$$X_I = \begin{bmatrix} \alpha & \beta \end{bmatrix}, \qquad Y_J = \begin{bmatrix} 1-\alpha \\ \gamma \end{bmatrix}.$$

Step 2 of Proposition 3.24 computes

$$M_R = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \qquad N_R(X_I, Y_J) = \begin{bmatrix} 0 \\ -\beta \end{bmatrix}.$$

Solving $M_R\widehat{Y} = N_R(X_I, Y_J)$ forces $\widehat{Y} = \begin{bmatrix} 0 \\ \delta \end{bmatrix}$ and $\beta = 0$. Therefore $\mathcal{S}_{\mathrm{R}} = \left\{ \left(\begin{bmatrix} \alpha & 0 \end{bmatrix}, \begin{bmatrix} 1-\alpha & 0 \\ \gamma & \delta \end{bmatrix} \right) : \alpha, \gamma, \delta \in \mathbb{F} \right\}$.

Next we describe and solve an analogous variant of the row restricted problem, where instead, we restrict the columns of the original tensors.

Problem ColumnRestricted.

Given: Tensors $R \in \mathbb{F}^{r\times b\times c}$, $S \in \mathbb{F}^{a\times s\times c}$, $T \in \mathbb{F}^{a\times b\times c}$, and index sets $I \subseteq [a]$, $J \subseteq [b]$ with $|I| = a'$ and $|J| = b'$.

Return: Return $\varnothing$, or the affine subspace

$$\mathcal{S}_{\mathrm{C}} := \{(X, Y_J) \in \mathcal{X} \times \mathcal{Y}_J : (\forall k \in [c])\ XR[:,J,k] + S[:,:,k]Y_J = T[:,J,k]\}.$$

Proposition 3.26. For fixed $I \subseteq [a]$ and $J \subseteq [b]$, ColumnRestricted is solved by solving DoubleRestricted and then LinearEqualsAffine($\mathbb{F}$), in cost

$$O\big(\mathtt{T}_{p1}(m = a'b'c,\ n = a'r + b's,\ \ell = 1) + d\,c\,(a-a')sb' + \mathtt{T}_{p2}(m = cb',\ n = r,\ \ell = a-a',\ p = d,\ q = q_C,\ \kappa = \kappa_C)\big).$$

where $a' = |I|$, $b' = |J|$, d is the dimension of the solution set of DoubleRestricted ($d = \dim \mathcal{S}_{\mathrm{DR}}$), and (q_C, κ_C) are the feasible parameter dimension and the left nullspace dimension parameter respectively in the LinearEqualsAffine($\mathbb{F}$) instance (these three parameters are all 0 if $\varnothing$ is returned)

Proof. **Algorithm. Step 1.** Solve DoubleRestricted. If it returns $\varnothing$, return $\varnothing$ and stop.

Step 2. Set $\widehat{I} := [a] \setminus I$ and define an affine map $N_C : \mathcal{S}_{\mathrm{DR}} \to \mathbb{F}^{(cb')\times(a-a')}$ as well as an element $M_C \in \mathbb{F}^{(cb')\times r}$ by

$$N_C(X_I, Y_J) := \begin{bmatrix} (T[\widehat{I}, J, 1] - S[\widehat{I}, :, 1]Y_J)^{\mathsf{T}} \\ \vdots \\ (T[\widehat{I}, J, c] - S[\widehat{I}, :, c]Y_J)^{\mathsf{T}} \end{bmatrix} \in \mathbb{F}^{(cb')\times(a-a')}, \qquad M_C := \begin{bmatrix} (R[:, J, 1])^{\mathsf{T}} \\ \vdots \\ (R[:, J, c])^{\mathsf{T}} \end{bmatrix} \in \mathbb{F}^{(cb')\times r}.$$

Apply LinearEqualsAffine($\mathbb{F}$) to

$$U = \mathbb{F}^{r\times(a-a')}, \quad W = \mathbb{F}^{(cb')\times(a-a')},$$
$$L_C : U \to W, \quad \widehat{X} \mapsto M_C\widehat{X}, \quad \Theta = \mathcal{S}_{\mathrm{DR}}, \quad N = N_C.$$

This returns either $\varnothing$, in which case our algorithm stops, or an affine subspace $\mathcal{A}_C \subseteq \mathcal{S}_{\mathrm{DR}} \times \mathbb{F}^{r\times(a-a')}$ of pairs $((X_I, Y_J), \widehat{X})$ satisfying $M_C\widehat{X} = N_C(X_I, Y_J)$.

Step 3. For each $((X_I, Y_J), \widehat{X}) \in \mathcal{A}_C$, form the unique $X \in \mathcal{X}$ such that $X[I, :] = X_I$ and $X[\widehat{I}, :] = \widehat{X}^{\mathsf{T}}$. Return $\mathcal{S}_{\mathrm{C}}$, the affine subspace of $\mathcal{X} \times \mathcal{Y}_J$ obtained by sending each $((X_I, Y_J), \widehat{X}) \in \mathcal{A}_C$ to (X, Y_J).

Timing. The same cost accounting as Proposition 3.24. Step 1 costs $\mathtt{T}_{\mathrm{DR}}(a, b, c, r, s, a', b')$, Step 2 has cost in $O\big(d\,c\,(a-a')sb' + \mathtt{T}_{p2}(m = cb',\ n = r,\ \ell = a-a',\ p = d,\ q = q_C,\ \kappa = \kappa_C)\big)$. Adding the three steps gives the desired cost.

Correctness. Same as Proposition 3.24. □

Example 3.27. Continue the running example with $I = J = \{1\}$, so $\widehat{I} = \{2\}$. From Example 3.23, write

$$X_I = \begin{bmatrix} \alpha & \beta \end{bmatrix}, \qquad Y_J = \begin{bmatrix} 1-\alpha \\ \gamma \end{bmatrix}.$$

Step 2 of Proposition 3.26 gives

$$M_C = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \qquad M_C\widehat{X} = N_C(X_I, Y_J) = \begin{bmatrix} 0 \\ -\gamma \end{bmatrix},$$

with $\widehat{X}$ as the unknown. Solving gives $\widehat{X} = \begin{bmatrix} 0 \\ \epsilon \end{bmatrix}$ and $\gamma = 0$. Therefore $\mathcal{S}_{\mathrm{C}} = \left\{ \left(\begin{bmatrix} \alpha & \beta \\ 0 & \epsilon \end{bmatrix}, \begin{bmatrix} 1-\alpha \\ 0 \end{bmatrix} \right) : \alpha, \beta, \epsilon \in \mathbb{F} \right\}$.

The final problem is the following.

Problem LiftRestricted.

Given: Solutions to RowRestricted and ColumnRestricted.
Return: Solution to SimultaneousSylvesterSystem.

Solving RowRestricted and ColumnRestricted gives $\mathcal{S}_{\mathrm{R}} \subseteq \mathcal{X}_I \times \mathcal{Y}$ and $\mathcal{S}_{\mathrm{C}} \subseteq \mathcal{X} \times \mathcal{Y}_J$. For $z_R = (X_I, Y) \in \mathcal{S}_{\mathrm{R}}$ and $z_C = (X, Y_J) \in \mathcal{S}_{\mathrm{C}}$, define projections

$$\pi_J : \mathcal{X}_I \times \mathcal{Y} \to \mathcal{X}_I \times \mathcal{Y}_J, \qquad \pi_J(z_R) := (X_I, Y[:, J]),$$
$$\pi_I : \mathcal{X} \times \mathcal{Y}_J \to \mathcal{X}_I \times \mathcal{Y}_J, \qquad \pi_I(z_C) := (X[I, :], Y_J),$$

for which we want restricted solutions to agree on.

Proposition 3.28. Let $\mathcal{S}_{\mathrm{R}} \subseteq \mathcal{X}_I \times \mathcal{Y}$, and $\mathcal{S}_{\mathrm{C}} \subseteq \mathcal{X} \times \mathcal{Y}_J$ be the solutions of RowRestricted and ColumnRestricted. Let the pullback of the projection maps be defined as

$$P := \{(z_R, z_C) \in \mathcal{S}_{\mathrm{R}} \times \mathcal{S}_{\mathrm{C}} : \pi_J(z_R) = \pi_I(z_C)\},$$

the output map be defined as

$$\Psi : P \to \mathcal{X} \times \mathcal{Y}, \qquad \Psi\big((X_I, Y), (X, Y_J)\big) := (X, Y),$$

and the error map be defined as

$$\mathrm{Err} : \mathcal{X} \times \mathcal{Y} \to \mathbb{F}^{a\times b\times c}, \qquad \mathrm{Err}(X,Y) := \big(XR[:,:,k] + S[:,:,k]Y - T[:,:,k]\big)_{k\in[c]}.$$

Then $\{\Psi(z) : z \in P \text{ and } \mathrm{Err}(\Psi(z)) = 0\}$ is the solution set to LiftRestricted.

Proof. The restricted pairs agreeing from $\mathcal{S}_\mathrm{R}$ and $\mathcal{S}_\mathrm{C}$ are exactly P. For each $z \in P$, $\Psi(z)$ is a candidate solution. This candidate is in the solution set of SimultaneousSylvesterSystem exactly when its residual is zero. □

Proposition 3.29. Let $\mathcal{S}_\mathrm{R}, \mathcal{S}_\mathrm{C}$ be the outputs of RowRestricted and ColumnRestricted. If $\mathcal{S}_\mathrm{R} \neq \varnothing$ and $\mathcal{S}_\mathrm{C} \neq \varnothing$, let

$$d := \max\{\dim \mathcal{S}_\mathrm{R}, \dim \mathcal{S}_\mathrm{C}\}, \qquad P := \{(z_R, z_C) \in \mathcal{S}_\mathrm{R} \times \mathcal{S}_\mathrm{C} : \pi_J(z_R) = \pi_I(z_C)\},$$
$$\tau := \dim P \text{ if } P \neq \varnothing, \text{ and } 0 \text{ otherwise.}$$

Otherwise set both d and τ equal to 0. Then LiftRestricted is solvable in cost

$$\mathtt{T}_{\mathrm{lift}}(a,b,c,r,s,a',b',d,\tau) \in O\big((a'r + b's)d^2 + (\tau+1)\,c\,a\,b\,(r+s) + \mathtt{T}_{p1}(m = abc,\ n = \tau,\ \ell = 1)\big).$$

Proof. **Algorithm.**

If either RowRestricted or ColumnRestricted returns $\varnothing$, then we return $\varnothing$ and stop.

Step 1 (compute the candidates). From the outputs of RowRestricted and ColumnRestricted, we get the affine frames $(z_{R,0}, \ldots, z_{R,d_R}) \subseteq \mathcal{S}_\mathrm{R}$, and $(z_{C,0}, \ldots, z_{C,d_C}) \subseteq \mathcal{S}_\mathrm{C}$. Solve AffinePullback($\mathbb{F}$) with

$$U = \mathbb{F}^{d_R}, \quad V = \mathbb{F}^{d_C}, \quad W = \mathcal{X}_I \times \mathcal{Y}_J,$$
$$f : U \to W,\ \lambda \mapsto \pi_J(\chi_R(\lambda)), \quad g : V \to W,\ \mu \mapsto \pi_I(\chi_C(\mu)),$$

where $\chi_R : \mathbb{F}^{d_R} \to \mathcal{S}_\mathrm{R}$ and $\chi_C : \mathbb{F}^{d_C} \to \mathcal{S}_\mathrm{C}$ are the coordinate maps. If $P = \varnothing$, return $\varnothing$. Otherwise output the frame $(p_0, \ldots, p_\tau) \subseteq \mathcal{S}_\mathrm{R} \times \mathcal{S}_\mathrm{C}$ from the coordinate pullback solutions.

Step 2 (compute the error map on a frame). For each $t \in \{0, \ldots, \tau\}$, compute

$$(X_t, Y_t) := \Psi(p_t), \qquad E_t := \mathrm{Err}(X_t, Y_t) \in \mathbb{F}^{a\times b\times c}, \qquad e_t := \mathrm{vec}(E_t) \in \mathbb{F}^{abc}.$$

If $\tau = 0$, this means at most one solution exists. We check if $e_0 = 0$. In the case $e_0 \neq 0$, return $\varnothing$. When $e_0 = 0$, the solution is the unique point $\Psi(p_0)$.

Else, $\tau > 0$, so we solve the inhomogeneous linear system $Mx = -e_0$ for $M := \begin{bmatrix} e_1 - e_0 & \cdots & e_\tau - e_0 \end{bmatrix} \in \mathbb{F}^{abc\times\tau}$ in the frame coordinates via LinSolve($\mathbb{F}$). If it returns $\varnothing$, return $\varnothing$. Otherwise output the frame of solutions in $\mathcal{X} \times \mathcal{Y}$ by applying Ψ to the coordinate solutions of the linear system.

Timing. Step 1 is one instance of AffinePullback($\mathbb{F}$), with $d = d_R$, $e = d_C$, and $m = a'r + sb'$. Under the assumptions that d_R, d_C are $O(d)$, Proposition 3.19 takes $O(d^2(a'r + sb'))$ operations. Evaluating Err at one point costs $O(c\,a\,b\,(r+s))$, so all $\tau+1$ evaluations cost $O((\tau+1)\,c\,a\,b\,(r+s))$. Solving $Mx = -e_0$ is one call to LinSolve($\mathbb{F}$) with $m = abc$, $n = \tau$, and $\ell = 1$, so it costs $\mathtt{T}_{p1}(m = abc,\ n = \tau,\ \ell = 1)$. □

Example 3.30. From Examples 3.25 and 3.27, a compatible pair in the pullback has the form $X = \begin{bmatrix} \alpha & 0 \\ 0 & \epsilon \end{bmatrix}$, $Y = \begin{bmatrix} 1-\alpha & 0 \\ 0 & \delta \end{bmatrix}$, for $\alpha, \epsilon, \delta \in \mathbb{F}$. Step 2 checks

$$\mathrm{Err}(X,Y) = \big(XR[:,:,k] + S[:,:,k]Y - T[:,:,k]\big)_{k\in[2]},$$

and the remaining condition is $\epsilon + \delta = 2$. Therefore LiftRestricted returns the same as the naive solution after renaming ϵ to β, namely $X = \mathrm{diag}(\alpha, \beta)$, and $Y = \mathrm{diag}(1-\alpha, 2-\beta)$ for $\alpha, \beta \in \mathbb{F}$.

Now we combine the above to solve SimultaneousSylvesterSystem.

Theorem 3.31. Let $I \subseteq [a]$ and $J \subseteq [b]$. Then SimultaneousSylvesterSystem is solvable in cost

$$\mathtt{T}_{\mathrm{DR}} + \mathtt{T}_{\mathrm{RR}} + \mathtt{T}_{\mathrm{CR}} + \mathtt{T}_{\mathrm{lift}},$$

where $\mathtt{T}_{\mathrm{DR}}$ is the cost for DoubleRestricted, $\mathtt{T}_{\mathrm{RR}}, \mathtt{T}_{\mathrm{CR}}$ are the costs from RowRestricted and ColumnRestricted, not including the double restricted solve, and $\mathtt{T}_{\mathrm{lift}}$ the cost for LiftRestricted.

Proof. By correctness of DoubleRestricted, we obtain the doubly-restricted solution family for (I, J). By Propositions 3.24 and 3.26, the row- and column-restricted families computed from that data are exactly the solutions of RowRestricted and ColumnRestricted. Then LiftRestricted is applied to these two families. Its correctness means we solve SimultaneousSylvesterSystem. □

An end-to-end example can be found in Example B.1.

While Theorem 3.31 gives the correct solution for any choice of I and J, we describe a class of problems on which this solve-and-lift algorithm performs best.

Definition 3.32. The *regularity* condition that is the hypothesis of Theorem 3.21 is when

(1) For some fixed integer n, that $a, b, c, r, s \in O(n)$.
(2) There are choices $I \subseteq [a], J \subseteq [b]$ such that $a' := |I| \in O(1)$, $b' := |J| \in O(1)$, and $a'b'c \geq a'r + b's$.
(3) The restricted systems have the property that $\dim \mathcal{S}_{\text{DR}} \in O(1)$, and that $N_R(\mathcal{S}_{\text{DR}}) \subseteq \text{im}(M_R)$, $N_C(\mathcal{S}_{\text{DR}}) \subseteq \text{im}(M_C)$, and $\text{Ker}(M_R) = \text{Ker}(M_C) = \{0\}$.

The above assumptions are plausible in the cubic case ($n \times n \times n$ tensors) with generic entries. The matrices M_R and M_C have dimensions $M_R \in \mathbb{F}^{(a'c)\times s}$ and $M_C \in \mathbb{F}^{(b'c)\times r}$ respectively. In the cubic regime, both the number of rows and columns are both in $O(n)$, and for generic slice data one expects full column rank, so $\text{Ker}(M_R) = \text{Ker}(M_C) = \{0\}$. The requrements $N_R(\mathcal{S}_{\text{DR}}) \subseteq \text{im}(M_R)$ and $N_C(\mathcal{S}_{\text{DR}}) \subseteq \text{im}(M_C)$ together with $\text{Ker}(M_R) = \text{Ker}(M_C) = \{0\}$ assert that all double restricted solutions can be uniquely lifted.

Proof of Theorem 3.21. Assume the regularity conditions. For the doubly-restricted step, the regularity hypothesis gives $a'b'c \geq a'r + b's$, so $\mathtt{T}_{\text{DR}} \in O\big((a'b'c)(a'r + b's)^2\big)$, which is $O(n^3)$ since $a', b' \in O(1)$ and $c, r, s \in O(n)$. Since $N_R(\mathcal{S}_{\text{DR}}) \subseteq \text{im}(M_R)$ and $N_C(\mathcal{S}_{\text{DR}}) \subseteq \text{im}(M_C)$, the feasible parameter spaces in the two lifting instances are all of $\mathcal{S}_{\text{DR}}$. Recalling that q_R and q_C denote their dimensions, so we have $q_R = q_C = \dim \mathcal{S}_{\text{DR}} \in O(1)$. For the row and column restricted steps, after excluding the already counted doubly-restricted solve, the remaining costs are

stage	$\mathtt{T}_{p2}$ **parameters**	**additional term**	**cost**
RowRestricted	$(m, n, \ell, p, q, \kappa) = (ca',\ s,\ b - b',\ d_{\text{DR}},\ q_R,\ \kappa_R)$	$d_{\text{DR}}\, c\, a'\, r\, (b - b')$	$O(n^3)$
ColRestricted	$(m, n, \ell, p, q, \kappa) = (cb',\ r,\ a - a',\ d_{\text{DR}},\ q_C,\ \kappa_C)$	$d_{\text{DR}}\, c\, (a - a') s b'$	$O(n^3)$

In each case above, $m, n, \ell \in O(n)$, $p, q \in O(1)$, and $\kappa \in O(n)$. Recall from Proposition 3.16 that $\mathtt{T}_{p2}$ has cost bounded by sum of the terms mn^2, $p\kappa m\ell$, $(\kappa\ell)p^2$, $n^2\ell$, $(q+1)pm\ell$, and $(q+1)mn\ell$. In both the row and column restricted case, $mn^2,\ p\kappa m\ell,\ n^2\ell,\ (q+1)mn\ell \in O(n^3)$, and $(\kappa\ell)p^2,\ (q+1)pm\ell \in O(n^2)$, so $\mathtt{T}_{p2} \in O(n^3)$. Each displayed additional term is also $O(n^3)$, so the cost of each stage is $O(n^3)$, as reflected in the cost column. Therefore $\mathtt{T}_{\text{RR}}, \mathtt{T}_{\text{CR}} \in O(n^3)$.

For the lift step, we have by assumption $\dim \mathcal{S}_{\text{R}} = \dim \mathcal{S}_{\text{C}} = \dim \mathcal{S}_{\text{DR}} = \tau \in O(1)$. Recalling that $d = \max\{\dim \mathcal{S}_{\text{R}}, \dim \mathcal{S}_{\text{C}}\}$, Proposition 3.29 gives

$$\mathtt{T}_{\text{lift}}(a, b, c, r, s, a', b', d, \tau) \in O\big((a'r + b's)d^2 + (\tau + 1)\, c\, a\, b\, (r + s) + \mathtt{T}_{p1}(m = abc,\ n = \tau,\ \ell = 1)\big).$$

Here $(a'r + b's)d^2 \in O(n)$, $(\tau + 1)\, c\, a\, b\, (r + s) \in O(n^4)$, and $\mathtt{T}_{p1}(m = abc,\ n = \tau,\ \ell = 1) \in O(n^3)$, so $\mathtt{T}_{\text{lift}} \in O(n^4)$. This gives the total solve-and-lift cost as $O(n^4)$.

Full deterministic verification checks all abc constraints and has cost $O(abc(r + s)) = O(n^4)$. Therefore SimultaneousSylvesterSystem is solved in the same order of arithmetic complexity as full deterministic verification. □

See Appendix A.1 for performance benchmarking of the solution strategy outlined by Theorem 3.21 as compared to solving one large linear system.

3.4. **Derivation Equations for** $\mathrm{Der}(T)$**.** In this section, we work towards Theorem 3.33 with a similar strategy as the above subsection. Throughout, a, b, c, r, s, t will be natural numbers, and we denote the spaces

$$\mathcal{X} := \mathbb{F}^{a\times r}, \qquad \mathcal{Y} := \mathbb{F}^{s\times b}, \qquad \mathcal{Z} := \mathbb{F}^{t\times c}.$$

This notation may have subscripts to indicate restricted solution spaces. As an example, for $I \subseteq [a]$, $\mathcal{X}_I$ is the space $\mathbb{F}^{|I|\times r}$. For a matrix $Z \in \mathbb{F}^{t\times c}$, define T^Z, the outer action of Z on T, by specifying its slices for $k \in [c]$ as

$$(T^Z)[:,:,k] := \sum_{j\in[t]} T[:,:,j]\, Z[j,k].$$

For $K \subseteq [c]$ with $|K| = c'$ and $Z_K := Z[:,K] \in \mathbb{F}^{t\times c'}$, we likewise write $T^{Z_K} \in \mathbb{F}^{a\times b\times c'}$, with slices indexed by $[c']$ in the induced order on K.

Problem DerivationSystem.

Given: Tensors $R \in \mathbb{F}^{r\times b\times c}$, $S \in \mathbb{F}^{a\times s\times c}$, $T \in \mathbb{F}^{a\times b\times t}$.
Return: Either $\varnothing$, or the affine subspace

$$\left\{(X,Y,Z) \in \mathcal{X}\times\mathcal{Y}\times\mathcal{Z} : (\forall k \in [c])\ X\,R[:,:,k] + S[:,:,k]\,Y = (T^Z)[:,:,k]\right\}.$$

This problem has a naive solution using LinSolve($\mathbb{F}$) with $m = abc$, $n = ar + bs + ct$, and $\ell = 1$. If $T \in \mathbb{F}^{a\times b\times c}$ and we take $r = a$, $s = b$, $t = c$, and $R = S = T$, then solutions to the equation correspond to the derivation algebra of T. We describe an alternative approach, culminating in the following.

Theorem 3.33. For cubic ($n \times n \times n$) tensors, under the regularity conditions (established in Definition 3.38), DerivationSystem is solved in cost $O(n^{4.5})$. This is an additional $n^{1/2}$ overhead compared to the deterministic verification cost of $O(n^4)$, as compared to the naive $O(n^7)$ cost by solving as a linear system of equations.

As in the above section, we solve the derivation system by solving on a subset of constraints, and then lifting those solutions to the full system.

Problem TripleRestrictedDer.

Given: Tensors $R \in \mathbb{F}^{r\times b\times c}$, $S \in \mathbb{F}^{a\times s\times c}$, $T \in \mathbb{F}^{a\times b\times t}$ and index sets $I \subseteq [a]$, $J \subseteq [b]$, and $K = \{k_1 < \cdots < k_{c'}\} \subseteq [c]$, with $|I| = a'$, $|J| = b'$, and $|K| = c'$.
Return: Either $\varnothing$, or the affine subspace

$$\mathcal{D}_{I,J,K} := \left\{(X_I, Y_J, Z_K) \in \mathcal{X}_I\times\mathcal{Y}_J\times\mathcal{Z}_K : (\forall j \in [c'])\ X_I\,R[:,J,k_j] + S[I,:,k_j]\,Y_J = (T^{Z_K})[I,J,j]\right\}.$$

This is a dense linear system with $ra' + sb' + tc'$ unknowns and $a'b'c'$ equations. Hence it is solvable in cost

$$\mathtt{T}_{p1}(m = a'b'c',\ n = ra' + sb' + tc',\ \ell = 1). \tag{3.2}$$

We now solve three restricted systems which give solution families for X, Y, and Z, respectively.

Problem ColDepthRestrictedDer.

Given: Tensors $R \in \mathbb{F}^{r\times b\times c}$, $S \in \mathbb{F}^{a\times s\times c}$, $T \in \mathbb{F}^{a\times b\times t}$, and index sets $I \subseteq [a]$, $J \subseteq [b]$, and $K = \{k_1 < \cdots < k_{c'}\} \subseteq [c]$ with $|I| = a'$, $|J| = b'$, and $|K| = c'$.
Return: Either $\varnothing$, or the affine subspace

$$\mathcal{D}_{[a],J,K} := \left\{(X, Y_J, Z_K) \in \mathcal{X}\times\mathcal{Y}_J\times\mathcal{Z}_K : (\forall j \in [c'])\ X\,R[:,J,k_j] + S[:,:,k_j]\,Y_J = (T^{Z_K})[:,J,j]\right\}.$$

Problem RowDepthRestrictedDer.

Given: Tensors $R \in \mathbb{F}^{r\times b\times c}$, $S \in \mathbb{F}^{a\times s\times c}$, $T \in \mathbb{F}^{a\times b\times t}$, and index sets $I \subseteq [a]$, $J \subseteq [b]$, and $K = \{k_1 < \cdots < k_{c'}\} \subseteq [c]$ with $|I| = a'$, $|J| = b'$, and $|K| = c'$.
Return: Either $\varnothing$, or the affine subspace

$$\mathcal{D}_{I,[b],K} := \left\{(X_I, Y, Z_K) \in \mathcal{X}_I\times\mathcal{Y}\times\mathcal{Z}_K : (\forall j \in [c'])\ X_I\,R[:,:,k_j] + S[I,:,k_j]\,Y = (T^{Z_K})[I,:,j]\right\}.$$

Problem RowColRestrictedDer.

Given: Tensors $R \in \mathbb{F}^{r\times b\times c}$, $S \in \mathbb{F}^{a\times s\times c}$, $T \in \mathbb{F}^{a\times b\times t}$, and index sets $I \subseteq [a]$, $J \subseteq [b]$, $K \subseteq [c]$ with $|I| = a'$, $|J| = b'$, and $|K| = c'$.

Return: Either $\varnothing$, or the affine subspace

$$\mathcal{D}_{I,J,[c]} := \left\{(X_I, Y_J, Z) \in \mathcal{X}_I \times \mathcal{Y}_J \times \mathcal{Z} : (\forall k \in [c])\ X_I\, R[:, J, k] + S[I, :, k]\, Y_J = (T^Z)[I, J, k]\right\}.$$

Proposition 3.34. Each of ColDepthRestrictedDer, RowDepthRestrictedDer, and RowColRestrictedDer are solved by TripleRestrictedDer followed by LinearEqualsAffine($\mathbb{F}$).

Proof. All three restricted problems are solved by first computing $\mathcal{D}_{I,J,K}$ using TripleRestrictedDer (this is only done once and the solution is shared), and then LinearEqualsAffine($\mathbb{F}$) to find the missing piece.

We detail RowDepthRestrictedDer. Let $\widehat{J} := [b] \setminus J$, and write $K = \{k_1 < \cdots < k_{c'}\}$. Define the affine map $N_R : \mathcal{D}_{I,J,K} \to \mathbb{F}^{(a'c')\times(b-b')}$ and $M_R \in \mathbb{F}^{a'c'\times s}$ by

(3.3)

$$N_R(X_I, Y_J, Z_K) := \begin{bmatrix} (T^{Z_K})[I, \widehat{J}, 1] - X_I\, R[:, \widehat{J}, k_1] \\ \vdots \\ (T^{Z_K})[I, \widehat{J}, c'] - X_I\, R[:, \widehat{J}, k_{c'}] \end{bmatrix} \in \mathbb{F}^{(a'c')\times(b-b')}, \qquad M_R := \begin{bmatrix} S[I, :, k_1] \\ \vdots \\ S[I, :, k_{c'}] \end{bmatrix} \in \mathbb{F}^{(a'c')\times s}.$$

Then, for each $(X_I, Y_J, Z_K) \in \mathcal{D}_{I,J,K}$, solve $M_R\widehat{Y} = N_R(X_I, Y_J, Z_K)$ as an instance of LinearEqualsAffine($\mathbb{F}$). The solution triple (X_I, Y, Z_K) is formed by combining Y_J and $\widehat{Y}$, in a fashion identical to the Sylvester case.

ColDepthRestrictedDer is solved similarly with matrices N_C and M_C, while RowColRestrictedDer requires some new notation to set up the system. Write $\widehat{K} = \{\ell_1 < \cdots < \ell_{c-c'}\} = [c] \setminus K$. Define a matrix $M_D \in \mathbb{F}^{a'b'\times t}$ by specifying its jth column $M_D[:, j] (j \in [t])$ as $\mathrm{vec}(T[I, J, j]) \in \mathbb{F}^{a'b'}$.

And define the affine map $N_D : \mathcal{D}_{I,J,K} \to \mathbb{F}^{a'b'\times(c-c')}$ by specifying the jth column of its output:

$$N_D(X_I, Y_J, Z_K)[:, j] := \mathrm{vec}\big(X_I\, R[:, J, \ell_j] + S[I, :, \ell_j]\, Y_J\big) \qquad (j \in [c - c'])$$

Now solving for $\widehat{Z} \in \mathbb{F}^{t\times(c-c')}$ in $M_D\widehat{Z} = N_D(X_I, Y_J, Z_K)$ as an instance of LinearEqualsAffine($\mathbb{F}$) concludes the solution. □

Proposition 3.35. Let $d := \dim \mathcal{D}_{I,J,K}$ and let (q_R, κ_R), (q_C, κ_C), and (q_Z, κ_Z) be the corresponding feasible parameter dimension / left nullspace dimension parameter pairs in the three LinearEqualsAffine($\mathbb{F}$) instances solving Proposition 3.34. Write $\mathtt{T}_{\mathrm{TR}} := \mathtt{T}_{p1}(m = a'b'c',\ n = a'r + sb' + tc',\ \ell = 1)$. Then

stage	step 1	step 2	step 3
RowDepth	$\mathtt{T}_{\mathrm{TR}}$	$d\,c'\,a'(b-b')\,(r+t)$	$\mathtt{T}_{p2}(m = a'c', n = s, \ell = b-b', p = d, q = q_R, \kappa = \kappa_R)$
ColDepth	$\mathtt{T}_{\mathrm{TR}}$	$d\,c'\,(a-a')\,b'\,(s+t)$	$\mathtt{T}_{p2}(m = b'c', n = r, \ell = a-a', p = d, q = q_C, \kappa = \kappa_C)$
RowCol	$\mathtt{T}_{\mathrm{TR}}$	$d\,(c-c')\,a'b'\,(r+s)$	$\mathtt{T}_{p2}(m = a'b', n = t, \ell = c-c', p = d, q = q_Z, \kappa = \kappa_Z)$

Summing each stage gives the cost of the corresponding restricted problem.

Proof. Each stage's cost is the sum of the contributions from the three steps. The first step is the shared triple restricted solve, the second is cost to build the affine map, and third the cost of the corresponding LinearEqualsAffine($\mathbb{F}$) instance.

We explain the step 2 cost for RowDepthRestrictedDer. Using Equation (3.3), we get for each $j \in [c']$, $X_I \in \mathbb{F}^{a'\times r}$ multiplies $R[:, \widehat{J}, k_j] \in \mathbb{F}^{r\times(b-b')}$, and $T^{Z_K}[I, \widehat{J}, j]$ is an $a' \times (b-b')$ block formed from $Z_K \in \mathbb{F}^{t\times c'}$ and the t slices of T. Hence each block costs $a'(b-b')(r+t)$, with c' blocks per frame point and d frame points, giving $d\,c'\,a'(b-b')(r+t)$. The ColDepthRestrictedDer and RowColRestrictedDer bounds are obtained in the same way from their stacked block dimensions, and summing across gives the stated total cost. □

Problem LiftRestrictedDer.

Given: Solutions to ColDepthRestrictedDer, RowDepthRestrictedDer, and RowColRestrictedDer.
Return: The solution to DerivationSystem.

The overall strategy is the same as the solution of LiftRestricted. Each lifted solution space has a restriction map to $\mathcal{D}_{I,J,K}$, for which we want the overall solution to agree on.

$$\begin{aligned}
\rho_X &: \mathcal{D}_{[a],J,K} \to \mathcal{D}_{I,J,K}, & (X, Y_J, Z_K) &\mapsto (X[I,:], Y_J, Z_K)\\
\rho_Y &: \mathcal{D}_{I,[b],K} \to \mathcal{D}_{I,J,K}, & (X_I, Y, Z_K) &\mapsto (X_I, Y[:,J], Z_K),\\
\rho_Z &: \mathcal{D}_{I,J,[c]} \to \mathcal{D}_{I,J,K}, & (X_I, Y_J, Z) &\mapsto (X_I, Y_J, Z[:,K]).
\end{aligned}$$

We first form the 2-way pullback as an instance of AffinePullback($\mathbb{F}$). Solving $Q_{XY} := \{(u,v) \in \mathcal{D}_{[a],J,K} \times \mathcal{D}_{I,[b],K} : \rho_X(u) = \rho_Y(v)\}$ gives compatible partial solution triples (X, Y, Z_K). Then as a second pullback calculation, solve for $Q := \{((u,v),w) \in Q_{XY} \times \mathcal{D}_{I,J,[c]} : \rho_X(u) = \rho_Y(v) = \rho_Z(w)\}$.

Define $\Psi_{\mathrm{Der}} : Q \to \mathcal{X} \times \mathcal{Y} \times \mathcal{Z}$ by $\Psi_{\mathrm{Der}}\big(((X, Y_J, Z_K), (X_I, Y, Z_K)), (X_I, Y_J, Z)\big) = (X, Y, Z)$, which is the derivation analogue of the map Ψ from Proposition 3.28. What remains is to enforce $\mathrm{Err} = 0$ on the candidate triples arising from Q. Define

$$\mathrm{Err} : \mathcal{X} \times \mathcal{Y} \times \mathcal{Z} \to \mathbb{F}^{a \times b \times c}, \qquad \mathrm{Err}(X, Y, Z)[:,:,k] := X\, R[:,:,k] + S[:,:,k]\, Y - (T^Z)[:,:,k],$$

and solve for the candidate solutions satisfying $\mathrm{Err} = 0$ in a linear system.

An end-to-end example can be found in Example B.2.

Proposition 3.36. Let $\mathcal{D}_{I,J,K}$, $\mathcal{D}_{[a],J,K}$, $\mathcal{D}_{I,[b],K}$, and $\mathcal{D}_{I,J,[c]}$ be the outputs of TripleRestrictedDer, ColDepthRestrictedDer, and RowDepthRestrictedDer, and RowColRestrictedDer respectively. If $\mathcal{D}_{I,J,K} \neq \varnothing$, $\mathcal{D}_{[a],J,K} \neq \varnothing$, $\mathcal{D}_{I,[b],K} \neq \varnothing$, and $\mathcal{D}_{I,J,[c]} \neq \varnothing$, let Q_{XY} and Q be the two pullbacks built in LiftRestrictedDer. Let

$$\begin{aligned}
&d := \dim \mathcal{D}_{I,J,K}, \quad d_X := \dim \mathcal{D}_{[a],J,K}, \quad d_Y := \dim \mathcal{D}_{I,[b],K}, \quad d_Z := \dim \mathcal{D}_{I,J,[c]},\\
&m_{\mathrm{TR}} := a'r + sb' + tc', \quad d_{XY} := \dim Q_{XY}, \quad \tau := \dim Q \text{ if } Q \neq \varnothing, \text{ and } 0 \text{ otherwise.}
\end{aligned}$$

Otherwise let each of d, d_X, d_Y, d_Z, m_{TR}, d_{XY}, and τ equal to 0.

Then LiftRestrictedDer is solvable in cost

$$\begin{aligned}
\mathtt{T}_{\mathrm{lift}} \in O\big(&\mathtt{T}_{p1}(m = m_{\mathrm{TR}},\ n = d_X + d_Y,\ \ell = 1) + \mathtt{T}_{p1}(m = m_{\mathrm{TR}},\ n = d_{XY} + d_Z,\ \ell = 1)\\
&+ (\tau + 1)\, abc(r + s + t) + \mathtt{T}_{p1}(m = abc,\ n = \tau,\ \ell = 1)\big).
\end{aligned}$$

Proof. If any of the four input spaces is $\varnothing$, return $\varnothing$ and stop. The first two $\mathtt{T}_{p1}$ summands are the costs of the two applications of AffinePullback($\mathbb{F}$). In both pullbacks, the common codomain is the ambient core space $\mathcal{X}_I \times \mathcal{Y}_J \times \mathcal{Z}_K \cong \mathbb{F}^{m_{\mathrm{TR}}}$ obtained by vectorizing the common restricted triple. After the second pullback, we evaluate the Err map on an affine frame $q_0, \ldots, q_\tau \in Q$ via Ψ_{Der}. Each evaluation costs $O(abc(r + s + t))$, since for each $k \in [c]$ we must compute $X\, R[:,:,k]$, $S[:,:,k]\, Y$, and $(T^Z)[:,:,k]$. Finally, solving for the coordinates of Q that satisfy $\mathrm{Err}(\Psi_{\mathrm{Der}}(q)) = 0$ requires an application of LinSolve($\mathbb{F}$) with $m = abc$, $n = \tau$, and $\ell = 1$. □

Theorem 3.37. Fix $I \subseteq [a]$, $J \subseteq [b]$, and $K \subseteq [c]$. Then DerivationSystem is solvable in cost

$$\mathtt{T}_{\mathrm{core}} + \mathtt{T}_{\mathrm{CD}} + \mathtt{T}_{\mathrm{RD}} + \mathtt{T}_{\mathrm{RC}}, + \mathtt{T}_{\mathrm{lift}}$$

where $\mathtt{T}_{\mathrm{core}}$ is the TripleRestrictedDer cost from Equation (3.2), $\mathtt{T}_{\mathrm{CD}}$, $\mathtt{T}_{\mathrm{RD}}$, and $\mathtt{T}_{\mathrm{RC}}$ are the three restricted lift costs from Proposition 3.35, not counting solving the triple restricted system, and $\mathtt{T}_{\mathrm{lift}}$ is the cost of computing compatible solutions from Proposition 3.36.

Proof. Same reasoning as Theorem 3.31. □

While Theorem 3.37 gives the correct solution for any choice of I, J and K, we describe a class of problems on which this solve-and-lift algorithm performs best.

Definition 3.38. The *regularity* condition that is the hypothesis of Theorem 3.33 is when

(1) For some fixed integer n, that $a, b, c, r, s, t \in O(n)$.
(2) There are choices $I \subseteq [a], J \subseteq [b], K \subseteq [c]$ such that $a' := |I| \in O(n^{1/2})$, $b' := |J| \in O(n^{1/2})$, $c' := |K| \in O(n^{1/2})$, and $a'b'c' \geq a'r + sb' + tc'$.
(3) The restricted systems have the property that $\dim \mathcal{D}_{I,J,K} \in O(1)$, $N_R(\mathcal{D}_{I,J,K}) \subseteq \mathrm{im}(M_R)$, $N_C(\mathcal{D}_{I,J,K}) \subseteq \mathrm{im}(M_C)$, $N_D(\mathcal{D}_{I,J,K}) \subseteq \mathrm{im}(M_D)$, and $\mathrm{Ker}(M_R) = \mathrm{Ker}(M_C) = \mathrm{Ker}(M_D) = \{0\}$.

The above assumptions are plausible in the cubic case ($n \times n \times n$ tensors) with generic entries. The matrices M_R, M_C, and M_D have dimensions $M_R \in \mathbb{F}^{(a'c') \times s}$, $M_C \in \mathbb{F}^{(b'c') \times r}$, and $M_D \in \mathbb{F}^{(a'b') \times t}$ respectively. In the cubic regime, both the number of rows and columns are in $O(n)$, and $a'b'c' \geq a'r + sb' + tc'$ implies $a'c' \geq s$, $b'c' \geq r$, and $a'b' \geq t$. Hence each matrix has at least as many rows as columns, and for generic slice data one expects full column rank, so $\mathrm{Ker}(M_R) = \mathrm{Ker}(M_C) = \mathrm{Ker}(M_D) = \{0\}$. The requirements $N_R(\mathcal{D}_{I,J,K}) \subseteq \mathrm{im}(M_R)$, $N_C(\mathcal{D}_{I,J,K}) \subseteq \mathrm{im}(M_C)$, and $N_D(\mathcal{D}_{I,J,K}) \subseteq \mathrm{im}(M_D)$ together with $\mathrm{Ker}(M_R) = \mathrm{Ker}(M_C) = \mathrm{Ker}(M_D) = \{0\}$ assert that all triple restricted solutions can be uniquely lifted.

Proof of Theorem 3.33. Assume the regularity conditions. The triple restricted derivation system has cost $\mathtt{T}_{p1}(m = a'b'c',\ n = ra' + sb' + tc',\ \ell = 1)$ by (3.2). The regularity hypothesis gives $a'b'c' \geq ra' + sb' + tc'$. With $a', b', c' \in O(n^{1/2})$ and $r, s, t \in O(n)$, the two matrix dimensions in Equation (3.2) are both in $O(n^{3/2})$. Therefore this step has cost $O(n^{4.5})$. Since $N_R(\mathcal{D}_{I,J,K}) \subseteq \mathrm{im}(M_R)$, $N_C(\mathcal{D}_{I,J,K}) \subseteq \mathrm{im}(M_C)$, and $N_D(\mathcal{D}_{I,J,K}) \subseteq \mathrm{im}(M_D)$, the feasible parameter spaces in the three lifting instances have dimension equal to $\mathcal{D}_{I,J,K}$. Hence $q_R = q_C = q_Z = \dim \mathcal{D}_{I,J,K} \in O(1)$. The three restricted lift families and the two pullback spaces all are uniquely determined by $\mathcal{D}_{I,J,K}$ as well, hence $d = d_X = d_Y = d_Z = d_{XY} = \tau = \dim \mathcal{D}_{I,J,K} \in O(1)$. For the three restricted lift stages, recall from Proposition 3.35 that, after excluding the already counted triple restricted solve, the remaining costs are

stage	$\mathtt{T}_{p2}$ **parameters**	**additional term**	**cost**
ColDepth	$(m, n, \ell, p, q, \kappa) = (b'c',\ r,\ a - a',\ d,\ q_C,\ \kappa_C)$	$d\,c'(a - a')b'(s + t)$	$O(n^3)$
RowDepth	$(m, n, \ell, p, q, \kappa) = (a'c',\ s,\ b - b',\ d,\ q_R,\ \kappa_R)$	$d\,c'a'(b - b')(r + t)$	$O(n^3)$
RowCol	$(m, n, \ell, p, q, \kappa) = (a'b',\ t,\ c - c',\ d,\ q_Z,\ \kappa_Z)$	$d(c - c')a'b'(r + s)$	$O(n^3)$

In each row of the table, $m, n, \ell \in O(n)$, $p, q \in O(1)$, and $\kappa \leq m \in O(n)$. Recall from Proposition 3.16 that $\mathtt{T}_{p2}$ has cost bounded by sum of the terms mn^2, $p\kappa m\ell$, $(\kappa\ell)p^2$, $n^2\ell$, $(q+1)pm\ell$, and $(q+1)mn\ell$. In all three stages, $mn^2,\ p\kappa m\ell,\ n^2\ell,\ (q+1)mn\ell \in O(n^3)$, and $(\kappa\ell)p^2,\ (q+1)pm\ell \in O(n^2)$, so $\mathtt{T}_{p2} \in O(n^3)$. Each additional term is also $O(n^3)$, therefore $\mathtt{T}_{\mathrm{CD}}$, $\mathtt{T}_{\mathrm{RD}}$,and $\mathtt{T}_{\mathrm{RC}}$ each is in $O(n^3)$.

For the final lifting stage, write $m_{\mathrm{TR}} := a'r + sb' + tc'$, so $m_{\mathrm{TR}} \in O(n^{3/2})$. Recall from Proposition 3.36 that

$$\begin{aligned}\mathtt{T}_{\mathrm{lift}} \in O\big(&\mathtt{T}_{p1}(m = m_{\mathrm{TR}},\ n = d_X + d_Y,\ \ell = 1) + \mathtt{T}_{p1}(m = m_{\mathrm{TR}},\ n = d_{XY} + d_Z,\ \ell = 1)\\ &+ (\tau + 1)\,abc(r + s + t) + \mathtt{T}_{p1}(m = abc,\ n = \tau,\ \ell = 1)\big).\end{aligned}$$

Here the first two $\mathtt{T}_{p1}$ terms are $O(n^{3/2})$, the evaluation of the Err function is $O(n^4)$, and the final linear solve ($\mathtt{T}_{p1}$) is $O(n^3)$ by the same bound with $\tau \in O(1)$, so $\mathtt{T}_{\mathrm{lift}} \in O(n^4)$.

Full deterministic verification checks all abc constraints at the cost $O(abc(r + s + t)) = O(n^4)$. The total solve-and-lift cost is dominated by the triple restricted solve cost of $O(n^{4.5})$, giving the stated additional $n^{1/2}$ overhead over verification. □

See Appendix A.2 for performance benchmarking of the solution strategy outlined in Theorem 3.33 as compared to solving one large linear system.

As a reminder, for a bilinear map $b : U \times V \rightarrowtail W$, its left nucleus is $\mathcal{L}(b) = \{(\alpha, \gamma) \in \mathrm{End}(U) \times \mathrm{End}(W) : b \bullet_2 \alpha = b \bullet_0 \gamma\}$. The middle and right nucleus, $\mathcal{M}(b), \mathcal{R}(b)$, are defined in the same way by matching the corresponding axis actions.

In this section, we work towards the below theorem.

Theorem 4.1. Let $s : U_2 \times U_1 \rightarrowtail U_0$ and $t : V_2 \times V_1 \rightarrowtail V_0$ be fully nondegenerate bilinear maps. Then

$$\mathcal{L}(s \otimes t) = \mathcal{L}(s) \boxtimes \mathcal{L}(t), \quad \mathcal{M}(s \otimes t) = \mathcal{M}(s) \boxtimes \mathcal{M}(t), \quad \mathcal{R}(s \otimes t) = \mathcal{R}(s) \boxtimes \mathcal{R}(t).$$

As a result of this theorem, we also have the following.

Corollary 4.2. Let $s : U_2 \times U_1 \rightarrowtail U_0$ and $t : V_2 \times V_1 \rightarrowtail V_0$ be fully nondegenerate bilinear maps. Then $\mathcal{C}(s \otimes t) = \mathcal{C}(s) \boxtimes \mathcal{C}(t)$.

4.1. Previous results.

Proposition 4.3 ([Wil09, Propositions 7.8 and 7.9])**.** Let $d : U \times U \rightarrowtail C$ be a nondegenerate Hermitian C-form and $b : V \times V \rightarrowtail W$ a k-bilinear map. Then $d \otimes b$ is Hermitian and $\mathrm{Adj}(d \otimes b) = \mathrm{Adj}(d) \otimes \mathrm{Adj}(b)$.

In our notation, $\mathrm{Adj}(b) = \mathcal{M}(b)$. In the Hermitian setting, the adjoint ring is faithfully represented on one axis, so the notation $\mathrm{Adj}(d) \otimes \mathrm{Adj}(b)$ is unambiguous. Theorem 4.1 recovers this case via $\mathcal{M}(d \otimes b) = \mathcal{M}(d) \boxtimes \mathcal{M}(b)$.

4.2. Algebra preliminaries.

Definition 4.4. Let U, V be finite-dimensional $\mathbb{F}$-vector spaces, and $t \in U \otimes V$. Define

$$\begin{aligned} \Phi_L &: U \otimes V \to \mathrm{Hom}(V^*, U), & \Phi_L(u \otimes v)(\varphi) &= \varphi(v)u, \\ \Phi_R &: U \otimes V \to \mathrm{Hom}(U^*, V), & \Phi_R(u \otimes v)(\psi) &= \psi(u)v \\ \mathrm{Im}_L(t) &\coloneqq \mathrm{Im}(\Phi_L(t)) \leq U, & \mathrm{Im}_R(t) &\coloneqq \mathrm{Im}(\Phi_R(t)) \leq V. \end{aligned}$$

Proposition 4.5. Let U, V be finite-dimensional $\mathbb{F}$-vector spaces and $t \in U \otimes V$. Then $t \in \mathrm{Im}_L(t) \otimes \mathrm{Im}_R(t) \leq U \otimes V$.

Proof. Let $A \coloneqq \mathrm{Im}_L(t) \leq U$, and $X \coloneqq \mathrm{Im}_R(t) \leq V$. Fix a basis $v_1, \ldots, v_n$ of V with dual basis $\varphi_1, \ldots, \varphi_n$. Then $t = \sum_{j=1}^n \Phi_L(t)(\varphi_j) \otimes v_j \in A \otimes V$, since each $\Phi_L(t)(\varphi_j)$ lies in A.

Fix a basis $u_1, \ldots, u_m$ of U with dual basis $\psi_1, \ldots, \psi_m$. Then $t = \sum_{i=1}^m u_i \otimes \Phi_R(t)(\psi_i) \in U \otimes X$, since each $\Phi_R(t)(\psi_i)$ lies in X. Therefore by Lemma 2.2, $t \in (A \otimes V) \cap (U \otimes X) = A \otimes X$. □

Remark 4.6. Let U, V be finite-dimensional $\mathbb{F}$-vector spaces. Let $\{u_i\}$ be an ordered basis of U and $\{v_j\}$ an ordered basis of V. Then t can be uniquely written as $t = \sum_{i,j} T_{ij}\, u_i \otimes v_j \in U \otimes V$. This identifies t with the matrix $T = (T_{ij})$.

Example 4.7. Let $U = \mathbb{F}^2$ with standard bases e_1, e_2 and $V = \mathbb{F}^3$ with standard basis f_1, f_2, f_3. Let $t = e_1 \otimes f_1 + e_2 \otimes f_1 + e_2 \otimes f_2$. Then t corresponds to the matrix $T = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \end{bmatrix}$.

Definition 4.8. Let U, V be finite-dimensional $\mathbb{F}$-vector spaces and $t \in U \otimes V$. The *rank* of t is the minimum number r such that t can be written as a sum $t = \sum_{i=1}^r u_i \otimes v_i$, for $u_i \in U$, and $v_i \in V$. The sum $\sum_{i=1}^r u_i \otimes v_i$ is called an *minimal rank decomposition* of t.

We'll use the observation that for $t \in U \otimes V$, the rank of t is less than or equal to the minimum of $\dim U$, and $\dim V$, as we can always expand t either in a basis of U or a basis of V.

Lemma 4.9. Let $t = \sum_{i=1}^r u_i \otimes v_i$ in $U \otimes V$, where $r = \mathrm{rank}(t)$. Then $\{u_1, \ldots, u_r\}$ and $\{v_1, \ldots, v_r\}$ are linearly independent, and $\mathrm{Im}_L(t) = \mathrm{span}\{u_1, \ldots, u_r\}$, $\mathrm{Im}_R(t) = \mathrm{span}\{v_1, \ldots, v_r\}$.

Proof. Set $W \coloneqq \operatorname{span}\{u_1, \dots, u_r\}$ and $X \coloneqq \operatorname{span}\{v_1, \dots, v_r\}$. Since $t \in W \otimes V$, the above observation gives $\operatorname{rank}(t) \le \dim W$. Since W is spanned by r vectors, $\dim W \le r$. Together with $\operatorname{rank}(t) = r$, this gives $\dim W = r$, so $\{u_1, \dots, u_r\}$ is linearly independent. An analogous argument gives $\dim X = r$, so $\{v_1, \dots, v_r\}$ is linearly independent.

Extend $v_1, \dots, v_r$ to a basis of V, and let $\varphi_1, \dots, \varphi_r \in V^*$ be the corresponding dual functionals. Then $\Phi_L(t)(\varphi_i) = u_i$, so $W \le \operatorname{Im}_L(t)$.

For the reverse inclusion, we see for all $\varphi \in V^*$, $\Phi_L(t)(\varphi) = \sum_{i=1}^r \varphi(v_i)\, u_i \in W$, so $\operatorname{Im}_L(t) \le W$. Hence $\operatorname{Im}_L(t) = W$. An analogous argument concludes $\operatorname{Im}_R(t) = X$. □

Remark 4.10. Suppose $\sum_{i=1}^r a_i \otimes x_i = \sum_{i=1}^r b_i \otimes y_i$ are two minimal rank decompositions of the same element $t \in U \otimes V$. Fix ordered bases of U and V, and let A, B be matrices whose columns are coordinates of a_i, b_i, and X, Y be matrices whose rows coordinates of x_i, y_i. In these fixed coordinates, we get the equality $AX = BY$. By Lemma 4.9, the columns of A and B are two ordered bases of the same space $\operatorname{Im}_L(t)$, so there is a unique $M \in \mathrm{GL}_r(\mathbb{F})$ such that $B = AM$. Then $AX = BY = AMY$, and since the columns of A are linearly independent, $X = MY$, so $Y = M^{-1}X$.

Example 4.11. Take $\mathbb{F} = \mathbb{F}_5$, $U = V = \mathbb{F}^3$, and $r = 2$. Define in the standard basis

$$A = \left[\begin{array}{c|c} a_1 & a_2 \end{array}\right] = \left[\begin{array}{c|c} 1 & 0 \\ 2 & 1 \\ 0 & 1 \end{array}\right], \quad X = \left[\begin{array}{c} x_1 \\ \hline x_2 \end{array}\right] = \left[\begin{array}{ccc} 1 & 2 & 0 \\ \hline 0 & 1 & 1 \end{array}\right]$$

$$B = \left[\begin{array}{c|c} b_1 & b_2 \end{array}\right] = \left[\begin{array}{c|c} 1 & 2 \\ 0 & 3 \\ 3 & 4 \end{array}\right], \quad Y = \left[\begin{array}{c} y_1 \\ \hline y_2 \end{array}\right] = \left[\begin{array}{ccc} 3 & 2 & 1 \\ \hline 4 & 0 & 2 \end{array}\right].$$

A direct multiplication verifies

$$AX = BY = \begin{bmatrix} 1 & 2 & 0 \\ 2 & 0 & 1 \\ 0 & 1 & 1 \end{bmatrix},$$

meaning $a_1 \otimes x_1 + a_2 \otimes x_2 = b_1 \otimes y_1 + b_2 \otimes y_2$. As per Remark 4.10, we find

$$M = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}, \qquad M^{-1} = \begin{bmatrix} 3 & 1 \\ 4 & 2 \end{bmatrix},$$

satisfying

$$AM = \begin{bmatrix} 1 & 2 \\ 0 & 3 \\ 3 & 4 \end{bmatrix} = B, \qquad M^{-1}X = \begin{bmatrix} 3 & 2 & 1 \\ 4 & 0 & 2 \end{bmatrix} = Y.$$

4.3. **Key reduction.** We'll use Lemma 2.11 to reduce an equality about the image of axis actions to a matrix factorization statement. As a reminder for vector spaces U, V, the map $\kappa : \operatorname{End}(U) \otimes \operatorname{End}(V) \to \operatorname{End}(U \otimes V)$ is the Kronecker isomorphism.

Lemma 4.12. Let $s : U_2 \times U_1 \rightarrowtail U_0$ and $t : V_2 \times V_1 \rightarrowtail V_0$ be fully nondegenerate bilinear maps. Let $K \subseteq \{2, 1, 0\}$ be nonempty, and $w \in \operatorname{Bil}(U_2, U_1; U_0) \otimes \operatorname{Bil}(V_2, V_1; V_0)$ be an element such that for each $k \in K$, there is some $x_k \in \operatorname{End}(U_k \otimes V_k)$ satisfying

$$w = \Xi\big((s \otimes t) \bullet_k x_k\big) \in \operatorname{Bil}(U_2, U_1; U_0) \otimes \operatorname{Bil}(V_2, V_1; V_0).$$

Fix a minimal rank decomposition $w = \sum_{i=1}^r u_i \otimes v_i$. Then for each $k \in K$, there exists $\{\alpha_{k,i} \in \operatorname{End}(U_k)\}_{i=1}^r$ and $\{f_{k,i} \in \operatorname{End}(V_k)\}_{i=1}^r$ such that for all $i \in \{1, \dots, r\}$, $u_i = s \bullet_k \alpha_{k,i}$, $v_i = t \bullet_k f_{k,i}$, and $x_k = \sum_{i=1}^r \kappa_k(\alpha_{k,i} \otimes f_{k,i})$.

Proof. Fix $k \in K$ and any decomposition $x_k = \sum_j \kappa_k(\alpha_j \otimes f_j)$. Equation (2.2) gives $w = \sum_j \varphi(s \bullet_k \alpha_j, t \bullet_k f_j)$, so $w \in \operatorname{Im}(\Lambda_k^s) \otimes \operatorname{Im}(\Lambda_k^t)$, with $\operatorname{Im}_L(w) \le \operatorname{Im}(\Lambda_k^s)$, and $\operatorname{Im}_R(w) \le \operatorname{Im}(\Lambda_k^t)$.

Since $w = \sum_i u_i \otimes v_i$ is a minimal rank decomposition, apply Lemma 4.9 to conclude for each i that the elements $u_i \in \operatorname{Im}_L(w) \le \operatorname{Im}(\Lambda_k^s)$ and $v_i \in \operatorname{Im}_R(w) \le \operatorname{Im}(\Lambda_k^t)$ are images of axis actions on s

and t respectively. Specifically, there exist $\alpha_{k,i} \in \mathrm{End}(U_k)$ and $f_{k,i} \in \mathrm{End}(V_k)$ such that $u_i = s \bullet_k \alpha_{k,i}$, and $v_i = t \bullet_k f_{k,i}$.

Finally, for each k, set $x'_k := \sum_{i=1}^r \kappa_k(\alpha_{k,i} \otimes f_{k,i})$. Since $(s \otimes t) \bullet_k x_k = (s \otimes t) \bullet_k x'_k$ and $\Lambda_k^{s \otimes t}$ is injective, we conclude $x_k = x'_k$. □

4.4. **Proof of Theorem 4.1.**

Proof. We first prove the middle nucleus case, $\mathcal{M}(s \otimes t) = \mathcal{M}(s) \boxtimes \mathcal{M}(t)$.

($\subseteq$) Let $(x, y) \in \mathcal{M}(s \otimes t)$, meaning $(s \otimes t) \bullet_2 x = (s \otimes t) \bullet_1 y$. Set $w := \Xi\big((s \otimes t) \bullet_2 x\big) = \Xi\big((s \otimes t) \bullet_1 y\big) \in \mathrm{Bil}(U_2, U_1; U_0) \otimes \mathrm{Bil}(V_2, V_1; V_0)$. Fix a minimal rank decomposition $w = \sum_{i=1}^r u_i \otimes v_i$. Apply Lemma 4.12 with $K = \{2, 1\}$. We obtain $\alpha_{2,i} =: \alpha_i \in \mathrm{End}(U_2)$, $\alpha_{1,i} =: \beta_i \in \mathrm{End}(U_1)$, $f_{2,i} =: f_i \in \mathrm{End}(V_2)$, $f_{1,i} =: g_i \in \mathrm{End}(V_1)$ such that $s \bullet_2 \alpha_i = u_i = s \bullet_1 \beta_i$, $t \bullet_2 f_i = v_i = t \bullet_1 g_i$, and $x = \sum_{i=1}^r \kappa_2(\alpha_i \otimes f_i)$, $y = \sum_{i=1}^r \kappa_1(\beta_i \otimes g_i)$.

Hence $(\alpha_i, \beta_i) \in \mathcal{M}(s)$ and $(f_i, g_i) \in \mathcal{M}(t)$ for each i, so we have the desired conclusion

$$(x, y) = \sum_{i=1}^r \big((\alpha_i, \beta_i) \boxtimes (f_i, g_i)\big) \in \mathcal{M}(s) \boxtimes \mathcal{M}(t).$$

($\supseteq$) Let $(\alpha, \beta) \in \mathcal{M}(s)$ and $(f, g) \in \mathcal{M}(t)$. Then

$$(s \otimes t) \bullet_2 \kappa_2(\alpha \otimes f) = (s \bullet_2 \alpha) \otimes (t \bullet_2 f) = (s \bullet_1 \beta) \otimes (t \bullet_1 g) = (s \otimes t) \bullet_1 \kappa_1(\beta \otimes g).$$

So $(\alpha, \beta) \boxtimes (f, g) = (\kappa_2(\alpha \otimes f), \kappa_1(\beta \otimes g)) \in \mathcal{M}(s \otimes t)$. Since $\mathcal{M}(s) \boxtimes \mathcal{M}(t)$ is the span of such $\boxtimes$ terms, we get $\mathcal{M}(s) \boxtimes \mathcal{M}(t) \subseteq \mathcal{M}(s \otimes t)$. The left and right nuclei case are analogous.

□

A worked-out nucleus decomposition example is in the appendix (Example D.1).

Now we wrap up this chapter with the proof that $\mathcal{C}(s \otimes t) = \mathcal{C}(s) \boxtimes \mathcal{C}(t)$.

Proof of Corollary 4.2. We prove the direction $\mathcal{C}(s \otimes t) \subseteq \mathcal{C}(s) \boxtimes \mathcal{C}(t)$. Let $(x_2, x_1, x_0) \in \mathcal{C}(s \otimes t)$. Then $(s \otimes t) \bullet_2 x_2 = (s \otimes t) \bullet_1 x_1 = (s \otimes t) \bullet_0 x_0$. Let $w := \Xi\big((s \otimes t) \bullet_2 x_2\big) \in \mathrm{Bil}(U_2, U_1; U_0) \otimes \mathrm{Bil}(V_2, V_1; V_0)$ be the element in common. Fix a minimal rank decomposition $w = \sum_{i=1}^r u_i \otimes v_i$, and apply Lemma 4.12 with $K = \{2, 1, 0\}$, then follow the same argument as the proof of Theorem 4.1.

The direction $\mathcal{C}(s \otimes t) \supseteq \mathcal{C}(s) \boxtimes \mathcal{C}(t)$ is analogous to the proof of $\mathcal{M}(s \otimes t) \supseteq \mathcal{M}(s) \boxtimes \mathcal{M}(t)$ in Theorem 4.1. □

5. Derivation of Product of Bilinear Maps

The main result of this section is the following.

Theorem 5.1. Let s and t be fully nondegenerate bilinear maps. Then

$$\begin{aligned} \mathrm{Der}(s \otimes t) = {} & \mathrm{Der}(s) \boxtimes \mathcal{C}(t) + \mathcal{C}(s) \boxtimes \mathrm{Der}(t) \\ & + \iota_{20}(\mathcal{L}(s) \boxtimes \mathcal{L}(t)) + \iota_{10}(\mathcal{R}(s) \boxtimes \mathcal{R}(t)) + \iota_{21}(\mathcal{M}(s) \boxtimes \mathcal{M}(t)), \end{aligned} \tag{5.1}$$

where $\iota_{20}(\alpha, \gamma) = (\alpha, 0, \gamma)$, $\iota_{10}(\beta, \gamma) = (0, \beta, \gamma)$, and $\iota_{21}(\alpha, \beta) = (-\alpha, \beta, 0)$.

The strategy is similar to the nuclei case, where we relate equations of axis actions on tensor product bilinear maps to equalities in the tensor product of vector spaces.

5.1. **Previous results.**

In this subsection, algebras are finite-dimensional over $\mathbb{F}$, and not necessarily associative. For an algebra A, its derivation algebra refers to

$$\mathcal{D}_{\mathrm{alg}}(A) := \{d \in \mathrm{End}(A) : \forall x, y \in A,\ d(xy) = d(x)y + xd(y)\}.$$

Brešar proved the following.

Theorem 5.2 (Brešar, Theorem 3.1 [Bre17])**.** Let R and S be nonassociative unital algebras. Suppose that either at least one of R and S is finite-dimensional or they are both finitely generated. Then every derivation d of $R \otimes S$ can be written as

$$d = \mathrm{ad}(u) + \sum_{j=1}^{p} \lambda_{z_j} \otimes f_j + \sum_{i=1}^{q} g_i \otimes \lambda_{w_i}, \tag{5.2}$$

where $u \in \mathrm{Nuc}(R) \otimes \mathrm{Nuc}(S)$, $z_j \in Z(R)$, $w_i \in Z(S)$, $f_j \in \mathcal{D}_{\mathsf{alg}}(S)$, and $g_i \in \mathcal{D}_{\mathsf{alg}}(R)$.

Proposition 5.3. Let R and S be unital algebras over $\mathbb{F}$, with multiplication maps μ_R, μ_S. Then Theorem 5.1 implies Theorem 5.2 in the case both R and S are finite dimensional.

Proof. See Appendix C. □

Specialization to Matrix Algebras. Let A be a unital finite-dimensional $\mathbb{F}$-algebra. As a vector space, $M_n(A) \cong \mathbb{F}^{n\times n} \otimes A$, and the multiplication on $M_n(A)$ is defined by $(X \otimes a)(Y \otimes b) = XY \otimes ab$.

Theorem 5.4 (Benkart–Osborn, Corollary 4.9 [BO81])**.** Assume $\mathrm{char}\,\mathbb{F} \neq 2, 3$ and $n > 3$. Then for any unital algebra A, $\mathcal{D}_{\mathsf{alg}}(M_n(A)) = \mathcal{D}_{\mathsf{alg}}(L'_n(A)) = \mathcal{D}_{\mathsf{alg}}(K_n(A)) = (\mathcal{D}_{\mathsf{alg}}(A))^{\#} + \mathrm{ad}_{M_n(N)}$.

Remark 5.5. Benkart–Osborn note that specifically for Corollary 4.9, no restrictions on $\mathrm{char}\,\mathbb{F}$ or n are needed. Notation wise, in Theorem 5.4, N denotes the nucleus of A, $L_n(A)$ is $M_n(A)$ under the Lie product, $L'_n(A) = [L_n(A), L_n(A)]$, and $K_n(A) = L'_n(A)/Z(L'_n(A))$. Also, $(\mathcal{D}_{\mathsf{alg}}(A))^{\#}$ denotes entrywise application of derivations of A. Thus in the notation of this chapter,

$$N = \mathrm{Nuc}(A), \qquad (\mathcal{D}_{\mathsf{alg}}(A))^{\#} = I_n \otimes \mathcal{D}_{\mathsf{alg}}(A), \qquad \mathrm{ad}_{M_n(N)} = \mathrm{ad}(M_n(\mathrm{Nuc}(A))).$$

So Theorem 5.4 proves $\mathcal{D}_{\mathsf{alg}}(M_n(A)) = \big(I_n \otimes \mathcal{D}_{\mathsf{alg}}(A)\big) + \mathrm{ad}(M_n(\mathrm{Nuc}(A)))$.

As a result, we have the following.

Proposition 5.6. Theorem 5.1 implies Theorem 5.4.

Proof. Theorem 5.1 implies Theorem 5.2 in the finite dimensional case by Proposition 5.3, and Theorem 5.2 implies Theorem 5.4 by [Bre17, Cor. 4.2]. □

5.2. **Preliminaries.** The following decomposition definition features prominently. Throughout this subsection, U and V are always finite dimensional $\mathbb{F}$-vector spaces.

Definition 5.7. Let $p, q, r \in U \otimes V$ be nonzero tensors satisfying $p + q = r$. Define

$$A := \mathrm{Im}_L(p), \quad B := \mathrm{Im}_L(q), \quad C := \mathrm{Im}_L(r),$$
$$X := \mathrm{Im}_R(p), \quad Y := \mathrm{Im}_R(q), \quad Z := \mathrm{Im}_R(r),$$

where $p \in A \otimes X$, $q \in B \otimes Y$, $r \in C \otimes Z$ by Proposition 4.5. A (p, q, r) *locally independent unified decomposition* (LIU decomposition) consists of a natural number n, and vectors

$$a_i \in A,\ b_i \in B,\ c_i \in C, \qquad x_i \in X,\ y_i \in Y,\ z_i \in Z,$$

for $i \in [n]$, satisfying the following:

- **Decomposition equality.**
$$p = \sum_{i=1}^{n} a_i \otimes x_i, \qquad q = \sum_{i=1}^{n} b_i \otimes y_i, \qquad r = \sum_{i=1}^{n} c_i \otimes z_i.$$
- **Local equality.** For all $i \in [n]$, $a_i \otimes x_i + b_i \otimes y_i = c_i \otimes z_i$.

Example 5.8. Let $U = V = \mathbb{F}^2$ with bases u_1, u_2 and v_1, v_2. Define $p := u_1 \otimes v_1 + u_1 \otimes v_2$, $q := u_2 \otimes v_1 - u_1 \otimes v_2$, and $r := u_1 \otimes v_1 + u_2 \otimes v_1$. Then $p + q = r$, and $A = \mathrm{span}\{u_1\}$, $B = U$, $C = U$ as subspaces of U, and $X = V$, $Y = V$, $Z = \mathrm{span}\{v_1\}$ as subspaces of V. Now let

$$a_1 = u_1,\ b_1 = u_2,\ c_1 = u_1 + u_2, \qquad x_1 = y_1 = z_1 = v_1,$$
$$a_2 = u_1,\ b_2 = -u_1,\ c_2 = 0, \qquad x_2 = y_2 = v_2,\ z_2 = v_1.$$

Then for each $i \in \{1,2\}$, $a_i \otimes x_i + b_i \otimes y_i = c_i \otimes z_i$, and

$$p = a_1 \otimes x_1 + a_2 \otimes x_2, \quad q = b_1 \otimes y_1 + b_2 \otimes y_2, \quad r = c_1 \otimes z_1 + c_2 \otimes z_2.$$

Thus this is an LIU decomposition of $p + q = r$.

5.3. Existence of a LIU decomposition of the equation $p + q = r$.

Convention 5.9. For the remainder of this subsection, whenever given nonzero tensors $p, q, r \in U \otimes V$ satisfying $p + q = r$, write

$$A \coloneqq \mathrm{Im}_L(p), \quad B \coloneqq \mathrm{Im}_L(q), \quad C \coloneqq \mathrm{Im}_L(r),$$
$$X \coloneqq \mathrm{Im}_R(p), \quad Y \coloneqq \mathrm{Im}_R(q), \quad Z \coloneqq \mathrm{Im}_R(r).$$

It is always the case that $A + B \leq U$ and $X + Y \leq V$. We also assume without loss of generality that $U \coloneqq A + B$, $V \coloneqq X + Y$, because it is always the case that

$$p \in A \otimes X \leq (A + B) \otimes (X + Y) \leq U \otimes V$$
$$q \in B \otimes Y \leq (A + B) \otimes (X + Y) \leq U \otimes V$$
$$r \in \mathrm{Im}_L(r) \otimes \mathrm{Im}_R(r).$$

Furthermore, $\mathrm{Im}_L(r), \mathrm{Im}_R(r)$ satisfies

$$\mathrm{Im}_L(r) = \mathrm{Im}_L(p + q) \leq \mathrm{Im}_L(p) + \mathrm{Im}_L(q) = A + B,$$
$$\mathrm{Im}_R(r) = \mathrm{Im}_R(p + q) \leq \mathrm{Im}_R(p) + \mathrm{Im}_R(q) = X + Y.$$

So we may redefine U to the space $A + B$, and redefine V to the space $X + Y$.

Lemma 5.10. Let U be a $\mathbb{F}$-vector space, and let $A, B, C \leq U$ with $U = A + B$. Define $U^{(1)} \leq U$ by

$$U^{(1)} \coloneqq A \cap B \cap C.$$

Define $U^{(2)} \leq A \cap B$, $U^{(3)} \leq A \cap C$, and $U^{(4)} \leq B \cap C$ by choosing complements

$$A \cap B = U^{(1)} \oplus U^{(2)}, \qquad A \cap C = U^{(1)} \oplus U^{(3)}, \qquad B \cap C = U^{(1)} \oplus U^{(4)}.$$

Define $U^{(5)} \leq A$ and $U^{(6)} \leq B$ by choosing complements

$$A = (U^{(1)} + U^{(2)} + U^{(3)}) \oplus U^{(5)}, \qquad B = (U^{(1)} + U^{(2)} + U^{(4)}) \oplus U^{(6)}.$$

Then

$$U = \bigoplus_{i=1}^{6} U^{(i)}, \qquad A = \bigoplus_{i \in \{1,2,3,5\}} U^{(i)}, \qquad B = \bigoplus_{i \in \{1,2,4,6\}} U^{(i)}.$$

Proof. Let $d_i \coloneqq \dim U^{(i)}$ for $1 \leq i \leq 6$. By construction, $\dim(A \cap B) = d_1 + d_2$, $\dim(A \cap C) = d_1 + d_3$. Using the fact $\dim(X + Y) = \dim X + \dim Y - \dim(X \cap Y)$, with $X = A \cap B$, $Y = A \cap C$, we get

$$\dim\big((A \cap B) + (A \cap C)\big) = (d_1 + d_2) + (d_1 + d_3) - d_1 = d_1 + d_2 + d_3.$$

Since $A = ((A \cap B) + (A \cap C)) \oplus U^{(5)}$, this gives $\dim A = d_1 + d_2 + d_3 + d_5$. Similarly, from $B = ((A \cap B) + (B \cap C)) \oplus U^{(6)}$, $\dim B = d_1 + d_2 + d_4 + d_6$. Finally, as $U = A + B$, $\dim U = \dim(A + B) = \dim A + \dim B - \dim(A \cap B) = \sum_{i=1}^{6} d_i$.

Recall

$$U = A + B = \sum_{i=1}^{6} U^{(i)}, \qquad A = \sum_{i \in \{1,2,3,5\}} U^{(i)}, \qquad B = \sum_{i \in \{1,2,4,6\}} U^{(i)},$$

and the fact that for any finite family of subspaces $\{W_i\}$, $\dim(\sum_i W_i) \leq \sum_i \dim W_i$, with equality if and only if the sum is direct. Applying this to the spaces U, A, B concludes

$$U = \bigoplus_{i=1}^{6} U^{(i)}, \qquad A = \bigoplus_{i \in \{1,2,3,5\}} U^{(i)}, \qquad B = \bigoplus_{i \in \{1,2,4,6\}} U^{(i)}. \qquad \square$$

The above lemma is to prepare for below, which is the key one in proving the existence of a LIU decomposition for the equation $p + q = r$.

Lemma 5.11. Let $p, q, r \in U \otimes V$ be nonzero elements satisfying $p + q = r$, and

$$A := \mathrm{Im}_L(p), \quad B := \mathrm{Im}_L(q), \quad C := \mathrm{Im}_L(r),$$
$$X := \mathrm{Im}_R(p), \quad Y := \mathrm{Im}_R(q), \quad Z := \mathrm{Im}_R(r).$$

Recall via Convention 5.9 that $U = A + B$, $V = X + Y$. With respect to the decomposition in Lemma 5.10, p, q, r satisfy:

$$p \in U^{(1)} \otimes V^{(1)} \oplus U^{(1)} \otimes V^{(\bullet)} \oplus U^{(\bullet)} \otimes V^{(1)} \oplus U^{(2)} \otimes V^{(2)} \oplus U^{(3)} \otimes V^{(3)},$$
$$q \in U^{(1)} \otimes V^{(1)} \oplus U^{(1)} \otimes V^{(\bullet)} \oplus U^{(\bullet)} \otimes V^{(1)} \oplus U^{(2)} \otimes V^{(2)} \oplus U^{(4)} \otimes V^{(4)},$$
$$r \in U^{(1)} \otimes V^{(1)} \oplus U^{(1)} \otimes V^{(\bullet)} \oplus U^{(\bullet)} \otimes V^{(1)} \oplus U^{(3)} \otimes V^{(3)} \oplus U^{(4)} \otimes V^{(4)},$$

where $U^{(\bullet)} := U^{(2)} \oplus \cdots \oplus U^{(6)}$ and $V^{(\bullet)} := V^{(2)} \oplus \cdots \oplus V^{(6)}$.

Proof. Apply Lemma 5.10 to the spaces $A, B, C \leq U$ and to the spaces $X, Y, Z \leq V$.

From the definitions, $A = U^{(1)} \oplus U^{(2)} \oplus U^{(3)} \oplus U^{(5)}$ and $B = U^{(1)} \oplus U^{(2)} \oplus U^{(4)} \oplus U^{(6)}$, and similarly $X = V^{(1)} \oplus V^{(2)} \oplus V^{(3)} \oplus V^{(5)}$ and $Y = V^{(1)} \oplus V^{(2)} \oplus V^{(4)} \oplus V^{(6)}$. Let $\pi_{ij} : U \otimes V \to U^{(i)} \otimes V^{(j)}$ be the block projections and write

$$\pi_{i\bullet} := \sum_{j=1}^{6} \pi_{ij}, \qquad \pi_{\bullet j} := \sum_{i=1}^{6} \pi_{ij}.$$

Hence $(\pi_{4\bullet} + \pi_{6\bullet})(p) = 0$ and $(\pi_{3\bullet} + \pi_{5\bullet})(q) = 0$, while $(\pi_{\bullet 4} + \pi_{\bullet 6})(p) = 0$ and $(\pi_{\bullet 3} + \pi_{\bullet 5})(q) = 0$. For $1 \leq i, j \leq 6$, set $p_{ij} := \pi_{ij}(p)$, $q_{ij} := \pi_{ij}(q)$, and $r_{ij} := \pi_{ij}(r)$.

Applying π_{ij} to $p + q = r$ gives for $(1 \leq i, j \leq 6)$, $p_{ij} + q_{ij} = r_{ij}$. This is a blockwise $p + q = r$ equality: if two entries in a block are zero, then the third is zero. Taking stock we visualize these relations below, where grey means *potentially nonzero*, white means *guaranteed to be zero*.



Row projections from known zero rows. Since $\pi_{3\bullet}(q) = 0$, define $s_3 := \pi_{3\bullet}(p) = \pi_{3\bullet}(r)$. Then $s_3 \in U^{(3)} \otimes X$ and $s_3 \in U^{(3)} \otimes Z$, hence $s_3 \in U^{(3)} \otimes (X \cap Z) = U^{(3)} \otimes \left(V^{(1)} \oplus V^{(3)}\right)$. So row $U^{(3)}$ of p, r has support only in columns $V^{(1)}, V^{(3)}$. The same argument with $U^{(5)}$ gives row $U^{(5)}$ of p, r also in $V^{(1)} \oplus V^{(3)}$.

Since $\pi_{4\bullet}(p) = 0$, define $t_4 := \pi_{4\bullet}(q) = \pi_{4\bullet}(r)$. Then $t_4 \in U^{(4)} \otimes Y$ and $t_4 \in U^{(4)} \otimes Z$, hence $t_4 \in U^{(4)} \otimes (Y \cap Z) = U^{(4)} \otimes \left(V^{(1)} \oplus V^{(4)}\right)$. So row $U^{(4)}$ of q, r has support only in columns $V^{(1)}, V^{(4)}$. The same argument with $U^{(6)}$ gives row $U^{(6)}$ of q, r also in $V^{(1)} \oplus V^{(4)}$.

After these row inferences, the potentially nonzero blocks are:



Column projections from known zero columns. Since $\pi_{\bullet 3}(q) = 0$, define $u_3 := \pi_{\bullet 3}(p) = \pi_{\bullet 3}(r)$. Then $u_3 \in A \otimes V^{(3)}$ and $u_3 \in C \otimes V^{(3)}$, hence $u_3 \in (A \cap C) \otimes V^{(3)} = \left(U^{(1)} \oplus U^{(3)}\right) \otimes V^{(3)}$. So column $V^{(3)}$ of p, r has support only in rows $U^{(1)}, U^{(3)}$.

Since $\pi_{\bullet 4}(p) = 0$, define $u_4 := \pi_{\bullet 4}(q) = \pi_{\bullet 4}(r)$. Then $u_4 \in B \otimes V^{(4)}$ and $u_4 \in C \otimes V^{(4)}$, hence $u_4 \in (B \cap C) \otimes V^{(4)} = \left(U^{(1)} \oplus U^{(4)}\right) \otimes V^{(4)}$. So column $V^{(4)}$ of q, r has support only in rows $U^{(1)}, U^{(4)}$.

Since $\pi_{\bullet 5}(q) = 0$, define $u_5 := \pi_{\bullet 5}(p) = \pi_{\bullet 5}(r)$. Then $u_5 \in (A \cap C) \otimes V^{(5)} = \big(U^{(1)} \oplus U^{(3)}\big) \otimes V^{(5)}$. Combining this with the row-$U^{(3)}$ and row-$U^{(5)}$ restrictions above forces support in column $V^{(5)}$ to lie only in row $U^{(1)}$.

Since $\pi_{\bullet 6}(p) = 0$, define $u_6 := \pi_{\bullet 6}(q) = \pi_{\bullet 6}(r)$. Then $u_6 \in (B \cap C) \otimes V^{(6)} = \big(U^{(1)} \oplus U^{(4)}\big) \otimes V^{(6)}$. Combining this with the row-$U^{(4)}$ and row-$U^{(6)}$ restrictions above forces support in column $V^{(6)}$ to lie only in row $U^{(1)}$.

After these column inferences, the potentially nonzero blocks are:



The $(2,2)$ block relation. Set $v_2 := \pi_{2\bullet}(r)$. By the previous row/column inferences, row $U^{(2)}$ of both p and q is supported in $V^{(1)} \oplus V^{(2)} = X \cap Y$, hence $v_2 \in U^{(2)} \otimes (X \cap Y)$. Also $r \in C \otimes Z$, so $v_2 \in U^{(2)} \otimes Z$. Therefore

$$v_2 \in U^{(2)} \otimes \big((X \cap Y) \cap Z\big) = U^{(2)} \otimes \big(X \cap Y \cap Z\big) = U^{(2)} \otimes V^{(1)}.$$

Thus $r_{22} = 0$, and the block equation gives $p_{22} + q_{22} = 0$.

What remains are the following support patterns, which we give as colored cells below. Here blue is $U^{(1)} \otimes V^{(1)}$, orange is $U^{(1)} \otimes V^{(\bullet)}$, cyan is $U^{(\bullet)} \otimes V^{(1)}$, green is $U^{(2)} \otimes V^{(2)}$, brown is $U^{(3)} \otimes V^{(3)}$, and pink is $U^{(4)} \otimes V^{(4)}$, and blank entries are zero.



This is the claimed support pattern

$$p \in U^{(1)} \otimes V^{(1)} \oplus U^{(1)} \otimes V^{(\bullet)} \oplus U^{(\bullet)} \otimes V^{(1)} \oplus U^{(2)} \otimes V^{(2)} \oplus U^{(3)} \otimes V^{(3)},$$
$$q \in U^{(1)} \otimes V^{(1)} \oplus U^{(1)} \otimes V^{(\bullet)} \oplus U^{(\bullet)} \otimes V^{(1)} \oplus U^{(2)} \otimes V^{(2)} \oplus U^{(4)} \otimes V^{(4)},$$
$$r \in U^{(1)} \otimes V^{(1)} \oplus U^{(1)} \otimes V^{(\bullet)} \oplus U^{(\bullet)} \otimes V^{(1)} \oplus U^{(3)} \otimes V^{(3)} \oplus U^{(4)} \otimes V^{(4)}. \qquad \square$$

Example 5.12. Let $\mathbb{F} = \mathbb{F}_{997}$, let $U = V = \mathbb{F}^6$ with standard bases, and identify $U \otimes V$ with $\mathbb{F}^{6\times 6}$ via $u \otimes v \mapsto [u][v]^T$ where $[u]$ is the column vector of coordinates.

Let $p, q, r \in U \otimes V$ correspond to the 6×6 matrices below.

$$p = \begin{bmatrix} 373 & 831 & 66 & 724 & 159 & 68 \\ 772 & 812 & 372 & 663 & 579 & 209 \\ 155 & 549 & 499 & 757 & 500 & 64 \\ 218 & 733 & 547 & 132 & 342 & 122 \\ 859 & 708 & 25 & 767 & 78 & 648 \\ 992 & 398 & 894 & 611 & 169 & 953 \end{bmatrix} \quad q = \begin{bmatrix} 176 & 355 & 631 & 671 & 759 & 955 \\ 252 & 645 & 740 & 978 & 500 & 638 \\ 308 & 485 & 963 & 867 & 212 & 39 \\ 146 & 830 & 155 & 854 & 264 & 14 \\ 30 & 375 & 193 & 887 & 38 & 692 \\ 107 & 281 & 218 & 640 & 676 & 450 \end{bmatrix} \quad r = \begin{bmatrix} 549 & 189 & 697 & 398 & 918 & 26 \\ 27 & 460 & 115 & 644 & 82 & 847 \\ 463 & 37 & 465 & 627 & 712 & 103 \\ 364 & 566 & 702 & 986 & 606 & 136 \\ 889 & 86 & 218 & 657 & 116 & 343 \\ 102 & 679 & 115 & 254 & 845 & 406 \end{bmatrix}$$

so that $p + q = r$ in $\mathbb{F}^{6\times 6}$. Using Magma, we find change-of-basis matrices M and N so that the new coordinates are adapted to the decomposition $U = \bigoplus_{i=1}^{6} U^{(i)}$ and $V = \bigoplus_{i=1}^{6} V^{(i)}$.

$$M = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 310 & 34 & 931 & 898 & 971 & 773 \\ 104 & 130 & 918 & 780 & 84 & 647 \\ 795 & 834 & 145 & 316 & 939 & 574 \\ 386 & 64 & 11 & 618 & 494 & 754 \\ 732 & 237 & 949 & 667 & 810 & 127 \end{bmatrix}, \quad N = \begin{bmatrix} 1 & 655 & 49 & 786 & 153 & 994 \\ 0 & 757 & 318 & 920 & 875 & 310 \\ 0 & 304 & 386 & 307 & 423 & 363 \\ 0 & 988 & 354 & 652 & 88 & 712 \\ 0 & 780 & 227 & 987 & 917 & 291 \\ 0 & 59 & 338 & 600 & 869 & 429 \end{bmatrix},$$

the transformed matrices $p' = MpN$, $q' = MqN$, $r' = MrN$ have the support patterns as described by Lemma 5.11.

$p' =$

	$V^{(1)}$	$V^{(2)}$	$V^{(3)}$	$V^{(4)}$	$V^{(5)}$	$V^{(6)}$
$U^{(1)}$	373	16	259	0	968	0
$U^{(2)}$	119	905	0	0	0	0
$U^{(3)}$	966	0	123	0	0	0
$U^{(4)}$	0	0	0	0	0	0
$U^{(5)}$	647	0	0	0	0	0
$U^{(6)}$	0	0	0	0	0	0

$q' =$

	$V^{(1)}$	$V^{(2)}$	$V^{(3)}$	$V^{(4)}$	$V^{(5)}$	$V^{(6)}$
$U^{(1)}$	176	828	0	553	0	244
$U^{(2)}$	607	92	0	0	0	0
$U^{(3)}$	0	0	0	0	0	0
$U^{(4)}$	68	0	0	815	0	0
$U^{(5)}$	0	0	0	0	0	0
$U^{(6)}$	970	0	0	0	0	0

$r' =$

	$V^{(1)}$	$V^{(2)}$	$V^{(3)}$	$V^{(4)}$	$V^{(5)}$	$V^{(6)}$
$U^{(1)}$	549	844	259	553	968	244
$U^{(2)}$	726	0	0	0	0	0
$U^{(3)}$	966	0	123	0	0	0
$U^{(4)}$	68	0	0	815	0	0
$U^{(5)}$	647	0	0	0	0	0
$U^{(6)}$	970	0	0	0	0	0

Lemma 5.13. Let $p, q, r \in U \otimes V$ be nonzero tensors satisfying $p + q = r$. Then (p, q, r) admits an LIU decomposition.

Proof. Recall from Convention 5.9 the spaces $A, B, C \leq U$, and $X, Y, Z \leq V$ where $U = A + B$ and $V = X + Y$. Decompose U and V by Lemma 5.10, getting

$$U \otimes V = \bigoplus_{i,j=1}^{6} U^{(i)} \otimes V^{(j)}.$$

Let $\pi_{ij} : U \otimes V \to U^{(i)} \otimes V^{(j)}$ be the corresponding projections. Define

$$p_{11} := \pi_{11}(p), \quad p_{1\bullet} := \sum_{j=2}^{6} \pi_{1j}(p), \quad p_{\bullet 1} := \sum_{i=2}^{6} \pi_{i1}(p), \quad p_{22} := \pi_{22}(p), \quad p_{33} := \pi_{33}(p),$$

and analogously define q_{11}, $q_{1\bullet}$, $q_{\bullet 1}$, q_{22}, q_{44}, and r_{11}, $r_{1\bullet}$, $r_{\bullet 1}$, r_{33}, r_{44}. By Lemma 5.11, all other blocks vanish, so

$$p = p_{11} + p_{1\bullet} + p_{\bullet 1} + p_{22} + p_{33}, \qquad q = q_{11} + q_{1\bullet} + q_{\bullet 1} + q_{22} + q_{44}, \qquad r = r_{11} + r_{1\bullet} + r_{\bullet 1} + r_{33} + r_{44}.$$

Applying the relevant projections to $p + q = r$ gives equations

$$p_{11} + q_{11} = r_{11} \qquad p_{1\bullet} + q_{1\bullet} = r_{1\bullet} \qquad p_{\bullet 1} + q_{\bullet 1} = r_{\bullet 1},$$
$$p_{22} + q_{22} = 0 \qquad p_{33} = r_{33} \qquad q_{44} = r_{44}.$$

We build an LIU decomposition for each block. After, summing the decompositions gives the desired result.

In $U^{(1)} \otimes V^{(1)}$ (blue block), choose bases of $U^{(1)}$ and $V^{(1)}$. Writing p_{11}, q_{11}, r_{11} in that basis, gives coefficientwise local equalities. The $U^{(2)} \otimes V^{(2)}$ (green block), $U^{(3)} \otimes V^{(3)}$ (brown block), and $U^{(4)} \otimes V^{(4)}$ (pink block) cases are specializations of the $U^{(1)} \otimes V^{(1)}$ (blue block) argument.

In $U^{(1)} \otimes V^{(\bullet)}$ (orange block), choose a basis $\{u_m\}_{m=1}^{d}$ of $U^{(1)}$ and write

$$p_{1\bullet} = \sum_{m=1}^{d} u_m \otimes f_m, \qquad q_{1\bullet} = \sum_{m=1}^{d} u_m \otimes g_m, \qquad r_{1\bullet} = \sum_{m=1}^{d} u_m \otimes h_m.$$

Since $p_{1\bullet} + q_{1\bullet} = r_{1\bullet}$, we get that for all m in $[d]$, $u_m \otimes f_m + u_m \otimes g_m = u_m \otimes h_m$. This gives local equalities with a common left factor in $A \cap B \cap C$. The $U^{(\bullet)} \otimes V^{(1)}$ (cyan block) case is analogous. □

Example 5.14. We demonstrate this splitting for Example 5.12. Recall all coefficients are in $\mathbb{F}_{997}$, and let u_i and v_j be basis vectors with $U^{(i)} = \langle u_i \rangle$ and $V^{(j)} = \langle v_j \rangle$.

For $U^{(1)} \otimes V^{(1)}$ (blue block), $p_{11} = 373\, u_1 \otimes v_1$, $q_{11} = 176\, u_1 \otimes v_1$, and $r_{11} = 549\, u_1 \otimes v_1$, so $p_{11} + q_{11} = (373 + 176)\, u_1 \otimes v_1 = 549\, u_1 \otimes v_1 = r_{11}$.

For $U^{(1)} \otimes V^{(\bullet)}$ (orange block), $p_{1\bullet} = u_1 \otimes (16v_2 + 259v_3 + 968v_5)$, and $q_{1\bullet} = u_1 \otimes (828v_2 + 553v_4 + 244v_6)$, so $p_{1\bullet} + q_{1\bullet} = u_1 \otimes (844v_2 + 259v_3 + 553v_4 + 968v_5 + 244v_6) = r_{1\bullet}$.

Likewise, for $U^{(\bullet)} \otimes V^{(1)}$ (cyan block), $p_{\bullet 1} = (119u_2 + 966u_3 + 647u_5) \otimes v_1$, and $q_{\bullet 1} = (607u_2 + 68u_4 + 970u_6) \otimes v_1$, hence $p_{\bullet 1} + q_{\bullet 1} = (726u_2 + 966u_3 + 68u_4 + 647u_5 + 970u_6) \otimes v_1 = r_{\bullet 1}$.

5.4. **Proof of Theorem 5.1.** We apply the LIU decomposition to the images of axis actions and match each block to one summand in Equation (5.1). First, we need to describe the partitioning of the index sets of a LIU decomposition.

Remark 5.15. Given a LIU decomposition of $p + q = r$ produced by Lemma 5.13, with

$$p = \sum_{i=1}^{n} a_i \otimes x_i, \qquad q = \sum_{i=1}^{n} b_i \otimes y_i, \qquad r = \sum_{i=1}^{n} c_i \otimes z_i,$$

and local equalities for all i in $[n]$, $a_i \otimes x_i + b_i \otimes y_i = c_i \otimes z_i$. Let $I_{11},\ I_{1\bullet},\ I_{\bullet 1},\ I_{22},\ I_{33},\ I_{44}$ be the index sets coming from the six blockwise decompositions in the proof of Lemma 5.13. Then these sets give a partition $[n] = I_{11} \sqcup I_{1\bullet} \sqcup I_{\bullet 1} \sqcup I_{22} \sqcup I_{33} \sqcup I_{44}$.

Proof. ($\subseteq$) We prove the inclusion

$$\mathrm{Der}(s \otimes t) \subseteq \mathrm{Der}(s) \boxtimes \mathcal{C}(t) + \mathcal{C}(s) \boxtimes \mathrm{Der}(t) + \iota_{20}(\mathcal{L}(s) \boxtimes \mathcal{L}(t)) + \iota_{10}(\mathcal{R}(s) \boxtimes \mathcal{R}(t)) + \iota_{21}(\mathcal{M}(s) \boxtimes \mathcal{M}(t)).$$

Let $(\delta^{(2)}, \delta^{(1)}, \delta^{(0)}) \in \mathrm{Der}(s \otimes t)$, and define

$$p := \Xi((s \otimes t) \bullet_2 \delta^{(2)}), \qquad q := \Xi((s \otimes t) \bullet_1 \delta^{(1)}), \qquad r := \Xi((s \otimes t) \bullet_0 \delta^{(0)}).$$

Then $p + q = r$ in $\mathrm{Bil}(U_2, U_1; U_0) \otimes \mathrm{Bil}(V_2, V_1; V_0)$.

If $r = 0$, then $(s \otimes t) \bullet_2 \delta^{(2)} = -(s \otimes t) \bullet_1 \delta^{(1)}$, so $(-\delta^{(2)}, \delta^{(1)}) \in \mathcal{M}(s \otimes t)$, and $\Lambda_0^{s \otimes t}$ is injective so $\delta^{(0)} = 0$. Thus $(\delta^{(2)}, \delta^{(1)}, \delta^{(0)}) = (\delta^{(2)}, \delta^{(1)}, 0) \in \iota_{21}(\mathcal{M}(s \otimes t))$. The cases $p = 0$ and $q = 0$ are analogous, obtaining elements of $\iota_{10}(\mathcal{R}(s \otimes t))$ and $\iota_{20}(\mathcal{L}(s \otimes t))$ respectively. Applying Theorem 4.1 proves the desired inclusion in all three cases. Hence for the remainder of the proof, assume p, q, r are all nonzero.

The strategy is to apply Lemma 5.13 to $p + q = r$, obtain a LIU decomposition, then lift the local equalities to the axis operators. As a reminder,

$$A := \mathrm{Im}_L(p), \quad B := \mathrm{Im}_L(q), \quad C := \mathrm{Im}_L(r),$$
$$X := \mathrm{Im}_R(p), \quad Y := \mathrm{Im}_R(q), \quad Z := \mathrm{Im}_R(r),$$

where $p \in A \otimes X$, $q \in B \otimes Y$, and $r \in C \otimes Z$ by Proposition 4.5. Let $(a_i, b_i, c_i, x_i, y_i, z_i)_{i=1}^n$ be the LIU decomposition by applying Lemma 5.13. By Remark 5.15, the LIU decomposition index set $[n]$ is partitioned into $[n] = I_{11} \sqcup I_{1\bullet} \sqcup I_{\bullet 1} \sqcup I_{22} \sqcup I_{33} \sqcup I_{44}$.

index block	local equality	support	corresponding summand
I_{11}	$a_i = b_i = c_i,\ x_i + y_i = z_i$	$U^{(1)} \otimes V^{(1)}$	$\mathcal{C}(s) \boxtimes \mathrm{Der}(t)$
$I_{1\bullet}$	$a_i = b_i = c_i,\ x_i + y_i = z_i$	$U^{(1)} \otimes V^{(\bullet)}$	$\mathcal{C}(s) \boxtimes \mathrm{Der}(t)$
$I_{\bullet 1}$	$x_i = y_i = z_i,\ a_i + b_i = c_i$	$U^{(\bullet)} \otimes V^{(1)}$	$\mathrm{Der}(s) \boxtimes \mathcal{C}(t)$
I_{22}	$a_i = b_i,\ x_i + y_i = 0$	$U^{(2)} \otimes V^{(2)}$	$\iota_{21}(\mathcal{M}(s) \boxtimes \mathcal{M}(t))$
I_{33}	$a_i = c_i,\ x_i = z_i$	$U^{(3)} \otimes V^{(3)}$	$\iota_{20}(\mathcal{L}(s) \boxtimes \mathcal{L}(t))$
I_{44}	$b_i = c_i,\ y_i = z_i$	$U^{(4)} \otimes V^{(4)}$	$\iota_{10}(\mathcal{R}(s) \boxtimes \mathcal{R}(t))$

- $i \in I_{1\bullet}$ ($U^{(1)} \otimes V^{(\bullet)}$) or $i \in I_{11}$ ($U^{(1)} \otimes V^{(1)}$): For each i, the local equality is

$$a_i = b_i = c_i, \qquad x_i + y_i = z_i,$$

with $a_i, b_i, c_i \in A \cap B \cap C$, $x_i \in X$, $y_i \in Y$, $z_i \in Z$. The I_{11} case deserves explaination. The coefficientwise decomposition gives a local equality $\lambda u \otimes v + \mu u \otimes v = \nu u \otimes v$, which we rewrite as $u \otimes (\lambda v) + u \otimes (\mu v) = u \otimes (\nu v)$.

By Lemma 2.11 and Proposition 4.5, we have

$$A \leq \mathrm{Im}(\Lambda_2^s), \quad B \leq \mathrm{Im}(\Lambda_1^s), \quad C \leq \mathrm{Im}(\Lambda_0^s),$$
$$X \leq \mathrm{Im}(\Lambda_2^t), \quad Y \leq \mathrm{Im}(\Lambda_1^t), \quad Z \leq \mathrm{Im}(\Lambda_0^t).$$

So each of $a_i, b_i, c_i, x_i, y_i, z_i$ has a preimage under the corresponding axis action maps. Choose

$$\alpha_i \in \mathrm{End}(U_2), \qquad \beta_i \in \mathrm{End}(U_1), \qquad \gamma_i \in \mathrm{End}(U_0)$$
$$f_i \in \mathrm{End}(V_2), \qquad g_i \in \mathrm{End}(V_1), \qquad h_i \in \mathrm{End}(V_0)$$

such that $a_i = s \bullet_2 \alpha_i,\ b_i = s \bullet_1 \beta_i,\ c_i = s \bullet_0 \gamma_i$, and $x_i = t \bullet_2 f_i,\ y_i = t \bullet_1 g_i,\ z_i = t \bullet_0 h_i$.

Then $(\alpha_i, \beta_i, \gamma_i) \in \mathcal{C}(s)$ and $(f_i, g_i, h_i) \in \mathrm{Der}(t)$, so $(\kappa_2(\alpha_i \otimes f_i), \kappa_1(\beta_i \otimes g_i), \kappa_0(\gamma_i \otimes h_i)) \in \mathcal{C}(s) \boxtimes \mathrm{Der}(t)$.

- $i \in I_{\bullet 1}$ ($U^{(\bullet)} \otimes V^{(1)}$): Here each local equality satisfies $x_i = y_i = z_i$ and $a_i + b_i = c_i$. Choose

$$\begin{aligned} &\alpha_i \in \mathrm{End}(U_2), && \beta_i \in \mathrm{End}(U_1), && \gamma_i \in \mathrm{End}(U_0) \\ &f_i \in \mathrm{End}(V_2), && g_i \in \mathrm{End}(V_1), && h_i \in \mathrm{End}(V_0) \end{aligned}$$

  such that $a_i = s \bullet_2 \alpha_i,\ b_i = s \bullet_1 \beta_i,\ c_i = s \bullet_0 \gamma_i$ and $x_i = t \bullet_2 f_i,\ y_i = t \bullet_1 g_i,\ z_i = t \bullet_0 h_i$. Then $(\alpha_i, \beta_i, \gamma_i) \in \mathrm{Der}(s)$ and $(f_i, g_i, h_i) \in \mathcal{C}(t)$, so $(\kappa_2(\alpha_i \otimes f_i), \kappa_1(\beta_i \otimes g_i), \kappa_0(\gamma_i \otimes h_i)) \in \mathrm{Der}(s) \boxtimes \mathcal{C}(t)$.
- $i \in I_{22}, I_{33}, I_{44}$: these are the nucleus cases with one side equal to 0. We show the I_{22} case, which has $c_i \otimes z_i = 0$. The local equality is $a_i \otimes x_i + b_i \otimes y_i = 0$, with $a_i, b_i \in A \cap B$ and $x_i, y_i \in X \cap Y$. Choose operators

$$\alpha_i \in \mathrm{End}(U_2),\ \beta_i \in \mathrm{End}(U_1), \qquad f_i \in \mathrm{End}(V_2),\ g_i \in \mathrm{End}(V_1)$$

  satisfying $a_i = s \bullet_2 \alpha_i,\ b_i = s \bullet_1 \beta_i$ and $x_i = t \bullet_2 f_i,\ y_i = t \bullet_1 g_i$. Then for all i,

$$(s \otimes t) \bullet_2 \kappa_2(\alpha_i \otimes f_i) = a_i \otimes x_i = -\, b_i \otimes y_i = -\,(s \otimes t) \bullet_1 \kappa_1(\beta_i \otimes g_i).$$

  Hence $\big(-\kappa_2(\alpha_i \otimes f_i),\ \kappa_1(\beta_i \otimes g_i)\big) \in \mathcal{M}(s \otimes t)$, and therefore $\iota_{21}\big(-\kappa_2(\alpha_i \otimes f_i), \kappa_1(\beta_i \otimes g_i)\big) \in \iota_{21}(\mathcal{M}(s \otimes t))$. Applying Theorem 4.1, $\iota_{21}(\mathcal{M}(s \otimes t)) = \iota_{21}(\mathcal{M}(s) \boxtimes \mathcal{M}(t))$, so the operator triple lies in one of the summands on the right-hand side of Equation (5.1). The I_{33} and I_{44} cases are analogous.

For each i, let $(\delta_i^{(2)}, \delta_i^{(1)}, \delta_i^{(0)}) \in \mathrm{End}(U_2 \otimes V_2) \times \mathrm{End}(U_1 \otimes V_1) \times \mathrm{End}(U_0 \otimes V_0)$ be the resulting operator triple constructed corresponding to each block. Summing over i,

$$\begin{aligned} \Xi\left((s \otimes t) \bullet_2 \sum_{i=1}^{n} \delta_i^{(2)}\right) &= p = \Xi((s \otimes t) \bullet_2 \delta^{(2)}), \\ \Xi\left((s \otimes t) \bullet_1 \sum_{i=1}^{n} \delta_i^{(1)}\right) &= q = \Xi((s \otimes t) \bullet_1 \delta^{(1)}), \\ \Xi\left((s \otimes t) \bullet_0 \sum_{i=1}^{n} \delta_i^{(0)}\right) &= r = \Xi((s \otimes t) \bullet_0 \delta^{(0)}). \end{aligned}$$

Since Ξ is an isomorphism and $\Lambda_2^{s \otimes t}, \Lambda_1^{s \otimes t}, \Lambda_0^{s \otimes t}$ are injective, we get

$$\sum_{i=1}^{n} \delta_i^{(2)} = \delta^{(2)}, \qquad \sum_{i=1}^{n} \delta_i^{(1)} = \delta^{(1)}, \qquad \sum_{i=1}^{n} \delta_i^{(0)} = \delta^{(0)}.$$

As each is summing over the same indices, $(\delta^{(2)}, \delta^{(1)}, \delta^{(0)}) = \sum_{i=1}^{n} (\delta_i^{(2)}, \delta_i^{(1)}, \delta_i^{(0)})$. This proves the desired inclusion.

($\supseteq$) We prove $\mathrm{Der}(s \otimes t) \supseteq \mathrm{Der}(s) \boxtimes \mathcal{C}(t) + \mathcal{C}(s) \boxtimes \mathrm{Der}(t) + \iota_{20}(\mathcal{L}(s) \boxtimes \mathcal{L}(t)) + \iota_{10}(\mathcal{R}(s) \boxtimes \mathcal{R}(t)) + \iota_{21}(\mathcal{M}(s) \boxtimes \mathcal{M}(t))$ by proving each summand on the right belongs to $\mathrm{Der}(s \otimes t)$.

For $\mathrm{Der}(s) \boxtimes \mathcal{C}(t)$, take $(\alpha, \beta, \gamma) \in \mathrm{Der}(s)$, and $(f, g, h) \in \mathcal{C}(t)$. This means $s \bullet_2 \alpha + s \bullet_1 \beta = s \bullet_0 \gamma$ and $t \bullet_2 f = t \bullet_1 g = t \bullet_0 h$.

By definition, $(\alpha, \beta, \gamma) \boxtimes (f, g, h) = \big(\kappa_2(\alpha \otimes f), \kappa_1(\beta \otimes g), \kappa_0(\gamma \otimes h)\big)$, hence

$$\begin{aligned} (s \otimes t) \bullet_2 \kappa_2(\alpha \otimes f) + (s \otimes t) \bullet_1 \kappa_1(\beta \otimes g) &= \big((s \bullet_2 \alpha) \otimes (t \bullet_2 f)\big) + \big((s \bullet_1 \beta) \otimes (t \bullet_1 g)\big) \\ &= \big((s \bullet_2 \alpha) + (s \bullet_1 \beta)\big) \otimes t \bullet_0 h \\ &= (s \bullet_0 \gamma) \otimes (t \bullet_0 h) \\ &= (s \otimes t) \bullet_0 \kappa_0(\gamma \otimes h). \end{aligned}$$

Taking span, $\mathrm{Der}(s) \boxtimes \mathcal{C}(t)$ lies in $\mathrm{Der}(s \otimes t)$. Hence $\mathrm{Der}(s) \boxtimes \mathcal{C}(t) \subseteq \mathrm{Der}(s \otimes t)$ by linearity.

The argument for $\mathcal{C}(s) \boxtimes \mathrm{Der}(t) \subseteq \mathrm{Der}(s \otimes t)$ is identical, and the argument for nucleus terms is also analogous after setting one of the three terms to 0. $\square$

A full derivation decomposition example is the Appendix (Example D.2).

5.5. **Non-existence in higher valence.** In this subsection, we demonstrate an obstruction to being able to decompose the product of trilinear maps' derivations using the techniques from this section. Let $\mathcal{U} := \mathrm{Tri}(U_3, U_2, U_1; U_0)$, and $\mathcal{V} := \mathrm{Tri}(V_3, V_2, V_1; V_0)$, where Tri indicates trilinear. Fix $t_1 \in \mathcal{U}$ and $t_2 \in \mathcal{V}$, and let $(\delta^{(3)}, \delta^{(2)}, \delta^{(1)}, \delta^{(0)}) \in \mathrm{Der}(t_1 \otimes t_2)$, where $t_1 \otimes t_2$ is the product of trilinear maps defined analogously to Definition 2.8. This gives tensors $p, q, r, s \in \mathcal{U} \otimes \mathcal{V}$ with $p + q + r = s$. We begin with the four-term definition of the LIU decomposition.

Definition 5.16. Let U and V be finite dimensional $\mathbb{F}$-vector spaces. Let $p, q, r, s \in U \otimes V$ be nonzero elements satisfying $p + q + r = s$. Let

$$A := \mathrm{Im}_L(p), \quad B := \mathrm{Im}_L(q), \quad C := \mathrm{Im}_L(r), \quad D := \mathrm{Im}_L(s),$$
$$X := \mathrm{Im}_R(p), \quad Y := \mathrm{Im}_R(q), \quad Z := \mathrm{Im}_R(r), \quad W := \mathrm{Im}_R(s),$$

where $p \in A \otimes X$, $q \in B \otimes Y$, $r \in C \otimes Z$, and $s \in D \otimes W$ by Proposition 4.5.

A (p, q, r, s) *LIU decomposition* consists of a natural number n, and vectors

$$a_i \in A,\ b_i \in B,\ c_i \in C,\ d_i \in D, \quad x_i \in X,\ y_i \in Y,\ z_i \in Z,\ w_i \in W,$$

for $i \in [n]$, satisfying the following:

- **Decomposition equality.**

$$p = \sum_{i=1}^{n} a_i \otimes x_i, \quad q = \sum_{i=1}^{n} b_i \otimes y_i, \quad r = \sum_{i=1}^{n} c_i \otimes z_i, \quad s = \sum_{i=1}^{n} d_i \otimes w_i.$$

- **Local equality.**

$$\forall i \in [n], \quad a_i \otimes x_i + b_i \otimes y_i + c_i \otimes z_i = d_i \otimes w_i.$$

We shall show when $p + q + r = s$, there may not exist a LIU decomposition of the quadruple. As a LIU decomposition corresponds to decomposing a derivation of a product of trilinear maps by the algebraic invariants of the individual pieces, its non-existence rules out an analogous decomposition theorem using this strategy.

A natural question is whether one can reduce the $p + q + r = s$ case to the three-term case by setting $\tilde{p} = p + q$, and applying Lemma 5.13 to both $\tilde{p} + r = s$ and $p + q = \tilde{p}$. However, separate LIU decompositions of $(\tilde{p}, r, s)$ and $(p, q, \tilde{p})$ do not guarantee a LIU decomposition of (p, q, r, s). Applying Lemma 5.13 to $(\tilde{p}, r, s)$ gives

$$\tilde{p} = \sum_i \tilde{a}_i \otimes \tilde{x}_i, \quad \tilde{a}_i \otimes \tilde{x}_i \in \mathrm{Im}_L(\tilde{p}) \otimes \mathrm{Im}_R(\tilde{p}).$$

Choose the corresponding local terms for r and s, so for each i we have $\tilde{a}_i \otimes \tilde{x}_i + c_i \otimes z_i = d_i \otimes w_i$. The equation $p + q = \tilde{p}$ means $\mathrm{Im}_L(\tilde{p}) \leq \mathrm{Im}_L(p) + \mathrm{Im}_L(q)$ and $\mathrm{Im}_R(\tilde{p}) \leq \mathrm{Im}_R(p) + \mathrm{Im}_R(q)$.

To recover a (p, q, r, s) LIU decomposition, for each i we would have to refine this local equality, with $c_i \otimes z_i \in \mathrm{Im}_L(r) \otimes \mathrm{Im}_R(r)$ and $d_i \otimes w_i \in \mathrm{Im}_L(s) \otimes \mathrm{Im}_R(s)$, to a local equality $a_i \otimes x_i + b_i \otimes y_i + c_i \otimes z_i = d_i \otimes w_i$. Therefore the following equation must be satisfied for each i:

$$\tilde{a}_i \otimes \tilde{x}_i = a_i \otimes x_i + b_i \otimes y_i \quad (a_i \in \mathrm{Im}_L(p),\ x_i \in \mathrm{Im}_R(p),\ b_i \in \mathrm{Im}_L(q),\ y_i \in \mathrm{Im}_R(q)),$$

but this is not guaranteed. Rank-one terms in $(A + B) \otimes (X + Y)$ can contain cross blocks terms in $A \otimes Y$ and $B \otimes X$. For example, with $A = \langle e_1 \rangle$, $B = \langle e_2 \rangle$, $X = \langle f_1 \rangle$, $Y = \langle f_2 \rangle$,

$$(e_1 + e_2) \otimes (f_1 + f_2) = e_1 \otimes f_1 + e_1 \otimes f_2 + e_2 \otimes f_1 + e_2 \otimes f_2,$$

so this rank-one tensor has nonzero $A \otimes Y$ and $B \otimes X$ parts and cannot be of the form $a \otimes x + b \otimes y$ with $a \in A$, $x \in X$, $b \in B$, $y \in Y$.

Lemma 5.17. Assume $s \neq 0$. Let p, q, r, s be as the above definitions. If a LIU decomposition exists, then the intersection space

$$H := (A \otimes X + B \otimes Y + C \otimes Z) \cap (D \otimes W)$$

contains a nonzero rank-one tensor.

Proof. Each local equality has right-hand side $d_i \otimes w_i \in D \otimes W$ and left-hand side in $A \otimes X + B \otimes Y + C \otimes Z$, hence $d_i \otimes w_i \in H$ for every i. Since $s = \sum_{i=1}^n d_i \otimes w_i \neq 0$, at least one $d_i \otimes w_i$ is nonzero, giving a nonzero rank-one tensor in H. □

We now show that LIU decompositions may not exist.

Theorem 5.18. Let $\mathbb{F}$ be a field such that $\mathrm{char}(\mathbb{F}) \neq 2$, and let $J \in \mathbb{F}^{2\times 2}$ be a matrix with an irreducible quadratic minimal polynomial over $\mathbb{F}$. Let $U = V = \mathbb{F}^4$ be vector spaces. Then there exist nonzero elements $p, q, r, s \in U \otimes V$ satisfying $p + q + r = s$ such that the quadruple (p, q, r, s) admits no LIU decomposition.

Proof. Take $U = V = \mathbb{F}^4$ with standard bases $\{e_i\}$ and $\{f_i\}$, and subspaces $D = \mathrm{span}\{e_1, e_2\} \leq U$, $D' = \mathrm{span}\{e_3, e_4\} \leq U$, $W = \mathrm{span}\{f_1, f_2\} \leq V$, $W' = \mathrm{span}\{f_3, f_4\} \leq V$. Define isomorphisms

$$\iota_D : D \to D',\ e_1 \mapsto e_3,\ e_2 \mapsto e_4, \qquad \iota_W : W \to W',\ f_1 \mapsto f_3,\ f_2 \mapsto f_4.$$

For $J \in \mathbb{F}^{2\times 2}$, use the fixed bases (e_1, e_2) of D, (e_3, e_4) of D', (f_1, f_2) of W, and (f_3, f_4) of W' to define linear maps $J_D : D \to D'$ and $J_W : W \to W'$, where J_D acts on columns vectors of coordinates, and J_W on rows vectors of coordinates using the entries of J. Write $Jd := J_D(d)$ and $wJ := J_W(w)$. With this notation, define the subspaces

$$A = \{(d, \iota_D d) : d \in D\} \leq U, \qquad B = \{(d, -\iota_D d) : d \in D\} \leq U, \qquad C = \{(d, Jd) : d \in D\} \leq U,$$
$$X = \{(w, \iota_W w) : w \in W\} \leq V, \qquad Y = \{(w, -\iota_W w) : w \in W\} \leq V, \qquad Z = \{(w, wJ) : w \in W\} \leq V.$$

As a reminder, we define the intersection space $H := (A \otimes X + B \otimes Y + C \otimes Z) \cap (D \otimes W)$. We prove all H is nonempty and nonzero elements in H are rank 2.

Let $P, Q, R, S \in \mathbb{F}^{2\times 2}$. Relative to $U = D \oplus D'$ and $V = W \oplus W'$, for $p \in A \otimes X$, $q \in B \otimes Y$, $r \in C \otimes Z$, and $s \in D \otimes W$, we have

$$p = \begin{bmatrix} P & P \\ P & P \end{bmatrix}, \qquad q = \begin{bmatrix} Q & -Q \\ -Q & Q \end{bmatrix}, \qquad r = \begin{bmatrix} R & RJ \\ JR & JRJ \end{bmatrix}, \qquad s = \begin{bmatrix} S & 0 \\ 0 & 0 \end{bmatrix}. \tag{5.3}$$

Hence $p + q + r = s$ is equivalent to the matrix equation

$$\begin{bmatrix} P + Q + R & P - Q + RJ \\ P - Q + JR & P + Q + JRJ \end{bmatrix} = \begin{bmatrix} S & 0 \\ 0 & 0 \end{bmatrix},$$

meaning

$$P + Q + R = S, \qquad P - Q = -RJ, \qquad P - Q = -JR, \qquad P + Q = -JRJ.$$

Using the second and third equations, $RJ - JR = 0$, so R and J commute. Using the commuting fact on the last equation, we get $P + Q = -RJ^2$. Substitution $-RJ^2$ for $P + Q$ in the first equation, we conclude $S = R - RJ^2$. This defines P, Q, and R in terms of the fixed J and S as

$$P = -\frac{1}{2}RJ(I_2 + J), \qquad Q = \frac{1}{2}RJ(I_2 - J), \qquad R = S(I_2 - J^2)^{-1}. \tag{5.4}$$

Since $\mathrm{char}(\mathbb{F}) \neq 2$ and the minimal polynomial of J is an irreducible quadratic, it is separable, so every matrix in $\mathbb{F}^{2\times 2}$ commuting with J is a polynomial in J by [EG04, Thm. 3.1]. Hence $R \in \mathbb{F}[J]$ and R is invertible whenever nonzero.

As well, the matrices J, $I_2 + J$, $I_2 - J$, and $I_2 - J^2$ are invertible as the minimal polynomial of J is an irreducible quadratic. Combining the above two facts with Equation (5.4), we conclude S, P, and Q are invertible as well. Thus every nonzero element of H has rank 2.

For the existence portion, fix $S = I_2$. Let $a_i := (e_i, \iota_D e_i)$, $b_i := (e_i, -\iota_D e_i)$, $c_i := (e_i, Je_i)$, $d_i := (e_i, 0)$, and $x_i := (f_i, \iota_W f_i)$, $y_i := (f_i, -\iota_W f_i)$, $z_i := (f_i, f_i J)$, $w_i := (f_i, 0)$ for $i \in [2]$. Then, for instance, $p = a_1 \otimes (P_{11}x_1 + P_{12}x_2) + a_2 \otimes (P_{21}x_1 + P_{22}x_2)$, and similarly for q, r, s. Because P, Q, R, S are invertible, the two left factors and the two corresponding right factors are linearly independent in each case, so these are minimal rank decompositions. Hence Lemma 4.9 gives $\mathrm{Im}_L(p) = A, \mathrm{Im}_L(q) = B, \mathrm{Im}_L(r) = C, \mathrm{Im}_L(s) = D$ and $\mathrm{Im}_R(p) = X, \mathrm{Im}_R(q) = Y, \mathrm{Im}_R(r) = Z, \mathrm{Im}_R(s) = W$.

From (5.3), we recover a quadruple (p, q, r, s) whose intersection space has all nonzero elements of rank greater than 1. Applying Lemma 5.17, (p, q, r, s) admits no LIU decomposition.

□

Example 5.19. We apply Theorem 5.18 to get a specific example. Fix $\mathbb{F} = \mathbb{F}_5$, $J = \begin{bmatrix} 0 & 4 \\ 1 & 4 \end{bmatrix}$, and $S = I_2$. We compute P, Q, R from Equation (5.4):

$$P = \begin{bmatrix} 1 & 1 \\ 4 & 2 \end{bmatrix}, \qquad Q = \begin{bmatrix} 3 & 2 \\ 3 & 0 \end{bmatrix}, \qquad R = (I_2 - J^2)^{-1} = \begin{bmatrix} 2 & 2 \\ 3 & 4 \end{bmatrix}.$$

In the standard coordinates, we write down a quadruple (p, q, r, s) from Equation (5.3),

$$p = \left[\begin{array}{cc|cc} 1 & 1 & 1 & 1 \\ 4 & 2 & 4 & 2 \\ \hline 1 & 1 & 1 & 1 \\ 4 & 2 & 4 & 2 \end{array}\right], \quad q = \left[\begin{array}{cc|cc} 3 & 2 & 2 & 3 \\ 3 & 0 & 2 & 0 \\ \hline 2 & 3 & 3 & 2 \\ 2 & 0 & 3 & 0 \end{array}\right], \quad r = \left[\begin{array}{cc|cc} 2 & 2 & 2 & 1 \\ 3 & 4 & 4 & 3 \\ \hline 2 & 1 & 1 & 2 \\ 4 & 3 & 3 & 3 \end{array}\right], \quad s = \left[\begin{array}{cc|cc} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array}\right],$$

where for instance, $p = \sum_{i,j=1}^{2} P_{ij} a_i \otimes x_j$ where $a_i = (e_i, \iota_D e_i)$, $x_i = (f_i, \iota_W f_i)$ and so on. This example was checked by hand, as well as through a Magma computation, which is available at this gist.

6. Conclusion

This dissertation studies algebraic invariants of tensors from two perspectives. First, for tensors satisfying regularity conditions (Definitions 3.32 and 3.38), we develop solve and lift algorithms to find the algebraic invariants of tensors at lower cost than the status quo; see Theorems 3.21 and 3.33. The main idea is that the linear equations defining the algebraic invariants admit solving a smaller subsystem and then lifting those solutions to full solutions.

Second, we study algebraic invariants of products of bilinear maps. Specifically, Theorem 4.1 proves that the nuclei of a product of bilinear maps is a product of the nuclei of the factors. For the derivation algebra, Theorem 5.1 proves that the derivation algebra of the product of bilinear maps is determined by the algebraic invariants of the factors. This result extends classical results in algebra [Corollary 4.9 [BO81]] by viewing algebras through their multiplication tensors. We also prove an impossibility theorem on extending our approach to trilinear maps in Theorem 5.18.

6.1. Future Directions.

The results in the previous chapters motivate some future research directions.

Randomized Algorithms for fast Sylvester and Derivation equation solving.

The purpose of the regularity conditions of Definitions 3.32 and 3.38 is to be able to prove deterministic arithmetic complexity improvements using the solve and lift approach. An alternative is to analyze the performance of the algorithms Theorems 3.31 and 3.37 allowing for randomization. For instance, Freivalds' algorithm for matrix product verification can be used in place of the deterministic verification procedure. It tests whether $AB = C$ in Monte Carlo $O(n^2)$ cost [Fre79]. Analyzing the arithmetic complexity of randomized variants of our solve and lift algorithms, and of algorithms assuming an average case model of coefficients, is a related future direction.

Faster algorithms for higher valence tensors.

The solve and lift algorithms for Sylvester and derivation equations have been developed for 3-tensors. A natural question asks whether there exist analogous solve and lift algorithms for higher valence tensors. Notation and theory have been established for higher valence tensors in [BMW20].

Analyzing the equation $p + q = r$ by Kronecker modules

Kronecker modules classify the indecomposables of a pair of matrices M, N up to change of bases [Kro90]. The equation $p + q = r$ in the tensor product space $U \otimes V$ has data specified by the pair of matrices p and q. Thus the spaces given by the decomposition of Lemma 5.13 are constrained by the Kronecker module classification. More work is needed to understand what constraints are placed on the dimension of the spaces $U^{(i)}$ and $V^{(j)}$. Furthermore, the case for triples of matrices is

considered wild [Rin10], meaning it is as hard as classifying representations over arbitrary finite-dimensional algebras. Does the contrast between pairs of matrices having a classification and triples of matrices being wild relate to our finding that an analogous decomposition theorem does not exist for trilinear maps?

Decomposing the product of bilinear maps.

Our work in Sections 4 and 5 deals with decomposing the algebraic invariants of the product of bilinear maps. A next question is detecting whether a given bilinear map admits a decomposition $r \cong s \otimes t$. There are examples in algebra that suggest uniqueness and classification challenges, for instance the algebra $\mathbb{M}_2(\mathbb{C})$ decomposes in two different ways as tensor products over $\mathbb{R}$: both $\mathbb{H} \otimes \mathbb{C}$ and $\mathbb{M}_2(\mathbb{R}) \otimes \mathbb{C}$ are isomorphic to $\mathbb{M}_2(\mathbb{C})$.

Generalized products of multilinear maps.

In [FMW20], a (P, Ω)-tensor product is defined that generalizes the usual binary tensor product. Given a sequence of vector spaces $U_1, \ldots, U_n$ and relations specified by polynomial recurrences using the ideal $P \subset \mathbb{F}[x_1, \ldots x_n]$ and operators from the set $\Omega \subset \prod_{i=1}^{n} \mathrm{End}(U_i)$, we form the vector subspace $\Xi(P, \Omega) \leq U_1 \otimes \cdots U_n$ described as the (P, Ω)-tensor product space. This method has been used to compress the search space for tensor isomorphism problems [BMW22].

Throughout this dissertation, the tensor product vector space we use is for the special case $P = (x_2 - x_1)$, and $\Omega \cong \mathbb{F}$. Future work can dig into what kinds of tensors arise from (P, Ω)-tensor products, and whether there are decomposition theorems for algebraic invariants of those tensors.

On a related front, we only analyze the derivation of products of bilinear maps due to the obstruction found in the trilinear map case. Defining a generalized (P, Ω)-product of trilinear and higher valence tensors may allow their derivation and other algebraic invariants to be analyzed.

References

[Aza08] Saeid Azam. "Derivations of tensor product of algebras". In: *Communications in Algebra* 36 (2008), pp. 905–927. URL: https://doi.org/10.1080/00927870701776672.

[BKW24] Peter A. Brooksbank, Martin D. Kassabov, and James B. Wilson. *Detecting null patterns in tensor data*. 2024. arXiv: 2408.17425 [math.NA]. URL: https://arxiv.org/abs/2408.17425.

[BL08] Peter A. Brooksbank and Eugene M. Luks. "Testing isomorphism of modules". In: *Journal of Algebra* 320.11 (2008), pp. 4020–4029.

[BMW17] Peter A. Brooksbank, Joshua Maglione, and James B. Wilson. "A fast isomorphism test for groups whose Lie algebra has genus 2". In: *J. Algebra* 473 (2017), pp. 545–590. ISSN: 0021-8693. DOI: 10.1016/j.jalgebra.2016.12.007. URL: https://doi.org/10.1016/j.jalgebra.2016.12.007.

[BMW20] Peter A. Brooksbank, Joshua Maglione, and James B. Wilson. "Exact sequences of inner automorphisms of tensors". In: *Journal of Algebra* 545 (Mar. 2020), pp. 43–63. ISSN: 0021-8693. DOI: 10.1016/j.jalgebra.2019.07.006. URL: http://dx.doi.org/10.1016/j.jalgebra.2019.07.006.

[BMW22] Peter A. Brooksbank, Joshua Maglione, and James B. Wilson. "Tensor isomorphism by conjugacy of Lie algebras". In: *Journal of Algebra* 604 (Aug. 2022), pp. 790–807. ISSN: 0021-8693. DOI: 10.1016/j.jalgebra.2022.04.014. URL: http://dx.doi.org/10.1016/j.jalgebra.2022.04.014.

[BO81] G. M. Benkart and J. M. Osborn. "Derivations and Automorphisms of Nonassociative Matrix Algebras". In: *Transactions of the American Mathematical Society* 263.2 (1981), pp. 411–430. ISSN: 00029947. URL: http://www.jstor.org/stable/1998359 (visited on 12/06/2025).

[Bre17] Matej Brešar. "Derivations of tensor products of nonassociative algebras". In: *Linear Algebra and Applications* 530 (2017), pp. 244–252. URL: https://www.sciencedirect.com/science/article/pii/S0024379517303105.

[Bro97] Rasmus Bro. "PARAFAC. Tutorial and applications". In: *Chemometrics and Intelligent Laboratory Systems* 38.2 (1997), pp. 149–171.

[BS72] R. H. Bartels and G. W. Stewart. "Algorithm 432 [C2]: Solution of the matrix equation AX + XB = C [F4]". In: *Commun. ACM* 15.9 (Sept. 1972), pp. 820–826. ISSN: 0001-0782. DOI: 10.1145/361573.361582. URL: https://doi.org/10.1145/361573.361582.

[BW12] Peter A. Brooksbank and James B. Wilson. "Intersecting two classical groups". In: *J. Algebra* 353 (2012), pp. 286–297. ISSN: 0021-8693. DOI: 10.1016/j.jalgebra.2011.12.004. URL: https://doi.org/10.1016/j.jalgebra.2011.12.004.

[BW14] Peter A. Brooksbank and James B. Wilson. "Groups acting on tensor products". In: *Journal of Pure and Applied Algebra* 218.3 (2014), pp. 405–416. ISSN: 0022-4049. DOI: https://doi.org/10.1016/j.jpaa.2013.06.011. URL: https://www.sciencedirect.com/science/article/pii/S0022404913001382.

[DLDMV00] Lieven De Lathauwer, Bart De Moor, and Joos Vandewalle. "A Multilinear Singular Value Decomposition". In: *SIAM Journal on Matrix Analysis and Applications* 21.4 (2000), pp. 1253–1278.

[EG04] Wayne Eberly and Mark Giesbrecht. "Efficient decomposition of separable algebras". In: *Journal of Symbolic Computation* 37.1 (2004), pp. 35–81. DOI: 10.1016/S0747-7171(03)00082-6.

[FMW20] Uriya First, Joshua Maglione, and James B. Wilson. *A spectral theory for transverse tensor operators*. 2020. arXiv: 1911.02518 [math.SP]. URL: https://arxiv.org/abs/1911.02518.

[Fre79] Rūsiņš Freivalds. "Fast Probabilistic Algorithms". In: *Mathematical Foundations of Computer Science 1979*. Ed. by Jiří Bečvář. Vol. 74. Lecture Notes in Computer Science. Springer Berlin, Heidelberg, 1979, pp. 57–69. DOI: 10.1007/3-540-09526-8_5. URL: https://doi.org/10.1007/3-540-09526-8_5.

[GFA10] Morton E. Gurtin, Eliot Fried, and Lallit Anand. *The Mechanics and Thermodynamics of Continua*. Cambridge University Press, 2010.

[IQ19] Gábor Ivanyos and Youming Qiao. "Algorithms Based on *-Algebras, and Their Applications to Isomorphism of Polynomials with One Secret, Group Isomorphism, and Polynomial Identity Testing". In: *SIAM Journal on Computing* 48.3 (2019), pp. 926–963.

[Jac37] Nathan Jacobson. "Abstract derivation and Lie algebras". In: *Transactions of the American Mathematical Society* 42.2 (1937), pp. 206–224. DOI: 10.1090/s0002-9947-1937-1501922-7. URL: https://doi.org/10.1090/s0002-9947-1937-1501922-7.

[Kar+10] Alexandros Karatzoglou et al. "Multiverse Recommendation: N-dimensional Tensor Factorization for Context-Aware Collaborative Filtering". In: *Proceedings of the Fourth ACM Conference on Recommender Systems*. RecSys '10. 2010, pp. 79–86. DOI: 10.1145/1864708.1864727.

[KB09] Tamara G. Kolda and Brett W. Bader. "Tensor Decompositions and Applications". In: *SIAM Review* 51.3 (2009), pp. 455–500.

[KBV09] Yehuda Koren, Robert Bell, and Chris Volinsky. "Matrix Factorization Techniques for Recommender Systems". In: *Computer* 42.8 (2009), pp. 30–37. DOI: 10.1109/MC.2009.263.

[Kro90] Leopold Kronecker. "Algebraische Reduction der Schaaren bilinearer Formen". In: *Sitzungsberichte der Königlich Preussischen Akademie der Wissenschaften zu Berlin* (1890), pp. 1225–1237.

[Lan12] J.M. Landsberg. *Tensors: Geometry and Applications*. Graduate studies in mathematics. American Mathematical Society, 2012. ISBN: 9780821884812. URL: https://books.google.com/books?id=JTjv3DTvxZIC.

[LL00] George F. Leger and Eugene M. Luks. "Generalized Derivations of Lie Algebras". In: *Journal of Algebra* 228.1 (2000), pp. 165–203. ISSN: 0021-8693. DOI: https://doi.org/10.1006/jabr.1999.8250. URL: https://www.sciencedirect.com/science/article/pii/S0021869399982509.

[MM10] Takanori Maehara and Kazuo Murota. "A numerical algorithm for block-diagonal decomposition of matrix *-algebras with general irreducible components". In: *Japan Journal of Industrial and Applied Mathematics* 27 (2010), pp. 263–293. URL: https://api.semanticscholar.org/CorpusID:7116115.

[Mya90] A. G. Myasnikov. "Definable invariants of bilinear mappings". In: *Siberian Mathematical Journal* 31.1 (1990), pp. 89–99.

[NH08] M. Novotný and J. Hřivnák. "On (α, β, γ)-derivations of Lie algebras and corresponding invariant functions". In: *Journal of Geometry and Physics* 58.2 (2008), pp. 208–217. DOI: 10.1016/j.geomphys.2007.09.005. URL: https://www.sciencedirect.com/science/article/pii/S0393044007001337.

[Rin10] Claus Michael Ringel. *Indecomposable Representations of the Kronecker Quivers*. 2010. DOI: 10.48550/arXiv.1009.5635. arXiv: 1009.5635 [math.RT]. URL: https://arxiv.org/abs/1009.5635.

[RS18] Elina Robeva and Anna Seigal. "Duality of graphical models and tensor networks". In: *Information and Inference: A Journal of the IMA* 8.2 (June 2018), pp. 273–288. ISSN: 2049-8772. DOI: 10.1093/imaiai/iay009. eprint: https://academic.oup.com/imaiai/article-pdf/8/2/273/28864933/iay009.pdf. URL: https://doi.org/10.1093/imaiai/iay009.

[Sch95] Richard D. Schafer. *An Introduction to Nonassociative Algebras*. New York: Dover Publications, 1995.

[Syl84] J. J. Sylvester. "Sur l'équation en matrices $px = xq$". In: *Comptes Rendus de l'Académie des Sciences de Paris* 99 (1884), pp. 67–71, 115–116.

[Tuc66] L. R. Tucker. "Some mathematical notes on three-mode factor analysis". In: *Psychometrika* 31 (1966c), pp. 279–311.

[Wil09] James B. Wilson. "Decomposing p-groups via Jordan algebras". In: *Journal of Algebra* 322.8 (2009), pp. 2642–2679.

[Wil12] James B. Wilson. "Existence, algorithms, and asymptotics of direct product decompositions, I". In: *Groups - Complexity - Cryptology* 4.1 (Jan. 2012). ISSN: 1869-6104. DOI: 10.1515/gcc-2012-0007. URL: http://dx.doi.org/10.1515/gcc-2012-0007.

[Wil15] James B. Wilson. *Skolem-Noether for nilpotent products*. 2015. arXiv: 1507.04406 [math.RT]. URL: https://arxiv.org/abs/1507.04406.

[Wil16] James B. Wilson. "On automorphisms of groups, rings, and algebras". In: *Communications in Algebra* 45.4 (Oct. 2016), pp. 1452–1478. ISSN: 1532-4125. DOI: 10.1080/00927872.2016.1175617. URL: http://dx.doi.org/10.1080/00927872.2016.1175617.

Appendix A. Implementation performance comparisons

We show some implementation results of Theorems 3.21 and 3.33. The code to reproduce this is available at https://github.com/cliu587/fast-der-solver. The timings are done on an Apple M1 CPU.

A.1. Simultaneous Sylvester Implementation Results. ↑

We create instances where the regularity assumptions of Definition 3.32 holds. The performance using the solution strategy described in Theorem 3.21 is compared to solving the large linear system as laid out in Example 3.20. See Figure 2 for the performance numbers.

A.2. Derivation Solver Implementation Results. ↑

We create instances where the regularity assumptions of Definition 3.38 holds. The performance using the solution strategy described in Theorem 3.33 is compared to solving the large linear system as laid out in an analogous fashion to Example 3.20. See Figure 3 for the performance numbers.

Appendix B. Examples for Simultaneous Sylvester and Derivation Equations

B.1. Simultaneous Sylvester Equation Example. ↑



FIGURE 2. Implementation performance comparison of the 2 algorithms. Both axes are on a logarithmic scale. We see the performance improvements of the fast solve and lift algorithm as compared to solving the full linear system. Each algorithm's timing was done as the median of 5 trials.

Example B.1. Let $\mathbb{F} = \mathbb{Q}$, with $a = b = r = s = 2$ and $c = 4$. Define

$$R[:,:,1] = \begin{bmatrix} 1 & 2 \\ 1 & 4 \end{bmatrix}, \quad R[:,:,2] = \begin{bmatrix} 3 & 1 \\ 4 & 3 \end{bmatrix}, \quad R[:,:,3] = \begin{bmatrix} 3 & 4 \\ 2 & 1 \end{bmatrix}, \quad R[:,:,4] = \begin{bmatrix} 0 & 2 \\ 1 & 0 \end{bmatrix},$$
$$S[:,:,1] = \begin{bmatrix} 2 & 1 \\ 4 & 1 \end{bmatrix}, \quad S[:,:,2] = \begin{bmatrix} 3 & 2 \\ 5 & 2 \end{bmatrix}, \quad S[:,:,3] = \begin{bmatrix} 1 & 0 \\ 2 & 0 \end{bmatrix}, \quad S[:,:,4] = \begin{bmatrix} 4 & 1 \\ 2 & 3 \end{bmatrix},$$
$$T[:,:,1] = \begin{bmatrix} 11 & 7 \\ 24 & 25 \end{bmatrix}, \quad T[:,:,2] = \begin{bmatrix} 19 & 10 \\ 44 & 23 \end{bmatrix}, \quad T[:,:,3] = \begin{bmatrix} 3 & -1 \\ 20 & 15 \end{bmatrix}, \quad T[:,:,4] = \begin{bmatrix} 20 & 7 \\ 20 & 11 \end{bmatrix}.$$

The full system has the unique solution

$$X = \begin{bmatrix} -1 & 1 \\ 2 & 3 \end{bmatrix}, \qquad Y = \begin{bmatrix} 4 & 2 \\ 3 & 1 \end{bmatrix}.$$

Now run the solve-and-lift algorithm with $I = J = \{1\}$.

Double restricted. Set $X_I = \begin{bmatrix} x_1 & x_2 \end{bmatrix}$, and $Y_J = \begin{bmatrix} y_1 & y_2 \end{bmatrix}^\mathsf{T}$. The restricted system is

$$\begin{bmatrix} 1 & 1 & 2 & 1 \\ 3 & 4 & 3 & 2 \\ 3 & 2 & 1 & 0 \\ 0 & 1 & 4 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ y_1 \\ y_2 \end{bmatrix} = \begin{bmatrix} 11 \\ 19 \\ 3 \\ 20 \end{bmatrix}.$$

It has the unique solution $X_I = \begin{bmatrix} -1 & 1 \end{bmatrix}$, and $Y_J = \begin{bmatrix} 4 & 3 \end{bmatrix}^\mathsf{T}$. Hence $\mathcal{S}_{\mathrm{DR}} = \{(X_I, Y_J)\}$.



FIGURE 3. Implementation performance of the derivation solve-and-lift strategy compared to solving the full linear system. Both axes are on a logarithmic scale. We see clearly the performance improvements of the fast solve and lift algorithm as compared to solving the full linear system. Each algorithm's timing was done as the median of 5 trials.

Row restricted. With $\widehat{J} = \{2\}$,

$$M_R = \begin{bmatrix} 2 & 1 \\ 3 & 2 \\ 1 & 0 \\ 4 & 1 \end{bmatrix}, \qquad N_R(X_I, Y_J) = \begin{bmatrix} 5 \\ 8 \\ 2 \\ 9 \end{bmatrix}.$$

Solving $M_R \widehat{Y} = N_R(X_I, Y_J)$ gives $\widehat{Y} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$, and $Y = \begin{bmatrix} 4 & 2 \\ 3 & 1 \end{bmatrix}$. So $\mathcal{S}_{\mathrm{R}} = \left\{ \left(\begin{bmatrix} -1 & 1 \end{bmatrix}, \begin{bmatrix} 4 & 2 \\ 3 & 1 \end{bmatrix} \right) \right\}$.

Column restricted. With $\widehat{I} = \{2\}$,

$$M_C = \begin{bmatrix} 1 & 1 \\ 3 & 4 \\ 3 & 2 \\ 0 & 1 \end{bmatrix}, \qquad N_C(X_I, Y_J) = \begin{bmatrix} 5 \\ 18 \\ 12 \\ 3 \end{bmatrix}.$$

Solving $M_C \widehat{X} = N_C(X_I, Y_J)$ gives $\widehat{X} = \begin{bmatrix} 2 \\ 3 \end{bmatrix}$, and $X = \begin{bmatrix} -1 & 1 \\ 2 & 3 \end{bmatrix}$. So $\mathcal{S}_{\mathrm{C}} = \left\{ \left(\begin{bmatrix} -1 & 1 \\ 2 & 3 \end{bmatrix}, \begin{bmatrix} 4 \\ 3 \end{bmatrix} \right) \right\}$.

Lift to full. The pullback is $P = \{(z_R, z_C) \in \mathcal{S}_{\mathrm{R}} \times \mathcal{S}_{\mathrm{C}} : \pi_J(z_R) = \pi_I(z_C)\} = \{p_0\}$, where

$$p_0 := \big((X_I, Y), (X, Y_J)\big), \qquad X_I = X[I, :], \;\; Y_J = Y[:, J].$$

Then $\mathrm{Err}(\Psi(p_0)) = \big(XR[:,:,k] + S[:,:,k]Y - T[:,:,k]\big)_{k \in [4]} = 0_{2\times 2\times 4}$. Therefore LiftRestricted returns the unique solution (X, Y) above.

B.2. **Derivation Equation.** ↑

Example B.2. We illustrate the solution strategy on a $2 \times 2 \times 2$ instance. Let $a = b = c = r = s = t = 2$, and $I = J = K = \{1\}$. Define the slices

$$R[:,:,1] = S[:,:,1] = T[:,:,1] = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \qquad R[:,:,2] = S[:,:,2] = T[:,:,2] = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}.$$

Write the unknowns as

$$X = \begin{bmatrix} x_{11} & x_{12} \\ x_{21} & x_{22} \end{bmatrix}, \qquad Y = \begin{bmatrix} y_{11} & y_{12} \\ y_{21} & y_{22} \end{bmatrix}, \qquad Z = \begin{bmatrix} z_{11} & z_{12} \\ z_{21} & z_{22} \end{bmatrix}.$$

We solve the derivation equations by brute force first. They are for $k \in \{1, 2\}$, $X\, R[:,:,k] + S[:,:,k]\, Y = (T^Z)[:,:,k]$.

For $k = 1$, this is $X \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} + \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} Y = z_{11} T[:,:,1] + z_{21} T[:,:,2]$. This gives the relations

$$\begin{bmatrix} x_{11} + y_{11} & y_{12} \\ x_{21} & 0 \end{bmatrix} = \begin{bmatrix} z_{11} & 0 \\ 0 & z_{21} \end{bmatrix},$$

so $y_{12} = x_{21} = z_{21} = 0$, and $x_{11} + y_{11} = z_{11}$. And for $k = 2$,

$$X \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} + \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} Y = z_{12} T[:,:,1] + z_{22} T[:,:,2],$$

gives the relations

$$\begin{bmatrix} 0 & x_{12} \\ y_{21} & x_{22} + y_{22} \end{bmatrix} = \begin{bmatrix} z_{12} & 0 \\ 0 & z_{22} \end{bmatrix},$$

so $z_{12} = x_{12} = y_{21} = 0$, and $x_{22} + y_{22} = z_{22}$. Combined, the solution space is

$$X = \begin{bmatrix} \alpha & 0 \\ 0 & \beta \end{bmatrix}, \qquad Y = \begin{bmatrix} \gamma & 0 \\ 0 & \delta \end{bmatrix}, \qquad Z = \begin{bmatrix} \alpha + \gamma & 0 \\ 0 & \beta + \delta \end{bmatrix}, \qquad \alpha, \beta, \gamma, \delta \in \mathbb{F},$$

We now construct the solution space using the steps described in this section.

Step 1: Triple-restricted solve. With $I = J = K = \{1\}$, the unknown blocks are $X_I = \begin{bmatrix} x_{11} & x_{12} \end{bmatrix}$, $Y_J = \begin{bmatrix} y_{11} & y_{21} \end{bmatrix}^\mathsf{T}$, and $Z_K = \begin{bmatrix} z_{11} & z_{21} \end{bmatrix}^\mathsf{T}$, and the triple-restricted equation $X_I R[:,J,1] + S[I,:,1] Y_J = (T^{Z_K})[I,J,1]$ is just the $(1,1)$-entry of the $k = 1$ equation computed above. Since

$$R[:,J,1] = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \qquad S[I,:,1] = \begin{bmatrix} 1 & 0 \end{bmatrix}, \qquad (T^{Z_K})[I,J,1] = \big(z_{11} T[:,:,1] + z_{21} T[:,:,2]\big)[1,1] = z_{11},$$

it reduces to $x_{11} + y_{11} = z_{11}$.

Step 2: Column-depth lift. Recall $\widehat{I} = [a] \setminus I = \{2\}$ and $\widehat{X} = X[\widehat{I},:]^\mathsf{T} = \begin{bmatrix} x_{21} & x_{22} \end{bmatrix}^\mathsf{T}$. Using the notation from above:

$$M_C = \big[(R[:,J,1])^\mathsf{T}\big] = \begin{bmatrix} 1 & 0 \end{bmatrix}, \qquad N_C(X_I, Y_J, Z_K) = \big[(T^{Z_K})[\widehat{I},J,1] - S[\widehat{I},:,1] Y_J\big]^\mathsf{T} = \begin{bmatrix} 0 \end{bmatrix}.$$

So the equation $M_C \widehat{X} = N_C(X_I, Y_J, Z_K)$ corresponds to the system $\begin{bmatrix} 1 & 0 \end{bmatrix} \begin{bmatrix} x_{21} & x_{22} \end{bmatrix}^\mathsf{T} = \begin{bmatrix} 0 \end{bmatrix}$, concluding $x_{21} = 0$.

Step 3: Row-depth lift. Recall $\widehat{J} = [b] \setminus J = \{2\}$ and $\widehat{Y} = Y[:,\widehat{J}] = \begin{bmatrix} y_{12} & y_{22} \end{bmatrix}^\mathsf{T}$. The matrices to solve in the lift are

$$M_R = \big[S[I,:,1]\big] = \begin{bmatrix} 1 & 0 \end{bmatrix}, \qquad N_R(X_I, Y_J, Z_K) = \big[(T^{Z_K})[I,\widehat{J},1] - X_I R[:,\widehat{J},1]\big] = \begin{bmatrix} 0 \end{bmatrix}.$$

So the equation $M_R \widehat{Y} = N_R(X_I, Y_J, Z_K)$ corresponds to the system $\begin{bmatrix} 1 & 0 \end{bmatrix} \begin{bmatrix} y_{12} & y_{22} \end{bmatrix}^\mathsf{T} = \begin{bmatrix} 0 \end{bmatrix}$, concluding $y_{12} = 0$.

Step 4: Row-col lift. Recall $\widehat{K} = [c] \setminus K = \{2\}$ and $\widehat{Z} = Z[:,\widehat{K}] = \begin{bmatrix} z_{12} & z_{22} \end{bmatrix}^\mathsf{T}$. The row-col lift uses

$$M_D = \big[\mathrm{vec}(T[I,J,1]) \quad \mathrm{vec}(T[I,J,2])\big] = \begin{bmatrix} 1 & 0 \end{bmatrix}, \qquad N_D(X_I, Y_J, Z_K) = \mathrm{vec}\big(X_I R[:,J,2] + S[I,:,2] Y_J\big) = \begin{bmatrix} 0 \end{bmatrix}.$$

So the equation $M_D \widehat{Z} = N_D(X_I, Y_J, Z_K)$ corresponds to the system $\begin{bmatrix} 1 & 0 \end{bmatrix} \begin{bmatrix} z_{12} & z_{22} \end{bmatrix}^\mathsf{T} = \begin{bmatrix} 0 \end{bmatrix}$, concluding $z_{12} = 0$.

Step 5: Pullback and final residual solve. Using the three lifts and the pullback conditions, the compatible candidates are

$$X = \begin{bmatrix} x_{11} & x_{12} \\ 0 & x_{22} \end{bmatrix}, \qquad Y = \begin{bmatrix} y_{11} & 0 \\ y_{21} & y_{22} \end{bmatrix}, \qquad Z = \begin{bmatrix} x_{11} + y_{11} & 0 \\ z_{21} & z_{22} \end{bmatrix}.$$

Here Step 1 gives $z_{11} = x_{11} + y_{11}$, Step 2 gives $x_{21} = 0$, Step 3 gives $y_{12} = 0$, and Step 4 gives $z_{12} = 0$. Now apply the final lift condition $\mathrm{Err}(X, Y, Z) = 0$, where $\mathrm{Err}(X, Y, Z)[:, :, k] = X\, R[:, :, k] + S[:, :, k]Y - (T^Z)[:, :, k]$. For the candidate solution above,

$$(T^Z)[:, :, 1] = \begin{bmatrix} x_{11} + y_{11} & 0 \\ 0 & z_{21} \end{bmatrix}, \qquad \mathrm{Err}(X, Y, Z)[:, :, 1] = \begin{bmatrix} 0 & 0 \\ 0 & -z_{21} \end{bmatrix},$$
$$(T^Z)[:, :, 2] = \begin{bmatrix} 0 & 0 \\ 0 & z_{22} \end{bmatrix}, \qquad \mathrm{Err}(X, Y, Z)[:, :, 2] = \begin{bmatrix} 0 & x_{12} \\ y_{21} & x_{22} + y_{22} - z_{22} \end{bmatrix}.$$

So $\mathrm{Err}(X, Y, Z) = 0$ requires $z_{21} = 0$, $x_{12} = 0$, $y_{21} = 0$, and $z_{22} = x_{22} + y_{22}$. Hence the full derivation space matches what we computed initially as

$$X = \begin{bmatrix} \alpha & 0 \\ 0 & \beta \end{bmatrix}, \qquad Y = \begin{bmatrix} \gamma & 0 \\ 0 & \delta \end{bmatrix}, \qquad Z = \begin{bmatrix} \alpha + \gamma & 0 \\ 0 & \beta + \delta \end{bmatrix}, \qquad \alpha, \beta, \gamma, \delta \in \mathbb{F},$$

Appendix C. Proof of Proposition 5.3

As a reminder, in Proposition 5.3, R and S are finite-dimensional unital non-associative algebras with multiplication maps μ_R and μ_S. Throughout this section, we will use the following notation.

$$X \coloneqq \mathrm{Der}(\mu_R) \boxtimes \mathcal{C}(\mu_S), \qquad Y \coloneqq \mathcal{C}(\mu_R) \boxtimes \mathrm{Der}(\mu_S),$$
$$N \coloneqq \iota_{20}(\mathcal{L}(\mu_R) \boxtimes \mathcal{L}(\mu_S)) + \iota_{10}(\mathcal{R}(\mu_R) \boxtimes \mathcal{R}(\mu_S)) + \iota_{21}(\mathcal{M}(\mu_R) \boxtimes \mathcal{M}(\mu_S)).$$

The argument starts from the decomposition $\mathrm{Der}(\mu_R \otimes \mu_S) = X + Y + N$ given by Theorem 5.1. Since algebra derivations are diagonal triples, it is enough to analyze the diagonal of $X + Y + N$. We first show that the diagonal of $X + Y + N$ is the sum of the diagonal subspaces of the three summands, and then identify each piece with the corresponding terms in Brešar's decomposition.

Lemma C.1. If A is a unital algebra and $\mu : A \times A \rightarrowtail A$ is its multiplication tensor, then μ is fully nondegenerate.

Proof. $\mu(a, 1) = a$ and $\mu(1, a) = a$ force both the left and right radical to be 0, and $\mathrm{span}(\mu(A, A)) = A$ since every $a \in A$ equals $\mu(a, 1)$. □

Applying Lemma C.1 to R and S, the multiplication tensors μ_R and μ_S are fully nondegenerate. The map $d \mapsto (d, d, d)$ identifies the derivation algebra of A, $\mathcal{D}_{\mathrm{alg}}(A)$, with the diagonal of $\mathrm{Der}(\mu_A)$. For $x \in A$, write $\lambda_x(a) \coloneqq xa$ and $\rho_x(a) \coloneqq ax$ for the left and right multiplication operators. Recall the definitions for a unital algebra A [Sch95, Ch. I, §2, p. 10]:

$$\begin{aligned}
&\text{(left nucleus)} && \mathrm{Nuc}_\ell(A) \coloneqq \{x \in A : \forall a, b \in A,\ (xa)b = x(ab)\}, \\
&\text{(middle nucleus)} && \mathrm{Nuc}_m(A) \coloneqq \{y \in A : \forall a, b \in A,\ (ay)b = a(yb)\}, \\
&\text{(right nucleus)} && \mathrm{Nuc}_r(A) \coloneqq \{z \in A : \forall a, b \in A,\ a(bz) = (ab)z\}, \\
&\text{(nucleus)} && \mathrm{Nuc}(A) \coloneqq \mathrm{Nuc}_\ell(A) \cap \mathrm{Nuc}_m(A) \cap \mathrm{Nuc}_r(A), \\
&\text{(center)} && Z(A) \coloneqq \{c \in \mathrm{Nuc}(A) : \forall a \in A,\ ca = ac\}.
\end{aligned}$$

We do not separately define the centroid of an algebra A, since in the unital case it identifies canonically with the center. The tensor versions are as follows.

Proposition C.2. Let A be a unital algebra with multiplication tensor μ. Then

$$\mathcal{M}(\mu) = \{(\rho_y, \lambda_y) : y \in \mathrm{Nuc}_m(A)\}, \qquad \mathcal{L}(\mu) = \{(\lambda_x, \lambda_x) : x \in \mathrm{Nuc}_\ell(A)\},$$
$$\mathcal{R}(\mu) = \{(\rho_z, \rho_z) : z \in \mathrm{Nuc}_r(A)\}, \qquad \mathcal{C}(\mu) = \{(\lambda_c, \lambda_c, \lambda_c) : c \in Z(A)\}.$$

Proof. We prove $\mathcal{M}(\mu) = \{(\rho_y, \lambda_y) : y \in \mathrm{Nuc}_m(A)\}$.

For $\subseteq$, let $(\alpha, \beta) \in \mathcal{M}(\mu)$. Then for all a, b in A, $(\alpha(a))b = a(\beta(b))$. Setting $b = 1$ gives $\alpha(a) = a\,\beta(1)$, so $\alpha = \rho_y$ with $y = \beta(1)$. Setting $a = 1$ gives $\beta(b) = \alpha(1)\,b$, so $\beta = \lambda_y$ with $y = \alpha(1)$. Hence $y = \alpha(1) = \beta(1)$ and $(ay)b = a(yb)$, so $y \in \mathrm{Nuc}_m(A)$. Therefore $(\alpha, \beta) = (\rho_y, \lambda_y)$ is in the right-hand side.

For $\supseteq$, let $y \in \mathrm{Nuc}_m(A)$. Then for all $a, b \in A$, $(\rho_y(a))b = (ay)b = a(yb) = a(\lambda_y(b))$, so $(\rho_y, \lambda_y) \in \mathcal{M}(\mu)$.

The equations for $\mathcal{L}(\mu)$ and $\mathcal{R}(\mu)$ are proved identically. Finally,

$$\begin{aligned}\mathcal{C}(\mu) &= \{(\alpha, \beta, \gamma) : \mu \bullet_2 \alpha = \mu \bullet_1 \beta = \mu \bullet_0 \gamma\} \\ &= \{(\alpha, \beta, \gamma) : (\alpha, \beta) \in \mathcal{M}(\mu),\ (\alpha, \gamma) \in \mathcal{L}(\mu),\ (\beta, \gamma) \in \mathcal{R}(\mu)\}.\end{aligned}$$

Using the previous three parts, let $(\alpha, \beta, \gamma) \in \mathcal{C}(\mu)$. Then $(\alpha, \gamma) = (\lambda_x, \lambda_x)$, $(\beta, \gamma) = (\rho_z, \rho_z)$, and $(\alpha, \beta) = (\rho_y, \lambda_y)$ for some $x \in \mathrm{Nuc}_\ell(A)$, $z \in \mathrm{Nuc}_r(A)$, $y \in \mathrm{Nuc}_m(A)$. Hence $\lambda_x = \rho_z$, $\lambda_x = \rho_y$, and $\rho_z = \lambda_y$, so $x = y = z =: c$ and $\lambda_c = \rho_c$. Thus $c \in \mathrm{Nuc}_\ell(A) \cap \mathrm{Nuc}_m(A) \cap \mathrm{Nuc}_r(A)$ and $ca = ac$ for all a. This means $c \in Z(A)$.

Conversely, if $c \in Z(A)$ then $(\lambda_c, \lambda_c, \lambda_c) \in \mathcal{C}(\mu)$. □

Definition C.3. Let V be a vector space. For a subspace $W \leq \mathrm{End}(V)^3$, define $W^{\mathrm{diag}} := \{d \in \mathrm{End}(V) : (d, d, d) \in W\}$.

For $u \in \mathrm{Nuc}(A)$, write $\mathrm{ad}(u) := \lambda_u - \rho_u$; then $\mathrm{ad}(u) \in \mathcal{D}_{\mathrm{alg}}(A)$. The following proposition is key to identifying the $\mathrm{ad}(u)$ piece of Brešar's decomposition.

Proposition C.4. Let A be a unital algebra with multiplication tensor μ, and let $N_A := \iota_{20}(\mathcal{L}(\mu)) + \iota_{10}(\mathcal{R}(\mu)) + \iota_{21}(\mathcal{M}(\mu))$. Then $N_A^{\mathrm{diag}} = \mathrm{ad}(\mathrm{Nuc}(A))$.

Proof. For $\supseteq$, let $u \in \mathrm{Nuc}(A)$. By Proposition C.2, $(\lambda_u, \lambda_u) \in \mathcal{L}(\mu)$, $(\rho_u, \rho_u) \in \mathcal{R}(\mu)$, and $(\rho_u, \lambda_u) \in \mathcal{M}(\mu)$. Hence $\iota_{20}(\lambda_u, \lambda_u) - \iota_{10}(\rho_u, \rho_u) + \iota_{21}(\rho_u, \lambda_u) = (\mathrm{ad}(u), \mathrm{ad}(u), \mathrm{ad}(u)) \in N_A$, so $\mathrm{ad}(u) \in N_A^{\mathrm{diag}}$.

For $\subseteq$, let $d \in N_A^{\mathrm{diag}}$. Then $(d, d, d) \in N_A$, so $(d, d, d) = \iota_{20}(\alpha, \gamma) + \iota_{10}(\beta, \gamma') + \iota_{21}(\alpha', \beta')$. By Proposition C.2, there exist $a \in \mathrm{Nuc}_\ell(A)$, $b \in \mathrm{Nuc}_r(A)$, and $c \in \mathrm{Nuc}_m(A)$ with

$$(\alpha, \gamma) = (\lambda_a, \lambda_a), \quad (\beta, \gamma') = (\rho_b, \rho_b), \quad (\alpha', \beta') = (\rho_c, \lambda_c).$$

Therefore $(d, d, d) = (\lambda_a - \rho_c,\ \rho_b + \lambda_c,\ \lambda_a + \rho_b)$. Comparing components gives $\rho_b = -\rho_c$, and $\lambda_a = \lambda_c$. Since A is unital, left and right multiplication are injective, so $a = c$, and $b = -c$. Hence $c \in \mathrm{Nuc}_\ell(A) \cap \mathrm{Nuc}_m(A) \cap \mathrm{Nuc}_r(A) = \mathrm{Nuc}(A)$ and $(d, d, d) = (\mathrm{ad}(c), \mathrm{ad}(c), \mathrm{ad}(c))$, so $d \in \mathrm{ad}(\mathrm{Nuc}(A))$. □

For the rest of this section, also let

$$N_R := \iota_{20}(\mathcal{L}(\mu_R)) + \iota_{10}(\mathcal{R}(\mu_R)) + \iota_{21}(\mathcal{M}(\mu_R)), \qquad N_S := \iota_{20}(\mathcal{L}(\mu_S)) + \iota_{10}(\mathcal{R}(\mu_S)) + \iota_{21}(\mathcal{M}(\mu_S)).$$

For a unital algebra A, define the *defect map*

$$\epsilon_A : \mathrm{End}(A)^3 \to A^2, \qquad \epsilon_A(\alpha, \beta, \gamma) := (\alpha(1_A), \beta(1_A)).$$

This definition extends to subspaces $U \leq \mathrm{End}(A)^3$ as $\epsilon_A(U) := \{\epsilon_A(u) : u \in U\}$. The following shows the triples in $\mathrm{Der}(\mu_A)$ with zero defect come from algebra derivations.

Lemma C.5. $(\mathrm{Der}(\mu_A) \cap \mathrm{Ker}(\epsilon_A))^{\mathrm{diag}} = \mathcal{D}_{\mathrm{alg}}(A)$.

Proof. Let $d \in (\mathrm{Der}(\mu_A) \cap \mathrm{Ker}(\epsilon_A))^{\mathrm{diag}}$. Then $(d, d, d) \in \mathrm{Der}(\mu_A)$ and $\epsilon_A(d, d, d) = 0$, so $d(1_A) = 0$. Since $(d, d, d) \in \mathrm{Der}(\mu_A)$, we have $d(x)y + xd(y) = d(xy)$ for all $x, y \in A$, hence $d \in \mathcal{D}_{\mathrm{alg}}(A)$.

Conversely, if $d \in \mathcal{D}_{\mathrm{alg}}(A)$, then $(d, d, d) \in \mathrm{Der}(\mu_A)$. Setting $y = 1_A$ in $d(xy) = d(x)y + xd(y)$ gives $d(x) = d(x) + xd(1_A)$ for all $x \in A$, so $xd(1_A) = 0$ for all x. Taking $x = 1_A$ gives $d(1_A) = 0$. Thus $(d, d, d) \in \mathrm{Ker}(\epsilon_A)$, so $d \in (\mathrm{Der}(\mu_A) \cap \mathrm{Ker}(\epsilon_A))^{\mathrm{diag}}$. □

Remark C.6. For any subspace $W \leq \mathrm{Der}(\mu_A)$, $W \cap \mathrm{Ker}(\epsilon_A) = \{(d, d, d) : d \in W^{\mathrm{diag}}\}$. Consequently, $(W \cap \mathrm{Ker}(\epsilon_A))^{\mathrm{diag}} = W^{\mathrm{diag}}$.

For $A = R \otimes S$, if $(\alpha, \beta, \gamma) \in \mathrm{End}(R)^3$ and $(f, g, h) \in \mathrm{End}(S)^3$, then evaluating ϵ_A at $(1_R \otimes 1_S)$ gives

$$\epsilon_A\big((\alpha, \beta, \gamma) \boxtimes (f, g, h)\big) = (\alpha(1_R) \otimes f(1_S),\ \beta(1_R) \otimes g(1_S)). \tag{C.1}$$

We shall also need the following for the proof.

Remark C.7. If $z \in Z(S)$, then $u \boxtimes (\lambda_z, \lambda_z, \lambda_z) \in N$ for every $u \in N_R$. Analogously, if $c \in Z(R)$, then $(\lambda_c, \lambda_c, \lambda_c) \boxtimes v \in N$ for every $v \in N_S$.

Proof. By Proposition C.2, $z \in Z(S)$ gives $(\lambda_z, \lambda_z) \in \mathcal{L}(\mu_S)$, $(\rho_z, \rho_z) \in \mathcal{R}(\mu_S)$, and $(\rho_z, \lambda_z) \in \mathcal{M}(\mu_S)$. Writing $u = u_{20} + u_{10} + u_{21}$ with $u_{20} \in \iota_{20}(\mathcal{L}(\mu_R))$, $u_{10} \in \iota_{10}(\mathcal{R}(\mu_R))$, and $u_{21} \in \iota_{21}(\mathcal{M}(\mu_R))$, each $u_{ij} \boxtimes (\lambda_z, \lambda_z, \lambda_z)$ lies in the matching summand of N, hence so does the sum. The second statement is analogous. □

The below key lemma decomposes the diagonal of $X + Y + N$ as the sum of the diagonal subspaces of the summands.

Lemma C.8. With the notation above,

$$(X + Y + N)^{\mathrm{diag}} = X^{\mathrm{diag}} + Y^{\mathrm{diag}} + N^{\mathrm{diag}}.$$

Proof. The inclusion $X^{\mathrm{diag}} + Y^{\mathrm{diag}} + N^{\mathrm{diag}} \subseteq (X + Y + N)^{\mathrm{diag}}$ is immediate.

We now prove the converse. Let $d \in (X + Y + N)^{\mathrm{diag}}$ and write $(d, d, d) = x + y + n$ with $x \in X$, $y \in Y$, and $n \in N$. Let $E_R := \epsilon_R(\mathrm{Der}(\mu_R))$, $F_R := \epsilon_R(N_R)$, $E_S := \epsilon_S(\mathrm{Der}(\mu_S))$, and $F_S := \epsilon_S(N_S)$. Thus E_R and E_S are the defects of the full derivation algebra, while F_R and F_S are the defects of the nucleus summands N_R and N_S. This gives the relations $F_R \le E_R$ and $F_S \le E_S$. Now view $E_R \otimes Z(S)$ as a subspace of A^2 by $(a, b) \otimes z \mapsto (a \otimes z, b \otimes z)$, and similarly for $Z(R) \otimes E_S$, $F_R \otimes S$, and $R \otimes F_S$. We prove a sequence of intermediate claims before putting it all together.

Claim 1. We prove

$$\epsilon_A(X) = E_R \otimes Z(S), \qquad \epsilon_A(Y) = Z(R) \otimes E_S. \tag{C.2}$$

We prove the direction $\epsilon_A(X) \le E_R \otimes Z(S)$. Every element of $X = \mathrm{Der}(\mu_R) \boxtimes \mathcal{C}(\mu_S)$ is a sum of terms $\delta \boxtimes (\lambda_z, \lambda_z, \lambda_z)$ with $\delta = (\alpha, \beta, \gamma) \in \mathrm{Der}(\mu_R)$ and $z \in Z(S)$. By (C.1), $\epsilon_A\big(\delta \boxtimes (\lambda_z, \lambda_z, \lambda_z)\big) = (\alpha(1_R) \otimes z, \beta(1_R) \otimes z) \in E_R \otimes Z(S)$, so $\epsilon_A(X) \le E_R \otimes Z(S)$.

Conversely, if $\epsilon_R(\delta) = (a, b) \in E_R$ and $z \in Z(S)$, then $(a \otimes z, b \otimes z) = \epsilon_A\big(\delta \boxtimes (\lambda_z, \lambda_z, \lambda_z)\big)$, and such pure tensors span $E_R \otimes Z(S)$. Hence $\epsilon_A(X) = E_R \otimes Z(S)$, and the formula for $\epsilon_A(Y)$ is analogous.

Claim 2. We prove

$$\epsilon_A(Y) + \epsilon_A(N) \le F_R \otimes S, \qquad \epsilon_A(X) + \epsilon_A(N) \le R \otimes F_S. \tag{C.3}$$

Let $Z(R)^2$ means $Z(R) \oplus Z(R) \le R^2$. We show $Z(R)^2 \le F_R$. Let $c, d \in Z(R) \subseteq \mathrm{Nuc}(R)$, then $(\lambda_c, \lambda_c) \in \mathcal{L}(\mu_R)$ and $(\rho_d, \rho_d) \in \mathcal{R}(\mu_R)$, so $(c, 0) = \epsilon_R(\iota_{20}(\lambda_c, \lambda_c))$ and $(0, d) = \epsilon_R(\iota_{10}(\rho_d, \rho_d))$, concluding $(c, d) \in F_R$. Every element of $Y = \mathcal{C}(\mu_R) \boxtimes \mathrm{Der}(\mu_S)$ is a sum of terms $(\lambda_c, \lambda_c, \lambda_c) \boxtimes \delta$ with $c \in Z(R)$ and $\delta = (f, g, h) \in \mathrm{Der}(\mu_S)$. By (C.1),

$$\epsilon_A\big((\lambda_c, \lambda_c, \lambda_c) \boxtimes \delta\big) = (c \otimes f(1_S), c \otimes g(1_S)) \in Z(R)^2 \otimes S \subseteq F_R \otimes S,$$

so $\epsilon_A(Y) \le F_R \otimes S$.

If $t = \iota_{20}(u \boxtimes v)$ with $u = (\alpha, \gamma) \in \mathcal{L}(\mu_R)$ and $v = (f, h) \in \mathcal{L}(\mu_S)$, then $\epsilon_A(t) = (\alpha(1_R) \otimes f(1_S), 0) \in \epsilon_R(\iota_{20}(u)) \otimes S \subseteq F_R \otimes S$. The ι_{10} and ι_{21} cases are the same, and the $R \otimes F_S$ case is analogous.

Claim 3. We prove

$$\epsilon_A(X \cap N) = F_R \otimes Z(S), \qquad \epsilon_A(Y \cap N) = Z(R) \otimes F_S. \tag{C.4}$$

For the inclusion $\epsilon_A(X \cap N) \subseteq F_R \otimes Z(S)$, use $X \cap N \le X$ and $X \cap N \le N$ together with (C.2), (C.3), and Lemma 2.2:

$$\epsilon_A(X \cap N) \subseteq (E_R \otimes Z(S)) \cap (F_R \otimes S) = F_R \otimes Z(S).$$

Let $u \in N_R$ and $z \in Z(S)$. Since $z \in Z(S)$, we have $\lambda_z = \rho_z$, so $(\lambda_z, \lambda_z, \lambda_z) \in \mathcal{C}(\mu_S)$. By Remark C.7, $u \boxtimes (\lambda_z, \lambda_z, \lambda_z) \in X \cap N$. Writing $\epsilon_R(u) = (a, b)$, (C.1) gives $\epsilon_A\big(u \boxtimes (\lambda_z, \lambda_z, \lambda_z)\big) = (a \otimes z, b \otimes z) \in$

$\epsilon_A(X \cap N)$. Since such simple tensors span $F_R \otimes Z(S)$, this gives the reverse inclusion. The second equality is analogous.

These claims imply

$$\epsilon_A(X) \cap \big(\epsilon_A(Y) + \epsilon_A(N)\big) = \epsilon_A(X \cap N), \qquad \epsilon_A(Y) \cap \big(\epsilon_A(X) + \epsilon_A(N)\big) = \epsilon_A(Y \cap N). \tag{C.5}$$

Indeed, $\epsilon_A(X) \cap \big(\epsilon_A(Y) + \epsilon_A(N)\big) \subseteq (E_R \otimes Z(S)) \cap (F_R \otimes S) = F_R \otimes Z(S) = \epsilon_A(X \cap N)$, and the reverse inclusion is immediate since $X \cap N \leq X$ and $X \cap N \leq N$. The second equality is analogous.

Putting it all together. Recall $(d, d, d) = x + y + n \in \mathrm{Ker}(\epsilon_A)$, so $\epsilon_A(x) = -\epsilon_A(y) - \epsilon_A(n) \in \epsilon_A(Y) + \epsilon_A(N)$ and $\epsilon_A(y) = -\epsilon_A(x) - \epsilon_A(n) \in \epsilon_A(X) + \epsilon_A(N)$. Thus by (C.5), $\epsilon_A(x) \in \epsilon_A(X \cap N)$ and $\epsilon_A(y) \in \epsilon_A(Y \cap N)$. Choose $x_N \in X \cap N$ and $y_N \in Y \cap N$ with $\epsilon_A(x_N) = \epsilon_A(x)$ and $\epsilon_A(y_N) = \epsilon_A(y)$.

Let $x_0 := x - x_N$, $y_0 := y - y_N$, and $n_0 := n + x_N + y_N$. Then $x_0 \in X \cap \mathrm{Ker}(\epsilon_A)$, $y_0 \in Y \cap \mathrm{Ker}(\epsilon_A)$, and $n_0 \in N$ with

$$\epsilon_A(n_0) = \epsilon_A(n) + \epsilon_A(x_N) + \epsilon_A(y_N) = \epsilon_A(n) + \epsilon_A(x) + \epsilon_A(y) = \epsilon_A(x + y + n) = 0,$$

so $n_0 \in N \cap \mathrm{Ker}(\epsilon_A)$. By Remark C.6, $x_0 = (d_X, d_X, d_X)$, $y_0 = (d_Y, d_Y, d_Y)$, and $n_0 = (d_N, d_N, d_N)$ for $d_X \in X^{\mathrm{diag}}, d_Y \in Y^{\mathrm{diag}}, d_N \in N^{\mathrm{diag}}$. Hence $d = d_X + d_Y + d_N \in X^{\mathrm{diag}} + Y^{\mathrm{diag}} + N^{\mathrm{diag}}$ as needed. □

Lemma C.9. With the notation above, $X^{\mathrm{diag}} = \mathcal{D}_{\mathrm{alg}}(R) \otimes Z(S)$ and $Y^{\mathrm{diag}} = Z(R) \otimes \mathcal{D}_{\mathrm{alg}}(S)$.

Proof. We prove the formula for X^{diag}. The inclusion $\mathcal{D}_{\mathrm{alg}}(R) \otimes Z(S) \subseteq X^{\mathrm{diag}}$ is immediate, since in the unital case $c \mapsto (\lambda_c, \lambda_c, \lambda_c)$ identifies $Z(S)$ with $\mathcal{C}(\mu_S)$.

For the reverse inclusion, let $u \in X^{\mathrm{diag}}$, so $(u, u, u) \in X$. Choose a basis $z_1, \ldots, z_m$ of $Z(S)$. By Proposition C.2, the triples $(\lambda_{z_i}, \lambda_{z_i}, \lambda_{z_i})$ form a basis of $\mathcal{C}(\mu_S)$. Hence there exist $d_i = (d_{2,i}, d_{1,i}, d_{0,i}) \in \mathrm{Der}(\mu_R)$ such that

$$(u, u, u) = \sum_{i=1}^{m} d_i \boxtimes (\lambda_{z_i}, \lambda_{z_i}, \lambda_{z_i}) = \Big(\sum_{i=1}^{m} d_{2,i} \otimes \lambda_{z_i}, \sum_{i=1}^{m} d_{1,i} \otimes \lambda_{z_i}, \sum_{i=1}^{m} d_{0,i} \otimes \lambda_{z_i}\Big).$$

Evaluating the three components on $r \otimes 1_S$ gives

$$\sum_{i=1}^{m} d_{2,i}(r) \otimes z_i = \sum_{i=1}^{m} d_{1,i}(r) \otimes z_i = \sum_{i=1}^{m} d_{0,i}(r) \otimes z_i$$

for all $r \in R$, so $d_{2,i} = d_{1,i} = d_{0,i}$ for each i. Thus each d_i is of the form $d_i = (\delta_i, \delta_i, \delta_i)$ with $\delta_i \in \mathcal{D}_{\mathrm{alg}}(R)$, and $u = \sum_{i=1}^{m} \delta_i \otimes z_i \in \mathcal{D}_{\mathrm{alg}}(R) \otimes Z(S)$. The formula for Y^{diag} is analogous. □

Proof of Proposition 5.3. By Theorem 5.1, $\mathrm{Der}(\mu_R \otimes \mu_S) = X + Y + N$. By the proof of Lemma C.5, diagonal triples in $\mathrm{Der}(\mu_R \otimes \mu_S)$ are algebra derivations of $R \otimes S$. Using Lemmas C.8 and C.9,

$$\mathcal{D}_{\mathrm{alg}}(R \otimes S) = (X + Y + N)^{\mathrm{diag}} = \mathcal{D}_{\mathrm{alg}}(R) \otimes Z(S) + Z(R) \otimes \mathcal{D}_{\mathrm{alg}}(S) + N^{\mathrm{diag}}.$$

By Theorem 4.1, $N = \iota_{20}(\mathcal{L}(\mu_R \otimes \mu_S)) + \iota_{10}(\mathcal{R}(\mu_R \otimes \mu_S)) + \iota_{21}(\mathcal{M}(\mu_R \otimes \mu_S))$, so Proposition C.4 gives $N^{\mathrm{diag}} = \mathrm{ad}(\mathrm{Nuc}(R \otimes S))$. Finally, the nucleus identity $\mathrm{Nuc}(R \otimes S) = \mathrm{Nuc}(R) \otimes \mathrm{Nuc}(S)$ [Bre17, p. 2] gives

$$N^{\mathrm{diag}} = \mathrm{ad}(\mathrm{Nuc}(R \otimes S)) = \mathrm{ad}(\mathrm{Nuc}(R) \otimes \mathrm{Nuc}(S)).$$

Therefore $\mathcal{D}_{\mathrm{alg}}(R \otimes S) = \mathcal{D}_{\mathrm{alg}}(R) \otimes Z(S) + Z(R) \otimes \mathcal{D}_{\mathrm{alg}}(S) + \mathrm{ad}(\mathrm{Nuc}(R) \otimes \mathrm{Nuc}(S))$, which is the subspace form of Equation (5.2). □

D.1. **Nucleus decomposition.** ↑

Example D.1. We decompose a term in $\mathcal{M}(s \otimes t)$ as sum of terms from $\mathcal{M}(s) \boxtimes \mathcal{M}(t)$. Let $\mathbb{F} = \mathbb{F}_5$, and let $s, t : \mathbb{F}^2 \times \mathbb{F}^2 \rightarrowtail \mathbb{F}$ be bilinear maps with

$$S = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}, \qquad T = I_2 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}.$$

Given matrices

$$X = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}, \qquad Y = \begin{bmatrix} 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 4 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}, \quad \text{and we verify that} X(S \otimes T) = (S \otimes T)Y = \begin{bmatrix} 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}.$$

Let x, y correspond to the matrices X and Y. Then $(s \otimes t) \bullet_2 x = (s \otimes t) \bullet_1 y$, so $(x, y) \in \mathcal{M}(s \otimes t)$. Set $w := \Xi\big((s \otimes t) \bullet_2 x\big) = \Xi\big((s \otimes t) \bullet_1 y\big)$. A direct calculation gives $w = u_1 \otimes v_1 + u_2 \otimes v_2$, where $u_1 = E_{11} + E_{12}$, $v_1 = E_{11}$, $u_2 = E_{22}$, and $v_2 = E_{22}$. Choose

$$\begin{aligned} (\alpha_1, \beta_1) &= (E_{11}, E_{11} + E_{12}), & (f_1, g_1) &= (E_{11}, E_{11}), \\ (\alpha_2, \beta_2) &= (E_{22}, 4E_{12} + E_{22}), & (f_2, g_2) &= (E_{22}, E_{22}). \end{aligned}$$

It suffices to check that $u_i = s \bullet_2 \alpha_i = s \bullet_1 \beta_i$ and $v_i = t \bullet_2 f_i = t \bullet_1 g_i$ for each $i \in \{1, 2\}$. Therefore $x = \kappa_2(\alpha_1 \otimes f_1) + \kappa_2(\alpha_2 \otimes f_2)$, and $y = \kappa_1(\beta_1 \otimes g_1) + \kappa_1(\beta_2 \otimes g_2)$.

To summarize, we've decomposed (x, y) as the sum $(\alpha_1, \beta_1) \boxtimes (f_1, g_1) + (\alpha_2, \beta_2) \boxtimes (f_2, g_2)$ in $\mathcal{M}(s) \boxtimes \mathcal{M}(t)$.

D.2. **Derivation decomposition.** ↑

Example D.2. We decompose $(\delta^{(2)}, \delta^{(1)}, \delta^{(0)}) \in \mathrm{Der}(s \otimes t)$ as sum of terms from $\mathrm{Der}(s) \boxtimes \mathcal{C}(t)$ and $\mathcal{M}(s) \boxtimes \mathcal{M}(t)$. Let $\mathbb{F} = \mathbb{F}_5$. Let $s, t : \mathbb{F}^2 \times \mathbb{F}^2 \rightarrowtail \mathbb{F}$ be given by 2×2 matrices $S = E_{11} + E_{21} + E_{22}$ and $T = I_2$, where E_{ij} denote matrix units. Consider $\Delta = (\Delta^{(2)}, \Delta^{(1)}, \Delta^{(0)})$.

$$\Delta^{(2)} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \end{bmatrix}, \qquad \Delta^{(1)} = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}, \qquad \Delta^{(0)} = \begin{bmatrix} 1 \end{bmatrix},$$

for which we verify $\Delta^{(2)}(S \otimes T) + (S \otimes T)\Delta^{(1)} = (S \otimes T)^{\Delta^{(0)}}$. Let $\delta = (\delta^{(2)}, \delta^{(1)}, \delta^{(0)}) \in \mathrm{Der}(s \otimes t)$ be the operators corresponding to Δ. Set $p := (s \otimes t) \bullet_2 \delta^{(2)}$, $q := (s \otimes t) \bullet_1 \delta^{(1)}$, and $r := (s \otimes t) \bullet_0 \delta^{(0)}$, where $p + q = r$. A direct calculation gives

$$p = (E_{11} + E_{21}) \otimes I_2 + E_{22} \otimes E_{11}, \qquad q = E_{22} \otimes I_2 - E_{22} \otimes E_{11}, \qquad r = (E_{11} + E_{21} + E_{22}) \otimes I_2 + 0 \otimes E_{11},$$

so $p + q = r$. We read off local equalities:

$$\begin{aligned} p &= a_1 \otimes x_1 + a_2 \otimes x_2, & a_1 &= E_{11} + E_{21},\ x_1 = I_2,\ a_2 = E_{22},\ x_2 = E_{11}, \\ q &= b_1 \otimes y_1 + b_2 \otimes y_2, & b_1 &= E_{22},\ y_1 = I_2,\ b_2 = -E_{22},\ y_2 = E_{11}, \\ r &= c_1 \otimes z_1 + c_2 \otimes z_2, & c_1 &= S,\ z_1 = I_2,\ c_2 = 0,\ z_2 = E_{11}. \end{aligned}$$

They are chosen so $a_i \otimes x_i + b_i \otimes y_i = c_i \otimes z_i$ for all $i \in \{1, 2\}$. Let $\alpha_D = E_{11} + E_{21},\ f_D = I_2,\ \beta_D = E_{22},\ g_D = I_2,\ \gamma_D = 1,\ h_D = 1$. Then

$$a_1 = s \bullet_2 \alpha_D,\ x_1 = t \bullet_2 f_D, \qquad b_1 = s \bullet_1 \beta_D,\ y_1 = t \bullet_1 g_D, \qquad c_1 = s \bullet_0 \gamma_D,\ z_1 = t \bullet_0 h_D,$$

with $(\alpha_D, \beta_D, \gamma_D) \in \mathrm{Der}(s)$ and $(f_D, g_D, h_D) \in \mathcal{C}(t)$.

Similarly, let $\alpha_N = 4E_{21} + E_{22},\ f_N = E_{11},\ \beta_N = E_{22},\ g_N = E_{11}$. Then $a_2 = s \bullet_2 \alpha_N,\ x_2 = t \bullet_2 f_N$, and $b_2 = -s \bullet_1 \beta_N,\ y_2 = t \bullet_1 g_N$, meaning $(\alpha_N, \beta_N) \in \mathcal{M}(s)$, $(f_N, g_N) \in \mathcal{M}(t)$.

Therefore

$$\delta^{(2)} = \kappa_2(\alpha_D \otimes f_D) + \kappa_2(\alpha_N \otimes f_N) \quad \delta^{(1)} = \kappa_1(\beta_D \otimes g_D) - \kappa_1(\beta_N \otimes g_N), \quad \delta^{(0)} = \kappa_0(\gamma_D \otimes h_D).$$

This demonstrates $(\delta^{(2)}, \delta^{(1)}, \delta^{(0)})$ is a sum of a term from $\mathrm{Der}(s) \boxtimes \mathcal{C}(t)$ and a term from $\iota_{21}\big(\mathcal{M}(s) \boxtimes \mathcal{M}(t)\big)$.