

**Theorem (2.3.67d).** *Let  $A$ , and  $B$  be sets with universal set  $U$ . Let  $f_{A \oplus B}$  be the characteristic function  $f_{A \oplus B} : U \Rightarrow \{0, 1\}$ . Let  $f_A$  be the characteristic function  $f_A : U \Rightarrow \{0, 1\}$ . Let  $f_B$  be the characteristic function  $f_B : U \Rightarrow \{0, 1\}$ .  $f_{A \oplus B}(x) = f_A(x) + f_B(x) - 2f_A(x)f_B(x)$ .*

*Proof.* There are two major cases to consider, each consisting of two sub cases. The major cases are where  $x$  is an element in  $A \oplus B$ , and the negation of that statement.

(i) Let  $x$  be an element in  $A \oplus B$ . By the definition for characteristic functions,  $f_{A \oplus B}(x) = 1$ . Since the definition for set symmetric difference says  $[(x \in A) \wedge (x \notin B)] \vee [(x \notin A) \wedge (x \in B)]$ , there are two sub cases that need to be taken under consideration.

(a) Suppose  $(x \in A) \wedge (x \notin B)$ . By the definition for characteristic functions  $f_A(x) = 1$  and  $f_B(x) = 0$ . This means that  $f_A(x) + f_B(x) - 2f_A(x)f_B(x) = 1 + 0 - 2(1)(0) = 1$ .

(b) Suppose  $(x \notin A) \wedge (x \in B)$ . Without loss of generality we arrive at the same result as that of case (a).

Thus, if  $x$  is an element in  $A \oplus B$ ,  $f_{A \oplus B}(x) = f_A(x) + f_B(x) - 2f_A(x)f_B(x)$ .

(ii) Suppose it were not the case that  $x$  were an element in  $A \oplus B$ . Then (c)  $x$  must either be an element in the intersection of  $A$  and  $B$ , or (d)  $x$  must be in the universe minus  $A \cup B$ .

(c) Suppose  $x \in (A \cap B)$ . By the definition for characteristic functions  $f_{A \oplus B}(x) = 0$ ,  $f_A(x) = 1$  and  $f_B(x) = 1$ . Thus,  $f_A(x) + f_B(x) - 2f_A(x)f_B(x) = 1 + 1 - 2(1)(1) = 0$ .

(d) Suppose  $x \in [U - (A \cup B)]$ . In this case, by the definition for characteristic functions,  $f_{A \oplus B}(x) = 0$ ,  $f_A(x) = 0$  and  $f_B(x) = 0$ . So,  $f_A(x) + f_B(x) - 2f_A(x)f_B(x) = 0 + 0 - 2(0)(0) = 0$ .

Thus, if  $x$  is not an element in  $A \oplus B$ ,  $f_{A \oplus B}(x) = f_A(x) + f_B(x) - 2f_A(x)f_B(x)$  is still a true statement; concludes the proof. ■