

# The Whitehall-Robins Report

December 2010 - Volume 19, Number 4

## The Role of Omega-3 Fatty Acids throughout the Life Cycle: Part 1

**Bruce J. Holub, PhD, University Professor Emeritus,  
Department of Human Health and Nutritional Sciences, University of Guelph  
and Scientific Director, DHA/EPA Omega-3 Institute, University of Guelph Research Park**

Canadian diets contain relatively high levels of polyunsaturated fats (fatty acids) representing approximately 4-6% of total energy intakes with 90% of the polyunsaturated fatty acids (PUFA) being consumed as the omega-6 PUFA (also known as n-6 PUFA)<sup>1</sup>. These are consumed mostly as linoleic acid via common vegetable oils plus minor amounts of arachidonic acid in animal-based food products. The so-called omega-3 PUFA (n-3 PUFA) typically represent only 10% of the total PUFA intake with approximately 90% of the omega-3 PUFA being consumed (approximately 1.6 gm daily) as the short-chain omega-3 fatty acid known as LNA (alpha-linolenic acid) as found in plant food sources such as soybean oil, canola oil, flax, English walnuts, others. The fish/fish oil-based omega-3 fatty acids represent very minor components (approximately 10%) of the total omega-3 intakes. These long-chain omega-3 fatty acids consist of EPA (eicosapentaenoic acid, 20:5 n-3) plus DHA (docosahexaenoic acid, 22:6 n-3) and are found in varying amounts and ratios in fish/seafood with lesser amounts being consumed in other animal-based foods or processed foods wherein DHA/EPA are added. The metabolic conversion of dietary LNA in the liver to EPA/DHA is both highly variable between individuals and very limited with overall efficiencies in adults (females and males) for LNA to DHA averaging only 3-4%. Thus, the direct consumption of pre-formed DHA/EPA in foods/diets is the most direct means for their provision to the body for health. Typical adult intakes of EPA plus DHA (combined) in Canada range from 100-150 mg/day which reflects very modest fish intakes of approximately one serving every 7-10 days. Much higher intakes are found (approaching 1200 mg/day) in certain countries such as Japan which have very high fish/seafood consumption. The present report will update the reader on recent evidence-based studies which focus on the potential health benefits of DHA/EPA omega-3 when consumed as fish/seafood, fish/algal oils, functional foods, and/or via supplementation in healthy or at risk populations.

### **Omega-3 Fatty Acids during Pregnancy and Lactation**

Docosahexaenoic acid (DHA) is recognized as a physiologically-essential fatty acid in the brain and retina of the eye where it is found in high levels as the predominant member of the omega-3 fatty acids found in the cell membranes therein. The high level of DHA, with its special structural and physical-chemical properties, supports optimal cognitive performance and visual acuity in the brain and retina, respectively, via the mediation of nerve transmission and other critical functions. DHA accumulates in neural and other tissues throughout pregnancy with the last trimester being a particularly active period of growth and neurodevelopment for the infant. DHA continues to accrue at a high rate via lactation during early infant development after birth in nervous and other tissues with the amount of DHA in the brain increasing approximately 30-fold from about 24 weeks of gestation to about 2 years of age<sup>2</sup>.

The ALSPAC study<sup>3</sup> found a significantly higher risk of sub-optimal outcomes for various development parameters (verbal IQ, behaviours) up to 8 yrs later in those children from mothers who consumed no

seafood as compared to those consuming 340 gms or more per week during pregnancy (340 gms/week estimated to provide an average of approx. 160 mg DHA/day). A recent Spanish study suggested that moderately high intakes of fish during pregnancy may benefit neurodevelopment in children at age 4 years<sup>4</sup>. Daily intake levels of 294 mg DHA as enhanced via supplementation by mothers (versus only 80 mg/day) were found to significantly improve infant visual acuity at four but not six months of age<sup>5</sup>. The PERLIP project (as charged by the European Commission) has recommended that pregnant women should aim to achieve an average intake of at least 200 mg DHA/day<sup>6</sup> which contrasts with an average intake of only 82 mg/day for pregnant Canadian women as directly assessed<sup>7</sup>. Very recently, a randomized controlled trial from Australia reported that DHA supplementation during the last half of pregnancy did not reduce postpartum depression in the mothers or selected measures of cognitive and language development in children as assessed at 18 months<sup>8</sup>. It remains to be determined whether more prolonged supplementation during pregnancy and throughout lactation, particularly in mothers with a low DHA

status, may have resulted in beneficial outcomes to the children based on a wide array of cognitive and visual measures.

### **Omega-3 Fatty Acids during Infancy, Childhood and Adolescence**

Some controlled clinical trials in full-term infants have indicated that the presence of DHA in breast milk at higher levels or in infant formula (eg., approx. 0.32% of milk fat which is close to the worldwide human milk level) results in improved cognitive and visual outcomes<sup>9</sup>. A literature review has supported an overall improvement in visual developments in term infants fed formula providing at least 100 mg daily<sup>10</sup> which can be expected by breast milk or infant formula having approx. 0.35% of milk fatty acids as DHA. Recently, the World Association of Perinatal Medicine recommended that lactating women consume at least 200 mg DHA/day<sup>11</sup> while the earlier ISSFAL-supported NIH Workshop advised at least 300 mg DHA/day. Such intakes can be expected to yield levels of approx. 0.30-0.35% DHA in breast milk fat.

The accretion of DHA in the brain progressively increases in the cerebral cortex up to 18 years of age<sup>12</sup>. The evidence

for the potential benefits of DHA/EPA supplementation in healthy children older than 2 years of age is generally promising in some studies but not yet conclusive. By supplementing healthy 4-year-old children with 400 mg DHA (or placebo) per day for 4 months, the higher blood levels of DHA which resulted were significantly and positively associated with improved scores on the Peabody Picture Vocabulary Test (a test of listening comprehension for the spoken word in Standard English)<sup>13</sup>. Another recent study on older children (10-12 years of age) which provided 400 or 800 mg supplemental DHA per day for 8 weeks did not find any significant beneficial effect on brain function in healthy children<sup>14</sup>. A formulated functional food (bread spread containing fish flour from a marine source) providing an average intake of 182 mg per day of DHA and EPA combined over 6 months was recently reported to moderately improve the learning ability and memory of children ages 7-9 years<sup>15</sup>. Various studies using DHA and EPA supplementation in attention-deficit-hyperactivity disorder have been inconsistent<sup>16</sup> while preliminary investigations on childhood depression<sup>17</sup> and neurological outcomes in children with phenylketonuria<sup>18</sup> have been encouraging.

Our lab has determined fatty acid intakes in a population of Canadian children aged 4-8 years by direct diet quantitation<sup>19</sup>. Very low intakes of DHA and EPA were found averaging 54 and 38 mg/child/day, respectively. Only 22% of the children met

the suggested adequate intake (DHA plus EPA) based on the Dietary Reference Intakes from the Institute of Medicine (Washington, DC) and only 51% met the recommended intakes for long-chain omega-3 fatty acids for children from Australia and New Zealand. The wide 'nutrition gap' between actual versus recommended intakes of DHA and EPA for young children emphasizes the need to recommend increased consumption of appropriate fish and seafood, fortified foods, and possibly supplements containing DHA and EPA.

### Summary and Conclusions

The overall evidence-based research to date support the importance of adequate long-chain omega-3 fatty acid intakes (particularly DHA) during pregnancy and lactation to support optimal cognitive and visual development both for the infant and throughout childhood. In addition, direct health benefits to the mother may also be forthcoming from

increased intakes of fish/seafood or alternative sources of DHA/EPA including various foods enriched in omega-3 fatty acids or supplementation. Unfortunately, almost all the research to date has not differentiated between those who may be at particularly high risk of omega-3 deprivation for optimal health and others (mother and child) who may have a sufficiently adequate omega-3 status at the physiological level. It is expected that specific recommendations for DHA and/or (DHA plus EPA) intakes in mg/day will be forthcoming from governmental sources in North America, in addition to current target intakes for LNA, in keeping with advisories from international committees and agencies.

Ongoing updates on forthcoming evidence-based research on DHA/EPA omega-3 in health and disease can be freely-accessed via the DHA/EPA Omega-3 Institute at [www.dhaomega3.org](http://www.dhaomega3.org).

The Whitehall-Robins Report is a Pfizer Consumer Healthcare, a division of Pfizer Canada Inc., publication that focuses on current issues on the role of vitamins and minerals in health promotion and disease prevention. Complimentary copies are distributed to Canadian health care professionals active or with a special interest in nutrition. Each issue is written and/or reviewed by independent health care professionals with expertise in the chosen topic.

**Editor: Pfizer Consumer Healthcare, a division of Pfizer Canada Inc.**

If you have any comments about the Whitehall-Robins Report or would like to be added to the mailing list, please write to:

**The Editor: The Whitehall-Robins Report, 5975 Whittle Road, Mississauga, ON L4Z 3M6**

For back issues or further information, visit the Centrum professional section of [www.centrum.ca](http://www.centrum.ca).

© 2010 – December. May be reproduced without permission provided source is recognized.

**References** 1. Holub BJ. Omega-3 fatty acids for health care. Whitehall-Robins Report 2005. 2. Mitmesser SH. *et al.* Roles of long-chain polyunsaturated fatty acids in the term infant: developmental benefits. Neonatal Net 2007; 229-234. 3. Hibbeln JR. *et al.* Maternal seafood consumption in pregnancy and neurodevelopmental outcomes in childhood (ALSPAC study): an observational cohort study. The Lancet 2007; 369:575-585. 4. Mendez MA. *et al.* Maternal fish and other seafood intakes during pregnancy and child neurodevelopment at age 4 years. Public Health Nutr 2009; 12:1702-1710. 5. Judge M. *et al.* A docosahexaenoic acid-functional food during pregnancy benefits infant visual acuity at four but not six months of age. Lipids 2007; 42:117-122. 6. Koletzko B. *et al.* Dietary fat intakes for pregnant and lactating women. Br J Nutr 2007; 98:873-877. 7. Denomme J. *et al.* Directly quantitated dietary (n-3) fatty acid intakes of pregnant Canadian women are lower than current dietary recommendations. J Nutr 2005; 135: 206-211. 8. Makrides M. *et al.* Effect of DHA supplementation during pregnancy on maternal depression and neurodevelopment of young children. JAMA 2010; 304(15):1675-1683. 9. Hoffman DR. *et al.* Toward optimizing vision and cognition in term infants by dietary docosahexaenoic and arachidonic acid supplementation: a review of randomized controlled trials. PLEFA 2009; 81: 151-158. 10. Eilander A. *et al.* Effects of n-3 long chain polyunsaturated fatty acids supplementations on visual and cognitive development throughout childhood: A review of human studies. Prostaglandins Leukot Essent Fatty Acids 2007; 76:189-203. 11. Koletzko B. *et al.* The roles of long-chain polyunsaturated fatty acids in pregnancy, lactation and infancy: review of current knowledge and consensus recommendations. J Perinat Med 2008; 36:548-549. 12. Carver J. *et al.* The relationship between age and the fatty acid composition of cerebral cortex and erythrocytes in human subjects. Brain Res Bull 2001; 56:79-85. 13. Ryan AS. *et al.* Assessing the effect of docosahexaenoic acid on cognitive functions in healthy, preschool children: a randomized, placebo-controlled, double-blind study. Clin Pediatr 2008; 47:355-62. 14. Kennedy DO. *et al.* Cognitive and mood effects of 8 weeks' supplementation with 400 mg or 1000 mg of the omega-3 essential fatty acid docosahexaenoic acid (DHA) in healthy children aged 10-12 years. Nutr Neurosci 2009; 12:48-56. 15. Dalton A. *et al.* A randomized control trial in schoolchildren showed improvement in cognitive function after consuming a bread spread, containing fish flour from a marine source. Prostaglandins Leukot Essent Fatty Acids 2009; 80: 143-149. 16. Raz R. *et al.* Essential fatty acids and attention-deficit-hyperactivity disorder: a systematic review. Dev Med Child Neurol 2009; 51: 580-592. 17. Nemets H. *et al.* Omega-3 treatment of childhood depression: a controlled, double-blind pilot study. Am J Psychiatry 2006; 163:1098-1100. 18. Koletzko B. *et al.* Does dietary DHA improve neural function in children? Observations in phenylketonuria. Prostaglandins Leukot Essent Fatty Acids 2009; 81:159-164. 19. Madden SM. *et al.* Direct diet quantification indicates low intakes of (n-3) fatty acids in children 4 to 8 years old. J Nutr 2009; 139:528-532.