
Narrowing the Edge Folkman Number Bounds

Independent Study Proposal

Christopher Wood

Advisor: Professor Stanisław Radziszowski

February 25, 2013

1 Background

Edge Folkman numbers, first introduced by Folkman in 1970 [1], are concerned with the study of graphs in which a monochromatic coloring of a particular subgraph always exists. We write $G \rightarrow (a_1, \dots, a_k; p)^e$ if for every edge coloring of an undirected simple graph G not containing K_p , there exists a monochromatic K_{a_i} in color i for some $i \in \{1, \dots, k\}$. The edge Folkman number is defined as $F_e(a_1, \dots, a_k) = \min\{|V(G)| : G \rightarrow (a_1, \dots, a_k; p)^e\}$. In 1970 Folkman proved that for all $k > \max(s, t)$, edge- and vertex- Folkman numbers $F_e(s, t; k)$ and $F_v(s, t; k)$ exist. Prior to this, Erdos and Hajnal pose the problem of finding $F_e(3, 3; 4)$, which can be informally stated as the following [2]:

What is the order of the smallest K_4 -free graph for which any 2-coloring of its edges must contain at least one monochromatic triangle?

This is equivalent to finding the smallest K_4 -free graph that is not the union of two triangle-free graphs. Since the proposition of this problem, there has been a significant amount of work towards aimed at narrowing the upper and lower bounds of $F_e(3, 3; 4)$. Table 1 enumerates the work on this problem and leads us to the current state of the field.

Table 1: History of $F_e(e, e; 4)$

Year	Bounds	Who	Ref.
1967	any?	Erdős-Hajnal	[2]
1970	exist	Folkman	[1]
1972	≥ 10	Lin	[3]
1975	$\leq 10^{10}?$	Erdős offers \$100 for proof	
1986	$\leq 8 \times 10^{11}$	Frankl-Rödl	[5]
1988	$\leq 3 \times 10^9$	Spencer	[6]
1999	≥ 16	Piwakowski et al (implicit)	[7]
2007	≥ 19	Radziszowski-Xu	[8]
2008	≤ 9697	Lu	[9]
2008	≤ 941	Dudek-Rödl	[10]
2012	≤ 786	Lange et al	TODO
2012	$\leq 100?$	Graham offers \$100 for proof	

2 Problems

2.1 Lower Bound

The current lower bound for $F_e(3, 3; 4)$ stands at 19 [8]. The proof used for this bound, which relies on the existence of independent sets of size k in a graph G on 18 vertices, where $k = 4$, cannot easily be extended to graphs on 19 vertices. It is estimated that more than 10^{19} nonisomorphic, K_4 -free graphs on 19 vertices exist [4].

The proof for this bound relies on the fact that $H \rightarrow (3, 3; 4)^v \Rightarrow H + x \rightarrow (3, 3; 5)^e$ and $F_e(3, 3; 5) = 15$. More specifically, using all nonisomorphic graphs on 18 vertices with contained independent sets of size $k = 4$, the authors were show that, for each candidate graph G , the graph

2.2 Upper Bound

The second segment of this project is to address the conjecture that $F_e(3, 3; 4) \leq 127$ by examining the graph $G_{127} = (\mathbb{Z}_{127}, E = \{(x, y) | x - y = \alpha^3 \pmod{127}\})$. It was first conjectured that $G_{127} \rightarrow (3, 3)^e$ in [8], where the intuition for this conjecture is that G_{127} has a large number of triangles and many small dense subgraphs. We will attempt to prove or disprove this conjecture by showing that $G_{127} \not\rightarrow (3, 3)^e$ using subgraph H extensions. More specifically, we will identify subgraphs H witnessing $H \not\rightarrow (3, 3)^e$ and then attempt to extend H to all of G . However, due to the computationally intensive nature of this procedure, we will reduce $\{H | H \not\rightarrow (3, 3)^e\}$ to 3-SAT and leverage the power of SAT solvers such as Minisat [11] and zChaff [12]. The reduction works by mapping the edges in $E(H)$ to variables in $\phi_G \in 3\text{-SAT}$, and for edge adding the following clauses to ϕ_G :

$$(x + y + z) \wedge (\bar{x} + \bar{y} + \bar{z})$$

Clearly, $H \not\vdash (3,3)^e \Leftrightarrow \phi_G$ is satisfiable.

TODO:

3 Summary of Proposed Study

The work for this project will be two-fold.

1. addressing lower bound
2. addressing upper bound

A major part of this work will be identifying candidate subgraphs H for extension that can easily be solved using the aforementioned 3-SAT solvers.

4 Project Goals and Deliverables

The following project goals have been identified for this work

- Research current methods and potential techniques for computing the upper and lower bounds of Folkman numbers (in particular, $F_e(3,3;4)$).
- Implement software to aid in the computation of Folkman number bounds.
- Begin a survey paper outlining the history of Folkman number research [13].

By achieving these goals I will generate the following deliverables:

- Publication-ready paper and more in-depth progress report for the entire project
- Report that weighs the performance of various SAT solvers for solving this arrowing problem with G127
- Report discussing the subgraph pruning techniques used for the lower bound

References

- [1] Jon Folkman. Graphs with monochromatic complete subgraphs in every edge coloring. *SIAM Journal of Applied Mathematics*. 18 (1970), 19-24.
- [2] P. Erdős, A. Hajnal. Research problem 2-5. *Journal of Combinatory Theory*, 2 (1967), 104.
- [3] Shen Lin. On Ramsey numbers and K_r -coloring of graphs. *Journal of Combinatorial Theory, Series B*, 12:82-92, 1972.

- [4] Brendan D. McKay and Stanisław P. Radziszowski. $R(4,5) = 25$. *Journal of Graph Theory*, 19(1995), 309-322.
- [5] Peter Frankl and Vojtech Rödl. Large triangle-free subgraphs in graphs with K_4 . *Graphs and Combinatorics*, 2:135-144, 1986.
- [6] Joel Spencer. Three hunder million points suffice. *Journal of Combinatorial Theory, Series A*, 49(2):210-217, 1988. Also see erratum by M. Hovey in Vol. 50, p. 323.
- [7] Konrad, Piwakowski, Stanisław P. Radziszowski, and Sebastian Urbański. Computation of the Folkman Number $F_e(3,3;5)$. *Journal of Graph Theory*, 32:41-49, 1999.
- [8] Stanisław P. Radziszowski and Xiaodong Xu. On the Most Wanted Folkman Graph. *Ge-combinatorics*, 16(4):367-381, 2007.
- [9] Linyuan Lu. Explicit Construction of Samll Folkman Graphs. *SIAM Journal on Discrete Mathematics*, 21(4):1053-1060, January 2008.
- [10] Andrzej Dudek and Vojtech Rödel. On the Folkman Number $f(2,3,4)$. *Experimental Mathematics*, 17(1):63-67, 2008.
- [11] Niklas Sorensson and Niklas Een. Minisat v1.13 - A SAT Solver with Conflict-Clause Minimization. *SAT 2005*, (2005): 53.
- [12] Yogesh Mahajan, Fu Zhaohui, and Sharad Malik. Zchaff2004: An Efficient SAT Solver. *Theory and Applications of Satisfiability Testing*. Springer Berlin/Heidelberg, 2005.
- [13] Stanisław P. Radziszowski. Small Ramsey Numbers. *Electronic Journal of Combinatorics*, 1 (2011).