

Avg SG: **114 mg/dL**

Time in range: **18% Above** 150 mg/dL


Estimated A1C ⁽¹⁾: **5.6%** calculated from SG values

66% in target range

16% Below 70 mg/dL

OBSERVED PATTERNS & SOME POSSIBLE CAUSES ⁽²⁾

1	2	3
Low SG Post-dinner 5:00 PM - 8:00 PM	High SG Overnight 11:00 PM - 6:00 AM	High SG Post-breakfast 6:00 AM - 10:00 AM
2 out of 4 days excursions observed: 1 day(s) 50 - 70 mg/dL 1 day(s) < 50 mg/dL	2 out of 4 days excursions observed: 2 day(s) 150 - 250 mg/dL 0 day(s) > 250 mg/dL	2 out of 4 days excursions observed: 1 day(s) 180 - 250 mg/dL 1 day(s) > 250 mg/dL
Oral medication(s) too high or incorrectly timed?	Oral medication(s) missed, too low, or incorrectly timed?	Oral medication(s) missed, too low, or incorrectly timed?
Basal insulin injections too high?	Basal insulin injection in evening(s) too low?	Pre-breakfast insulin incorrectly timed, too low, or missed?
Pre-dinner insulin too high?	Insulin to carbohydrate ratio not optimal in prior evening(s)?	Insulin to carbohydrate ratio not optimal for pre-breakfast insulin?
Insulin to carbohydrate ratio not optimal for pre-dinner insulin?	High calorie or high fat foods in prior evening(s)?	High calorie or high carbohydrate foods?
Less food intake than usual?	Late evening snack?	
Exercised around dinner?		



(1) Estimated A1C does not replace Lab measurement and is calculated from limited SG data.

(2) Suggested considerations are limited and do not replace the opinion or advice of the healthcare provider. Please see User Guide on how patterns and possible causes are identified.