
## Polyrate 2023:A Computer Program for the Calculation of Chemical Reaction Rates for Polyatomics. New Version Announcement



Polyrate 2023:A Computer Program for the Calculation of Chemical Reaction Rates for Polyatomics. New Version Announcement

Rubén Meana-Pañeda, Jingjing Zheng, Junwei Lucas Bao, Shuxia Zhang, Benjamin J. Lynch et al.


PII: S0010-4655(23)00278-3

DOI: https://doi.org/10.1016/j.cpc.2023.108933

Reference: COMPHY 108933

To appear in: Computer Physics Communications

Received date: 5 August 2023 Revised date: 5 September 2023 Accepted date: 8 September 2023



Please cite this article as: R. Meana-Pañeda, J. Zheng, J.L. Bao et al., *Polyrate 2023*:A Computer Program for the Calculation of Chemical Reaction Rates for Polyatomics. New Version Announcement, *Computer Physics Communications*, 108933, doi: https://doi.org/10.1016/j.cpc.2023.108933.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2023 Published by Elsevier.

# New version announcement to be submitted to *Computer Physics Communications*

# **Polyrate 2023:** A Computer Program for the Calculation of Chemical Reaction Rates for Polyatomics. New Version Announcement

#### Rubén Meana-Pañeda,

Department of Chemistry, Chemical Theory Center, and Minnesota Supercomputing Institute, University of Minnesota, Minneapolis, MN 55455-0431, USA and Center for Structural Biology, Center for Cancer Research, National Cancer Institute, Frederick, MD, 21702, USA

### Jingjing Zheng, Junwei Lucas Bao, Shuxia Zhang,

Benjamin J. Lynch, José C. Corchado, Yao-Yuan Chuang, Patton L. Fast, Wei-Ping Hu, Yi-Ping Liu, Gillian C. Lynch, Kiet A. Nguyen, Charles F. Jackels, Antonio Fernández-Ramos, Benjamin A. Ellingson, Vasilios S. Melissas, Jordi Villà, Ivan Rossi, Elena L. Coitiño, Jingzhi Pu, Titus V. Albu

Department of Chemistry, Chemical Theory Center, and Minnesota Supercomputing Institute, University of Minnesota, Minneapolis, MN 55455-0431, USA

### Rui Ming Zhang and Xuefei Xu\*

Center for Combustion Energy, Department of Energy and Power Engineering, and Key Laboratory for Thermal Science and Power Engineering of Ministry of Education, Tsinghua University, Beijing 100084, China

#### Artur Ratkiewicz

Institute of Chemistry, University of Bialystok, Poland

#### Rozeanne Steckler

Northwest Alliance for Computational Science & Engineering, Oregon State University, Corvallis, OR 97331, USA

#### Bruce C. Garrett

Environmental Molecular Sciences Laboratory, Pacific Northwest National Laboratory, Richland, WA 99352 USA

#### Alan D. Isaacson

Department of Chemistry and Biochemistry, Miami University, Oxford, OH 45056, USA

#### and Donald G. Truhlar\*

Department of Chemistry, Chemical Theory Center, and Minnesota Supercomputing Institute, University of Minnesota, Minneapolis, MN 55455 USA

<sup>\*</sup>email: xuxuefei@gmail.com, truhlar@umn.edu

#### **ABSTRACT**

Polyrate is a suite of computer programs for the calculation of chemical reaction rates of polyatomic species (including atoms and diatoms as special cases) by variational transition state theory (VTST); conventional transition state theory is also supported. *Polyrate* can calculate the rate constants for both bimolecular reactions and unimolecular reactions, and it can be applied to reactions in the gas phase, liquid-solution phase, or solid-state and to reactions at gas-solid interfaces. *Polyrate* can perform VTST calculations on gas-phase reactions with both tight and loose transition states. For tight transition states it uses the reaction-path (RP) variational transition state theory developed by Garrett and Truhlar, and for loose transition states it uses variable-reaction-coordinate (VRC) variational transition state theory developed by Georgievskii and Klippenstein. The RP methods used for tight transition states are conventional transition state theory, canonical variational transition state theory (CVT), and microcanonical variational transition state theory (µVT) with multidimensional semiclassical approximations for tunneling and nonclassical reflection. For VRC calculations, rate constants may be calculated for canonical or microcanonical ensembles or energy-andtotal-angular-momentum resolved microcanonical ensembles. Pressure-dependent rate constants for elementary reactions can be computed using system-specific quantum RRK theory (SS-QRRK) with the information obtained from high-pressure-limit VTST calculation as input by using the SS-QRRK utility code. Alternatively, Polyrate 2023 may be interfaced with TUMME 2023 for a master-equation treatment of pressure dependence or to obtain phenomenological rate constants for complex mechanisms. Potential energy surfaces may be analytic functions evaluated by subroutines, or they may be implicit surfaces defined by electronic structure input files or interface subroutines containing energies, gradients, and force constants (Hessians). For the latter, *Polyrate* can be used in conjunction with various interfaces to electronic structure programs for direct dynamics, and it has routines designed to make such interfacing straightforward. *Polyrate* supports six options for direct dynamics, namely (i) straight single-level direct dynamics, (ii) zero-order interpolated variational transition state theory (IVTST-0), (iii) first-order interpolated variational transition state theory (IVTST-1), (iv) interpolated variational transition state theory by mapping (IVTST-M), (v) variational transition state theory with interpolated single-point energies (VTST-ISPE), and (vi) variational transition state theory with interpolated optimized corrections (VTST-IOC). Polyrate can handle multistructural and torsional-potential anharmonicity in conjunction with the MSTor program. Polyrate 2023 contains 112 test runs, and 46 of these are for direct dynamics calculations; 85 of the test runs are single-level runs, and 27 are dual-level calculations.

#### Keywords:

chemical reaction rates, stationary-point analysis, reaction path, variational transition state theory, multidimensional tunneling, kinetics, surface science, pressure-dependence

### New version program summary

Program Title: Polyrate 2023

CPC Library link to program files: https://doi.org/10.17632/vpf3y9cbhf.1

Developer's repository link: https://doi.org/10.5281/zenodo.8213313

Licensing Provisions: Apache-2.0 for the program and CC-BY-4.0 for the manual

Programming language: Fortran 90

Does the new version supersede the previous version? Yes

Journal references of previous versions: Comput. Phys. Commun. 47 (1987) 91 [1]; Comput. Phys. Commun. 71 (1992) 235 [2]; Comput. Phys. Commun. 88 (1995) 341 [3]

Nature of problem: Polyrate 2023 can calculate unimolecular and bimolecular chemical reaction rates of polyatomic species (and also of atoms and diatoms as special cases) by using variational transition state theory with multidimensional tunneling [4-8] or by conventional transition state theory [9]. Rate constants may be calculated for canonical or microcanonical ensembles with multidimensional semiclassical approximations for tunneling and nonclassical reflection and for reactions in the gas phase or solid state or at gas-solid interfaces. In addition, rates may be calculated for vibrationally adiabatic and vibrationally diabatic state-selected reactions in which one or more of the vibrational modes is restricted to the ground state or to the first excited state, while the translational, rotational, and remaining vibrational modes are treated thermally [10,11]. The program may also be used to find stationary geometries of reactants, products, and transition states and to calculate reaction paths, Arrhenius parameters, and equilibrium constants.

Solution method: Energies, gradients, and Hessians are obtained from analytic potential energy functions or from electronic structure input data, where the input data may be supplied in files or by direct interface to electronic structure codes. A number of methods are available for calculating the reaction path. Nontorsional anharmonicity may be included under the assumption of independent generalized normal modes. Torsional potential anharmonicity and multiple-structure anharmonicity can be included along the reaction path using the formulas of the multi-structural torsional (MS-T) method with one input structure [12-16]. Small-curvature [2,17], large-curvature [18,19], and optimized multidimensional [20,21] semiclassical tunneling methods are available as options to calculate tunneling contributions to transmission coefficients. Pressure-dependent rate constants for elementary reactions can be computed using system-specific quantum Rice–Ramsperger–Kassel theory (SS-QRRK) [22]. Phenomenological rate constants for complex reactions can be obtained in conjunction with the TUMME 2023 program [23-25].

Interfaces to other programs: Polyrate 2023 can run as a standalone program, and many interface options are also available [24], including: CRATE, GAMESSPLUSRATE, GAUSSRATE, JAGUARATE, MC-TINKERRATE, MORATE, MULTILEVELRATE, NWCHEMRATE.

Reason for new version: The new version greatly increases the functionality, stability, and

accuracy; the code has been cleaned up, and some bugs have been fixed. *Summary of revisions:* 

- 1. The variable-reaction-coordinate method with either single-faceted or multi-faceted dividing surfaces [26–30] has been implemented for barrierless association reactions and dissociation reactions without an intrinsic barrier, i.e., reactions without a barrier when considered in the downhill direction. This implementation is parallelized using MPI. In calculating the total number of available states for this method, the integration over the projection on the principal axis of the total angular momentum is calculated analytically. The user now has a choice of using the original variational transition state theory based on a reaction path (VTST-RP) or using variational transition state theory based on variable reaction coordinate (VTST-VRC).
- 2. Options to carry out VTST-RP calculations with generalized normal modes in curvilinear coordinates in nonredundant [31–33] or redundant [34] internal coordinates were added. All standard force constant types (stretches, nondegenerate bends, torsions, and degenerate bends) are available. Curvilinear coordinates are recommended for all calculations based on reaction paths.
- 3. The number-of-states calculation that is required for microcanonical variational theory  $(\mu VT)$  [35] is now programmed much more efficiently by the extended Beyer-Swinehart method [36,37]. Now there is no significant difference between CVT and  $\mu VT$  as far as computational cost.
- 4. The improved canonical variational theory (ICVT) option is removed. Whenever one needs a more accurate treatment of the threshold, one can simply use  $\mu$ VT without significantly increasing the computational cost, so there is no longer any motivation for using ICVT.
- 5. The first-order interpolated variational transition state theory (IVTST-1) method [38], interpolated variational transition state theory by mapping (IVTST-M) [39,40], including a spline-under-tension option, and several variational-transition-state-theory-with-interpolated-corrections (VTST-IC) [40] algorithms were added. IVTST-M allows arbitrary numbers of points along the reaction path and arbitrary numbers of points at which Hessians are computed, and it leads to faster convergence of direct dynamics calculations. VTST-IC methods include variational transition state theory with interpolated single point energies (VTST-ISPE).
- 6. Interfaces for direct dynamics were made more convenient. The PTUMME keyword to print an interface file readable by the TUMME 2023 program [23-25] has been added.
- 7. An option for performing calculations with classical rather than quantal partition functions was added.
- 8. The code has been modified to calculate wells in the reactant and/or product channels. One can optimize the reactants, the products, and the saddle point in separate runs. Quasi-Newton optimization with BFGS updating [41–44] was added. The eigenvector following

- (EF) algorithm [45–47] was added for geometry optimization of transition states and other stationary states.
- 9. The canonical unified statistical model (CUS) [6,48–50] was added.
- 10. The re-orientation of the dividing surface (RODS) [51,52] and variational reaction path (VRP) [53] algorithms were implemented. These methods allow VTST calculations with larger step sizes.
- 11. An option was added to perform nonequilibrium solvation calculations [54].
- 12. Version 4 of the large-curvature ground state tunneling method (LCG4) [55] was added as the default large-curvature tunneling (LCT) method, and it is now also used in the optimized multidimensional tunneling (OMT) [56] and microcanonical optimized multidimensional (μOMT) [20] tunneling methods. The interpolated algorithm for efficient large-curvature tunneling calculations (ILCT) [6,35] was also added.
- 13. New WKB vibrational anharmonicity [58,59] capability was added.
- 14. An option was added to scale the vibrational frequencies [60,61].
- 15. A keyword was added to allow users to raise low frequencies to a certain number (e.g., 50 or 100 cm<sup>-1</sup>) to approximately account for anharmonicity [62,63].
- 16. Several separable options for including anharmonicity in hindered internal rotations were added [64].
- 17. The single-structure MS-T method [12,14] was added with coupled torsional potentials [16]. This capability is added both for saddle point and for including torsional anharmonicity along a reaction path. Multi-structural [13] and multi-path [65–67] variational transition state theory calculations can be performed by combining the data from multiple *Polyrate 2023* and *MSTor* outputs.
- 18. The SS-QRRK code for pressure-dependent rate constants [22,68,69] has been included as a utility code. This method can include variational effects, multi-dimensional tunneling, and various vibrational and torsional anharmonicity options into the computed pressure-dependent rate constants and the SS-QRRK microcanonical rate constants. The FE integral for the normalized Boltzmann-weighted number of unimolecular states above the threshold energy can be evaluated via numerical integration instead of using the approximated analytical formula [70].
- 19. The standard-state free energy of activation profile is calculated along a reaction path using a standard state of 1 bar rather than an unconventional pressure.
- 20. The generalized free energy of activation can now be calculated using logarithms of the vibrational partition functions to avoid underflow. Similarly, a shifting constant is used to solve the overflow and underflow problem in the case that the imaginary action integral (the semiclassical tunneling integral) is too large.
- 21. The code was rearranged to allow parallel processing using MPI.

- 22. All common blocks were replaced by Fortran 90 modules. Memory for arrays is dynamically allocated. Therefore, compiling the code with a predefined system size (e.g., number of atoms) is no longer required.
- 23. Various bug fixes have been made, and some errors have been corrected. The restart option was improved. The code was made more portable, many options were made more convenient, defaults or keywords were improved, the output and manual were improved, and many new test runs were added.

#### References

- [1] A. D. Isaacson, D. G. Truhlar, S. N. Rai, R. Steckler, G. C. Hancock, B. C. Garrett, and M. J. Redmon, POLYRATE: A General Computer Program for Variational Transition State Theory and Semiclassical Tunneling Calculations of Chemical Reaction Rates, Comput. Phys. Commun. 47 (1987) 91–102. doi.org/10.1016/0010-4655(87)90069-5
- [2] D.-h. Lu, T. N. Truong, V. S. Melissas, G. C. Lynch, Y.-P. Liu, B. C. Garrett, R. Steckler, A. D. Isaacson, S. N. Rai, G. C. Hancock, J. G. Lauderdale, T. Joseph, and D. G. Truhlar, POLYRATE 4: A New Version of a Computer Program for the Calculation of Chemical Reaction Rates for Polyatomics, Comput. Phys. Commun. 71 (1992) 235-262. doi.org/10.1016/0010-4655(92)90012-N
- [3] R. Steckler, W.-P. Hu, Y.-P. Liu, G. C. Lynch, B. C. Garrett, A. D. Isaacson, V. S. Melissas, D.-H. Lu, T. N. Truong, S. N. Rai, G. C. Hancock, J. G. Lauderdale, T. Joseph, and D. G. Truhlar, POLYRATE 6.5: A New Version of Computer Program for the Calculation of Chemical Reaction Rates for Polyatomics, Comput. Phys. Commun. 88 (1995) 341-343 doi.org/10.1016/0010-4655(95)00037-G
- [4] D. G. Truhlar and B. C. Garrett, Variational Transition-State Theory, Acc. Chem. Res. 13 (1980) 440-448. doi.org/10.1021/ar50156a002
- [5] D. G. Truhlar and B. C. Garrett, Variational Transition State Theory, Annu. Rev. Phys. Chem. 35, (1984) 159-189. doi.org/10.1146/annurev.pc.35.100184.001111
- [6] D. G. Truhlar, A. D. Isaacson, and B. C. Garrett, Generalized Transition State Theory, in: Theory of Chemical Reaction Dynamics, edited by M. Baer, CRC Press, Boca Raton, FL, 1985, Vol. 4, pp. 65-137.
- [7] A. Fernandez-Ramos, B. A. Ellingson, B. C. Garrett, and D. G. Truhlar, Variational Transition State Theory with Multidimensional Tunneling, Rev. Comput. Chem. 23 (2007) 125-232. doi.org/10.1002/9780470116449.ch3
- [8] J. L. Bao and D. G. Truhlar, Variational Transition State Theory: Theoretical Framework and Recent Developments, Chem. Soc. Rev. 46 (2017) 7548-7596. doi.org/10.1039/C7CS00602K
- [9] S. Glasstone, K. J. Laidler, and H. Eyring, The Theory of Rate Processes, McGraw-Hill Book Co., New York, 1941.

- [10] D. G. Truhlar and A. D. Isaacson, Statistical-Diabatic Model for State-Selected Reaction Rates. Theory and Application of Vibrational-Mode Correlation Analysis to  $OH(n_{OH}) + H2(n_{HH}) \rightarrow H_2O + H$ , J. Chem. Phys. 77 (1982) 3516-3522. doi.org/10.1063/1.444297
- [11] R. Steckler, D. G. Truhlar, and B. C. Garrett, Tests of the Extension of Variational Transition State Theory to Calculate Reaction Rates for Molecules in Selected Excited Vibrational States, J. Chem. Phys. 84 (1986) 6712-6718. doi.org/10.1063/1.450673
- [12] J. Zheng, T. Yu, E. Papajak, I. M. Alecu, S. L. Mielke, and D. G. Truhlar, Practical Methods for Including Torsional Anharmonicity in Thermochemical Calculations of Complex Molecules: The Internal-Coordinate Multi-Structural Approximation, Phys. Chem. Phys. 13 (2011) 10885-10907. doi.org/10.1039/C0CP02644A
- [13] T. Yu, J. Zheng, and D. G. Truhlar, Multi-Structural Variational Transition State Theory. Kinetics of the 1,4-Hydrogen Shift Isomerization of the Pentyl Radical with Torsional Anharmonicity, Chem. Sci. 2 (2011) 2199-2213. doi.org/10.1039/C1SC00225B
- [14] J. Zheng, S. L Mielke, K. L Clarkson, and D. G. Truhlar, MSTor: A Program for Calculating Partition Functions, Free Energies, Enthalpies, Entropies, and Heat Capacities of Complex Molecules Including Torsional Anharmonicity, Comput. Phys. Commun. 183 (2012) 1803-1812. doi.org/10.1016/j.cpc.2012.03.007
- [15] W. Chen, J. Zheng, J. L. Bao, D. G. Truhlar, and X. Xu, MSTor 2023: A New Version of the Computer Code for Multistructural Torsional Anharmonicity, Now with Automatic Torsional Identification using Redundant Internal Coordinates, Comput. Phys. Commun. 288 (2023) 108740. doi.org/10.1016/j.cpc.2023.108740
- [16] J. Zheng and D. G. Truhlar, Quantum Thermochemistry: Multi-Structural Method with Torsional Anharmonicity Based on a Coupled Torsional Potential, J. Chem. Theory Comput. 9 (2013) 1356-1367. doi.org/10.1021/ct3010722
- [17] Y.-P. Liu, G. C. Lynch, T. N. Truong, D.-h. Lu, D. G. Truhlar, and B. C. Garrett, Molecular Modeling of the Kinetic Isotope Effect for the [1,5]-Sigmatropic Rearrangement of *cis*-1,3-Pentadiene, J. Am. Chem. Soc. 115 (1993) 2408-2415. doi.org/10.1021/ja00059a041
- [18] A. Fernandez-Ramos and D. G. Truhlar, Improved Algorithm for Corner Cutting Calculations, J. Chem. Phys. 114 (2001) 1491-1496. doi.org/10.1063/1.1329893
- [19] A. Fernández-Ramos and D. G. Truhlar, A New Algorithm for Efficient Direct Dynamics Calculations of Large-Curvature Tunneling and Its Application to Radical Reactions with 9–15 Atoms, J. Chem. Theory Comput. 1 (2005) 1063-1078. doi.org/10.1021/ct050153i
- [20] Y.-P. Liu, D.-h. Lu, A. Gonzàlez-Lafont, D. G. Truhlar, and B. C. Garrett, Direct Dynamics Calculation of the Kinetic Isotope Effect for an Organic Hydrogen-Transfer Reaction, Including Corner-Cutting Tunneling in 21 Dimensions, J. Am. Chem. Soc. 115 (1993) 7806-7817. doi.org/10.1021/ja00070a029
- [21] A. Fernández-Ramos, J. A. Miller, S. J. Klippenstein, and D. G. Truhlar, Modeling the Kinetics of Bimolecular Reactions, Chem. Rev. 106 (2006) 4518-4584. doi.org/10.1021/cr050205w

- [22] J. L. Bao, J. Zheng, and D. G. Truhlar, Kinetics of Hydrogen Radical Reactions with Toluene by Chemical Activation Theory Employing System-Specific Quantum RRK Theory Calibrated by Variational Transition State Theory, J. Am. Chem. Soc. 138 (2016) 2690-2704. doi.org/10.1021/jacs.5b11938
- [23] R. M. Zhang, X. Xu, and Donald G. Truhlar, TUMME 2023: An Open-Source Master Equation Program for Chemical Kinetics. Zenodo. https://doi.org/10.5281/zenodo.7943284
- [24] R. M. Zhang, X. Xu, and Donald G. Truhlar, TUMME 2023: Tsinghua University Minnesota Master Equation Program. New Version Announcement, Comput. Phys. Commun., submitted.
- [25] https://truhlar.chem.umn.edu/computational-tools/software
- [26] Y. Georgievskii and S. J. Klippenstein, Variable Reaction Coordinate Transition State Theory: Analytic Results and Application to the C<sub>2</sub>H<sub>3</sub> + H → C<sub>2</sub>H<sub>4</sub> Reaction, J. Chem. Phys. 118 (2003) 5442-5455. doi.org/10.1063/1.1539035
- [27] Y. Georgievskii and S. J. Klippenstein, Transition State Theory for Multichannel Addition Reactions: Multifaceted Dividing Surfaces, J. Phys. Chem. A 107 (2003) 9776-9781. doi.org/10.1021/jp034564b
- [28] J. Zheng, S. Zhang, and D. G. Truhlar, Density Functional Study of Methyl Radical Association Reaction Kinetics, J. Phys. Chem. A 112 (2008) 11509-11513. doi.org/10.1021/jp806617m
- [29] J. L. Bao, X. Zhang, and D. G. Truhlar, Association of CF<sub>2</sub> and Dissociation of C<sub>2</sub>F<sub>4</sub> by Variational Transition State Theory and System-Specific Quantum RRK Theory, Proc. Nat. Acad. Sci. U.S.A. 113 (2016) 13606-13611. doi.org/10.1073/pnas.1616208113
- [30] L. Zhang, D. G. Truhlar, and S. Sun, Association of Cl with C<sub>2</sub>H<sub>2</sub> by Unified Variable-Reaction-Coordinate and Reaction-Path Variational Transition State Theory, Proc. Nat. Acad. Sci. U.S.A. 117 (2020) 5610-5616. doi.org/10.1073/pnas.1920018117
- [31] C. F. Jackels, Z. Gu, and D. G. Truhlar, Reaction-Path Potential and Vibrational Frequencies in Terms of Curvilinear Internal Coordinates, J. Chem. Phys. 102 (1995) 3188-3201. doi.org/10.1063/1.468630
- [32] K. A. Nguyen, C. F. Jackels, and D. G. Truhlar, Reaction-Path Dynamics in Curvilinear Internal Coordinates Including Torsions, J. Chem. Phys. 104 (1996) 6491-6496. doi.org/10.1063/1.471369
- [33] Y.-Y. Chuang and D. G. Truhlar, Reaction-Path Dynamics with Harmonic Vibration Frequencies in Curvilinear Internal Coordinates: H + trans-N<sub>2</sub>H<sub>2</sub> → NH<sub>2</sub> + H<sub>2</sub>, J. Chem. Phys. 107 (1997) 83-89. doi.org/10.1063/1.474377
- [34] Y.-Y. Chuang and D. G. Truhlar, Reaction-Path Dynamics in Redundant Internal Coordinates, J. Phys. Chem. A 102 (1998) 242-247. doi.org/10.1021/jp9724028
- [35] A. D. Isaacson, M. T. Sund, S. N. Rai, and D. G. Truhlar, Improved Canonical and Microcanonical Variational Transition State Theory Calculations for a Polyatomic System: OH + H<sub>2</sub> → H<sub>2</sub>O + H, J. Chem. Phys. 82 (1985) 1338-1340. doi.org/10.1063/1.448963

- [36] T. Beyer and D. F. Swinehart, Algorithm 448: Number of Multiply-Restricted Partitions, Commun. ACM 16 (1973) 379. doi.org/10.1145/362248.362275
- [37] J. L. Bao, J. Zheng, and D. G. Truhlar, Kinetics of Hydrogen Radical Reactions with Toluene by Chemical Activation Theory Employing System-Specific Quantum RRK Theory Calibrated by Variational Transition State Theory, J. Am. Chem. Soc. 138 (2016) 2690-2704. doi.org/10.1021/jacs.5b11938
- [38] A. Gonzàlez-Lafont, T. N. Truong, and D. G. Truhlar, Interpolated Variational Transition State Theory: Practical Methods for Estimating Variational Transition State Properties and Tunneling Contributions to Chemical Reaction Rates from Electronic Structure Calculations, J. Chem. Phys. 95 (1991) 8875-8894. doi.org/10.1063/1.461221
- [39] J. C. Corchado () E. L. Coitiño, Y.-Y. Chuang, P. L. Fast, and D. G. Truhlar, Interpolated Variational Transition State Theory by Mapping, J. Phys. Chem. A 102 (1998) 2424-2438. doi.org/10.1021/jp9801267
- [40] Y.-Y. Chuang, J. C. Corchado, and D. G. Truhlar, Mapped Interpolation Scheme for Single-Point Energy Corrections in Reaction Rate Calculations and a Critical Evaluation of Dual-Level Reaction-Path Dynamics Methods, J. Phys. Chem. A 103 (1999) 1140-1149. doi.org/10.1021/jp9842493
- [41] C. J. Broyden, The Convergence of a Class of Double-Rank Minimization Algorithms, Inst. Math. App. 6 (1970) 76–90. doi.org/10.1093/imamat/6.1.76
- [42] R. Fletcher, New Approach to Variable Metric Algorithms, Comput J. (UK) 13 (1970) 317–322. doi.org/10.1093/comjnl/13.3.317
- [43] D. Goldfarb, A Family of Variable-Metric Methods Derived by Variational Means, Math. Comput. 24 (1970) 23–26. doi.org/10.1090/S0025-5718-1970-0258249-6
- [44] D. F. Shannon, Conditioning of Quasi-Newton Methods for Function Minimization, Math. Comput. 24 (1970) 647–656. doi.org/ 10.1090/S0025-5718-1970-0274029-X
- [45] A. Banerjee, N. Adams, J. Simons, and R. Shepard, Search for Stationary Points on Surfaces, J. Phys. Chem. 89 (1985) 52-57. doi.org/10.1021/j100247a015
- [46] J. Baker, An Algorithm for the Location of Transition States, J. Comput. Chem. 7 (1986) 385-395. doi.org/10.1002/jcc.540070402
- [47] F. Jensen, Locating Transition Structures by Mode Following: A Comparison of Six Methods on the Ar<sub>8</sub> Lennard-Jones Potential. J. Chem. Phys. 102 (1995) 6706-6718. doi.org/10.1063/1.469144
- [48] B. C. Garrett and D. G. Truhlar, Canonical Unified Statistical Model. Classical Mechanical Theory and Applications to Collinear Reactions, J. Chem. Phys. 76 (1982) 1853-1858. doi.org/10.1063/1.443157
- [49] X. G. Zhao, D.-h. Lu, Y.-P. Liu, G. C. Lynch, and D. G. Truhlar, Use of an Improved Ion-Solvent Potential Energy Function to Calculate the Reaction Rate and α-Deuterium and Microsolvation Kinetic Isotope Effects for the Gas-Phase S<sub>N</sub>2 Reaction of Cl<sup>-</sup>(H<sub>2</sub>O) with CH<sub>3</sub>Cl, J. Chem. Phys. 97 (1992) 6369-6383. doi.org/10.1063/1.463698

- [50] W.-P. Hu and D. G. Truhlar, Deuterium Kinetic Isotope Effects and their Temperature Dependence in the Gas-Phase  $S_N2$  Reactions  $X^- + CH_3Y \rightarrow CH_3X + Y^-$  (X, Y = Cl, Br, I), J. Am. Chem. Soc. 117 (1995) 10726-10734. doi.org/10.1021/ja00148a015
- [51] J. Villà and D. G. Truhlar, Variational Transition State Theory Without the Minimum Energy Path, Theor. Chem. Acc. 97 (1997) 317-323. doi.org/10.1007/s002140050267
- [52] A. González-Lafont, J. Villà, J. M. Lluch, J. Bertrán, R. Steckler, and D. G. Truhlar, Variational Transition State Theory and Tunneling Calculations with Reorientation of the Generalized Transition States for Methyl Cation Transfer, J. Phys. Chem. A 102 (1998) 3420-3428. doi.org/10.1021/jp9807672
- [53] P. L. Fast and D. G. Truhlar, Variational Reaction Path Algorithm, J. Chem. Phys. 109 (1998) 3721-3729. doi.org/10.1063/1.476973
- [54] Y.-Y. Chuang and D. G. Truhlar, Nonequilibrium Solvation Effects for a Polyatomic Reaction in Solution, J. Am. Chem. Soc. 121 (1999) 10157-10167. doi.org/10.1021/ja991809i
- [55] A. Fernandez-Ramos and D. G. Truhlar, Improved Algorithm for Corner Cutting Calculations, J. Chem. Phys. 114 (2001) 1491-1496. doi.org/10.1063/1.1329893
- [56] D. G. Truhlar, D.-h. Lu, S. C. Tucker, X. G. Zhao, A. Gonzàlez-Lafont, T. N. Truong, D. Maurice, Y-.P. Liu, and G. C. Lynch, Variational Transition State Theory with Multidimensional Semiclassical Ground-State Transmission Coefficients: Applications to Secondary Deuterium Kinetic Isotope Effects in Reactions Involving Methane and Chloromethane, Amer. Chem. Soc. Symp. Ser. 502 (1992) 16-36. doi.org/10.1021/bk-1992-0502.ch002
- [57] A. Fernandez-Ramos, D. G. Truhlar, J. C. Corchado, and J. Espinosa-Garcia, Interpolated Algorithms for Large-Curvature Tunneling Calculations, J. Phys. Chem. A 106 (2002) 4957-4960. doi.org/10.1021/jp014204t
- [58] B. C. Garrett and D. G. Truhlar, WKB Approximation for the Reaction-Path Hamiltonian: Application to Variational Transition State Theory, Vibrationally Adiabatic Excited-State Barrier Heights, and Resonance Calculations, J. Chem. Phys. 81 (2001) 309-317 (x). doi.org/10.1063/1.447386
- [59] B. C. Garrett and D. G. Truhlar, Thermal and State-Selected Rate Coefficients for the O(<sup>3</sup>P) + HCl Reaction and New Calculations of the Barrier Height and Width, S. Skokov, S. Zou, J. M. Bowman, T. C. Allison, D. G. Truhlar, Y. Lin, B. Ramachandran, B. C. Garrett, and B. J. Lynch, J. Phys. Chem. A 105 (2001) 2298-2307. doi.org/10.1021/jp003783j
- [60] J. Pu and D. G. Truhlar, Parameterized Direct Dynamics Study of Rate Constants of H with CH<sub>4</sub> from 250 to 2400 K, J. Pu and D. G. Truhlar, J. Chem. Phys. 116 (2002) 1468-1478. doi.org/10.1063/1.1427917
- [61] H. S. Yu, L. J. Fiedler, I. M. Alecu, and D. G. Truhlar, Computational Thermochemistry: Automated Generation of Scale Factors for Vibrational Frequencies Calculated by Electronic Structure Model Chemistries, Comput. Phys. Commun. 210 (2017) 132-138.

- doi.org/10.1016/j.cpc.2016.09.004
- [62] L. M. Pratt, D. G. Truhlar, C. J. Cramer, S. R. Kass, J. D. Thompson, and J. D. Xidos, Aggregation of Alkyllithiums in Tetrahydrofuran, J. Org. Chem. 72 (2007) 2962-2966. doi.org/10.1021/jo0625570
- [63] A. Marenich, W. Ding, C. J. Cramer, and D. G. Truhlar, Resolution of a Challenge for Solvation Modeling: Calculation of Dicarboxylic Acid Dissociation Constants Using Mixed Discrete – Continuum Solvation Models, J. Phys. Chem. Lett. 3 (2012) 1437-1442. doi.org/10.1021/jz300416r
- [64] Y.-Y. Chuang and D. G. Truhlar, Statistical Thermodynamics of Bond Torsion Modes, J. Chem. Phys. 124 (2006) 179903. doi.org/10.1063/1.2190219
- [65] T. Yu, J. Zheng, and D. G. Truhlar, Multipath Variational Transition State Theory. Rate Constant of the 1,4-Hydrogen Shift Isomerization of the 2-Cyclohexylethyl Radical, J. Phys. Chem. A 116 (2012) 297-308. doi.org/10.1021/jp209146b
- [66] J. Zheng and D. G. Truhlar, Multi-path Variational Transition State Theory for Chemical Reaction Rates of Complex Polyatomic Species: Ethanol + OH Reactions, Faraday Discuss. 157 (2012) 59-88. doi.org/10.1039/C2FD20012K.
- [67] J. L. Bao, R. Meana-Pañeda, and D. G. Truhlar, Multi-Path Variational Transition State Theory for Chiral Molecules: The Site-Dependent Kinetics for Abstraction of Hydrogen from Hydroperoxyl Radical, Analysis of Hydrogen Bonding in the Transition State, and Dramatic Temperature Dependence of the Activation Energy, Chem. Sci. 6 (2015) 5866-5881. doi.org/10.1039/C5SC01848J
- [68] J. L. Bao and D. G. Truhlar, Silane-Initiated Nucleation in Chemically Active Plasmas: Validation of Density Functionals, Mechanisms, and Pressure-Dependent Variational Transition State Calculations, Phys. Chem. Chem. Phys. 18 (2016) 10097-10108. doi.org/10.1039/c6cp00816j
- [69] J. L. Bao, X. Zhang, and D. G. Truhlar, Predicting Pressure-Dependent Unimolecular Rate Constants Using Variational Transition State Theory with Multidimensional Tunneling Combined with System-Specific Quantum RRK Theory: A Definitive Test for Fluoroform Dissociation, Phys. Chem. Chem. Phys. 18 (2016) 16659-16670. doi.org/10.1039/C6CP02765B
- [70] J. L. Bao and D. G. Truhlar, Effect of Energy Dependence of the Density of States on Pressure-Dependent Rate Constants, Phys. Chem. Chem. Phys. 20 (2018) 30475-30479. doi.org/10.1039/C8CP05915B

### **Declaration of competing interest**

The authors declare that they have no competing financial interests or personal relationships that could influence or appear to influence the work reported in this paper.

### Acknowledgments

This work was supported in part by the National Natural Science Foundation of China (award 21973053) and by the U.S. Department of Energy, Office of Science, Office of Basic Energy Sciences under Award DE-SC0015997.

### AUTHOR INFORMATION

#### Corresponding Authors

\*E-mail: truhlar@umn.edu (DGT),xuxuefei@tsinghua.edu.cn (XX)

### **ORCID**

Rui Ming Zhang: 0000-0002-4880-6391 Donald G. Truhlar: 0000-0002-7742-7294

Xuefei Xu: 0000-0002-2009-0483

### **Declaration of competing interest**

The authors declare that they have no competing financial interests or personal relationships that could influence or appear to influence the work reported in this paper.