

Advanced Engineering Mathematics Complex Analysis by Dennis G. Zill Notes

Chris Doble

February 2024

Contents

17 Functions of a Complex Variable	1
17.1 Complex Numbers	1
17.2 Powers and Roots	3

17 Functions of a Complex Variable

17.1 Complex Numbers

- A **complex number** is any number of the form

$$z = a + ib$$

where a and b are real numbers and i is the imaginary unit such that $i^2 = -1$.

- The real number a in the above complex number z is called the **real part** of z and the real number b (not ib) is called the **imaginary part** of z .
- The real and imaginary parts of a complex number z are denoted $\text{Re}(z)$ and $\text{Im}(z)$, respectively.
- A real constant multiple of the imaginary unit, e.g. $6i$ is called a **pure imaginary number**.
- Two complex numbers are equal if their real and imaginary parts are equal.
- The addition and subtraction of complex numbers occur between the real and imaginary parts, e.g.

$$(a + bi) + (c + di) = (a + c) + (b + d)i.$$

- The multiplication of complex numbers occurs elementwise as normal, e.g.

$$(a + bi)(c + di) = ac + adi + bci - bd.$$

- The **conjugate** of a complex number $z = a + ib$ is

$$\bar{z} = a - ib.$$

- The division of complex numbers occurs by multiplying the numerator and denominator by the conjugate of the denominator, e.g.

$$\begin{aligned}\frac{a + bi}{c + di} &= \frac{(a + bi)(c - di)}{(c + di)(c - di)} \\ &= \frac{ac - adi + bci + bd}{c^2 + d^2} \\ &= \frac{ac + bd}{c^2 + d^2} + i \frac{bc - ad}{c^2 + d^2}.\end{aligned}$$

- Conjugates have several interesting properties:

$$\begin{aligned}\overline{z_1 + z_2} &= \bar{z}_1 + \bar{z}_2 \\ \overline{z_1 - z_2} &= \bar{z}_1 - \bar{z}_2 \\ \overline{z_1 z_2} &= \bar{z}_1 \bar{z}_2 \\ \frac{z_1}{z_2} &= \frac{\bar{z}_1}{\bar{z}_2}.\end{aligned}$$

- The sum and product of a complex number $z = x + iy$ with its conjugate are real numbers

$$\begin{aligned}z + \bar{z} &= 2x \\ z\bar{z} &= x^2 + y^2\end{aligned}$$

while the difference between a complex number and its conjugate is a pure imaginary number

$$z - \bar{z} = 2iy.$$

- The above properties let us define

$$\operatorname{Re}(z) = \frac{z + \bar{z}}{2} \text{ and } \operatorname{Im}(z) = \frac{z - \bar{z}}{2i}.$$

- The **complex plane** or **z -plane** is a coordinate system where the horizontal or x -axis is called the **real axis** and the vertical or y -axis is called the **imaginary axis**. Complex numbers can be plotted in this coordinate system by considering their real and imaginary parts an ordered pair corresponding their position.

- The **modulus** or **absolute value** of a complex number $z = x + iy$ denoted by $|z|$ is the real number

$$|z| = \sqrt{x^2 + y^2} = \sqrt{z\bar{z}}.$$

This is the distance between z and the origin in the complex plane.

- If you consider two numbers in the complex plane as vectors, the length of their sum can't be longer than their individual lengths combined

$$|z_1 + z_2| \leq |z_1| + |z_2|.$$

This extends to any finite sum

$$|z_1 + z_2 + \cdots + z_n| \leq |z_1| + |z_2| + \cdots + |z_n|$$

and is known as the **triangle inequality**.

17.2 Powers and Roots

- A complex number can be expressed in **polar form**

$$z = (r \cos \theta) + i(r \sin \theta)$$

where $r = |z|$ is the nonnegative modulus of z and $\theta = \arg z$ is the **argument** of z — the angle between z and the positive real axis measured in the counterclockwise direction.

- The argument of a complex number z isn't unique as any multiply of 2π can be added to it. The **principle argument** of z denoted $\text{Arg } z$ is the argument of z restricted to the interval $-\pi \leq \text{Arg } z \leq \pi$.
- Multiplication and division of complex numbers is simpler in polar form. For two complex numbers $z_1 = r_1(\cos \theta_1 + i \sin \theta_1)$ and $z_2 = r_2(\cos \theta_2 + i \sin \theta_2)$ we get

$$\begin{aligned} z_1 z_2 &= r_1 r_2 [\cos(\theta_1 + \theta_2) + i \sin(\theta_1 + \theta_2)] \\ \frac{z_1}{z_2} &= \frac{r_1}{r_2} [\cos(\theta_1 - \theta_2) + i \sin(\theta_1 - \theta_2)]. \end{aligned}$$

- The above formulas can be used to find integer powers of a complex number z

$$z^n = r^n (\cos n\theta + i \sin n\theta)$$

where n is an integer (including negative integers).

- **DeMoivre's formula** is a special case of the above where $r = 1$ so

$$z^n = (\cos \theta + i \sin \theta)^n = \cos n\theta + i \sin n\theta.$$

- A number w is said to be an **n th root** of a nonzero complex number z if $w^n = z$. The n th roots of a nonzero complex number $z = r(\cos \theta + i \sin \theta)$ are

$$w_k = r^{1/n} \left[\cos \left(\frac{\theta + 2k\pi}{n} \right) + i \sin \left(\frac{\theta + 2k\pi}{n} \right) \right]$$

where $k = 0, 1, 2, \dots, n-1$.

- The root w of a complex number z obtained by using the principle argument of z with $k = 0$ is called the **principle n th root** of z .
- Since the n th roots of a complex number have the same modulus they lie on a circle of radius $r^{1/n}$. The arguments of subsequent roots differ by $2\pi/n$ so they're also equally spaced around the circle.