

Advanced Engineering Mathematics Partial
Differential Equations by Dennis G. Zill Notes

Chris Doble

November 2023

Contents

12 Orthogonal Functions and Fourier Series 1
12.1 Orthogonal Functions 1

12 Orthogonal Functions and Fourier Series

12.1 Orthogonal Functions

- The **inner product** of two functions f_1 and f_2 on an interval $[a, b]$ is the number

$$(f_1, f_2) = \int_a^b f_1(x)f_2(x) \, dx.$$

- Two functions f_1 and f_2 are said to be orthogonal on an interval if $(f_1, f_2) = 0$.
- A set of real-valued functions $\{\phi_1(x), \phi_2(x), \dots, \phi_n(x)\}$ is said to be **orthogonal** on an interval if

$$(\phi_i, \phi_j) = 0 \text{ for } i \neq j.$$

- The **square norm** of a function is

$$||\phi_n(x)||^2 = (\phi_n, \phi_n)$$

and thus its **norm** is

$$||\phi_n(x)|| = \sqrt{(\phi_n, \phi_n)}.$$

- An **orthonormal set** of functions is an orthogonal set of functions that all have a norm of 1.
- An orthogonal set can be made into an orthonormal set by dividing each member by its norm.

- If $\{\phi_n(x)\}$ is an infinite orthogonal set of functions on an interval $[a, b]$ and $f(x)$ is an arbitrary function, then it's possible to determine a set of coefficients $c_n, n = 0, 1, 2, \dots$ such that

$$f(x) = \sum_{n=0}^{\infty} c_n \phi_n(x) = c_0 \phi_0(x) + c_1 \phi_1(x) + \dots + c_n \phi_n(x) + \dots$$

This is called an **orthogonal series expansion** of f or a **generalized Fourier series** where the coefficients are given by

$$c_n = \frac{(f, \phi_n)}{||\phi_n||^2}.$$

- A set of real-valued functions $\{\phi_n(x)\}$ is said to be **orthogonal with respect to a weight function** $w(x)$ on the interval $[a, b]$ if

$$\int_a^b w(x) \phi_m(x) \phi_n(x) dx = 0, \quad m \neq n.$$