Quantum Computation and Quantum Information by Michael A. Nielsen and Isaac L. Chuang Problems

Chris Doble

June 2024

Contents

| I Fundamental concepts | 1 |
|--------------------------------|---|
| 2 Linear algebra | 1 |
| Part I Fundamental concepts | |
| 2 Linear algebra | |
| Exercise 2.1 | |
| (1,-1) + (1,2) - (2,1) = (0,0) | |

Exercise 2.2

$$A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$$

Using the basis $|+\rangle = (|0\rangle + |1\rangle)/\sqrt{2}$ and $|-\rangle = (|0\rangle - |1\rangle)/\sqrt{2}$ we get

$$|0\rangle = \frac{|+\rangle + |-\rangle}{\sqrt{2}}$$

$$|1\rangle = \frac{|+\rangle - |-\rangle}{\sqrt{2}}$$

$$\begin{bmatrix} a_{00} & a_{01} \\ a_{10} & a_{11} \end{bmatrix} \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

$$a_{00} + a_{01} = 1$$

$$a_{10} + a_{11} = -1$$

$$\begin{bmatrix} a_{00} & a_{01} \\ a_{10} & a_{11} \end{bmatrix} \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ -1 \end{bmatrix} = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

$$a_{00} - a_{01} = 1$$

$$a_{10} - a_{11} = 1$$

$$A = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$$