

Introduction to Quantum Mechanics by David J. Griffiths Notes

Chris Doble

March 2024

Contents

I	Theory	1
1	The Wave Function	1
1.1	The Schrödinger Equation	1
1.2	The Statistical Interpretation	2
1.3	Probability	2
1.3.1	Discrete Variables	2
1.3.2	Continuous Variables	4

Part I

Theory

1 The Wave Function

1.1 The Schrödinger Equation

- The Schrödinger equation

$$i\hbar \frac{\partial \Psi}{\partial t} = -\frac{\hbar^2}{2m} \frac{\partial^2 \Psi}{\partial x^2} + V\Psi$$

is to quantum mechanics what Newton's second law is to classical mechanics. Given suitable initial conditions — typically $\Psi(x, 0)$ — the Schrödinger equation determines $\Psi(x, t)$ for all future time.

1.2 The Statistical Interpretation

- The **Born rule** states that $|\Psi(x, t)|^2$ gives the probability of finding the particle at point x at time t or

$$\int_a^b |\Psi(x, t)|^2 dx$$

gives the probability of finding the particle between a and b at time t .

- This statistical interpretation introduces indeterminacy to quantum mechanics — we can't predict with certainty the particle's position.
- Suppose we measure a particle's position to be C . Where was it before we took the measurement? In the past there were three main schools of thought:
 1. The **realist** position believes that the particle was at C but Ψ doesn't give us enough information to determine that — there's another **hidden variable** that would allow us to.
 2. The **orthodox** position (also known as the **Copenhagen interpretation**) believes that the particle didn't have a definite position but the act of measuring it forced it to do so.
 3. The **agnostic** position believes that it doesn't matter and is potentially unknowable.
- **Bell's theorem** confirms the orthodox interpretation.
- If we take two consecutive measurements of a particle, they will both yield the same result. The first measurement causes the wavefunction to **collapse** such that it is peaked only at the particle's measured location. If the system is allowed to evolve between the measurements the wavefunction will "spread out" but if done in quick succession the result won't change.

1.3 Probability

1.3.1 Discrete Variables

- The average value of a discrete variable j is

$$\langle j \rangle = \frac{\sum jN(j)}{N} = \sum_{j=0}^{\infty} jP(j)$$

where N is the size of the population, $N(j)$ is the number of j s in the population, and $P(j)$ is the probability of randomly selecting a j from the population.

- In quantum mechanics the average is usually the quantity of interest and it is called the **expectation value** (even though it may not be the most probable value).
- The average value of a function f of a discrete variable j is

$$\langle f(j) \rangle = \sum_{j=0}^{\infty} f(j)P(j).$$

- Two distributions could have the same median, mean, mode, and size but be spread out differently. One way to quantify this could be to calculate how far each element is from the average

$$\Delta j = j - \langle j \rangle$$

and calculate the average of Δj . However, it ends up being zero

$$\begin{aligned} \langle \Delta j \rangle &= \sum (j - \langle j \rangle)P(j) \\ &= \sum jP(j) - \langle j \rangle \sum P(j) \\ &= \langle j \rangle - \langle j \rangle \\ &= 0. \end{aligned}$$

To avoid this we calculate the average of the square of Δj

$$\sigma^2 = \langle (\Delta j)^2 \rangle$$

which is known as the **variance** of the distribution.

- The square root of the variance is called the **standard deviation**.
- A useful theorem on variances is

$$\begin{aligned} \sigma^2 &= \langle (\Delta j)^2 \rangle \\ &= \sum (\Delta j)^2 P(j) \\ &= \sum (j - \langle j \rangle)^2 P(j) \\ &= \sum (j^2 - 2j\langle j \rangle + \langle j \rangle^2) P(j) \\ &= \sum j^2 P(j) - 2\langle j \rangle \sum jP(j) + \langle j \rangle^2 \sum P(j) \\ &= \langle j^2 \rangle - 2\langle j \rangle^2 + \langle j \rangle^2 \\ &= \langle j^2 \rangle - \langle j \rangle^2 \end{aligned}$$

and thus the standard deviation can be calculated as

$$\sigma = \sqrt{\langle j^2 \rangle - \langle j \rangle^2}$$

which is usually easier than the full formula.

1.3.2 Continuous Variables

- The equations above translate as expected to continuous variables:

$$P_{ab} = \int_a^b \rho(x) dx$$

$$1 = \int_{-\infty}^{+\infty} \rho(x) dx$$

$$\langle x \rangle = \int_{-\infty}^{+\infty} x \rho(x) dx$$

$$\langle f(x) \rangle = \int_{-\infty}^{+\infty} f(x) \rho(x) dx$$

$$\sigma^2 = \langle x^2 \rangle - \langle x \rangle^2$$