BECA / Dr. Huson / Geometry 06-Analytic-geometry Name: pset ID: 92


6-8HW-tangent-trig-ratios

- 1. Express the result to the nearest thousandth.
 - (a) $\tan 60^{\circ} =$

(c) $\tan 23^{\circ} =$

(b) $\tan 67^{\circ} =$


- (d) $\tan 45^{\circ} =$
- 2. Given right $\triangle JKL$ with $\overline{JK} \perp \overline{KL}$, JK = 7, $m \angle J = 20^{\circ}$.




(a) Use the tangent function to find the length KL

(b) Use the Pythagorean formula to find the length JL


3. Given right $\triangle ABC$ with $AC=6, BC=2.8, \ m\angle C=90^{\circ}$. Find the value of $m\angle A$, expressed as a decimal to the nearest thousandth.



4. Given right $\triangle JKL$ with $\overline{JK} \perp \overline{KL}$, JK = 7, $m \angle J = 20^{\circ}$. Find the length KL.



5. Given right $\triangle ABC$ with $AC=10, BC=4, \ m\angle C=90^{\circ}$. Find the value of $m\angle A$, expressed as a decimal to the nearest thousandth.



- 6. Spicy: Given a rectangle with area 35, width x, and length x+2.
 - (a) Find x.
 - (b) Find the perimeter of the rectangle.