2.28 Polynomials and Rational Functions: **End-of-Unit Assessment**

You may use a four-function or scientific calculator, but not a graphing calculator.

1. Select the expression that is equivalent to $\frac{2x^3+11x^2-21x}{x^2+3x}$ for $x \neq -3$ or 0.

A.
$$2x + 5 - \frac{6}{x+3}$$

B.
$$2x + 17 - \frac{20}{x+3}$$


C.
$$2x + 17 - \frac{36}{x^2 + 3x}$$

D.
$$2x + 5 - \frac{36}{x+3}$$

$$\frac{x^{2}+3x}{\chi^{2}+3x} \frac{2\chi+5}{2\chi^{3}+11\chi^{2}-21\chi}$$

$$2\chi^{3}+6\chi^{2}$$


F. Bf. 7 Solve problems using the Geometric Sin 2. Here are the first three stages of a sequence of dots. If we drew each of the first 15 $\frac{1}{2}$ stages, how many dots would we have to draw? Select all of the expressions representing this number.



C.
$$(1-3)(1+3+3^2+...+3^{14})$$


D) $3\frac{1-3^{15}}{1-3}$

E. $3(1-3)(1+3+3^2+...+3^{14})$




F. IF. 7d Graph rabing / Fundings (with #6)

3. Which figure below is a graph of $r(x) = 2 + \frac{x+1}{x-3}$?



A.



Vertical
asymptote


x=3

(hor. Fontal)
y=3

B.

-10 -8 -6 -6


C.



D.

A. REI, 4 Silve quadratic equations

4. Find all values of x that make the equation true.



$$\frac{3}{x-4} = \frac{x-5}{x}$$

$$3x = (x-4)(x-5)$$

$$= x^2 - 9x + 70$$

$$x^2 - 12x + 70 = 0$$

$$(x-7)(x-10) = 0$$

$$x=7$$

5. Kiran and Mai are trying to figure out if this equation is an identity:

Fig. 7 denite $|exp|_{x=a}^{x=a} (a-b)^4 = a^4 - 4a^3b + 6a^2b^2 - 4ab^3 + b^4$ Geometric Sum formula Kiran picks several different values for a and b to test. For each pair he tries, the left

side and the right side have the same value. Since his tests all worked, Kiran thinks the equation is an identity.

Mai started by using the distributive property on the expression on the left side. She then arranged the result to look the same as the right side. Since she can write both sides the same way, Mai thinks the equation is an identity.

Who do you think had a better argument for why the equation is an identity? Explain your reasoning.


Mais argument is better. It is general because it applies to all possible values of a RB.

Kiran's method may have missel Certain values which would Violate the equality. F. IF.7d (Rept patienal expressions

6. A plane is going to fly 300 miles at a planned speed of 530 mph. The flight will have an average headwind of w miles per hour the entire time, meaning the plane is flying directly against the wind. The time T, in hours, of the flight is a function of the speed of the headwind w, in mph, and can be modeled by

$$T(w) = \frac{300}{530 - w}$$

Here is the graph of y = T(w):



a. What does T(110) mean in this situation?

The time to fly 300 miles if the wind speed

b. At what value of w does the graph have a vertical asymptote? Explain how you know and what this asymptote means in the situation.

W= 530 mph (wind Speed)

Mg. bro, cells at that value the Denimotor

is zero and the finction unletimed. OR

with the plane's flight Speed equal

to the wind Speed, It would never

CC BY 2019 by Illustrative Mathematics®

AMNE

4




A. APR. 6 Rewrite ratinal expressions

7. There are many square prisms with volume 125 in^3 . Let wrepresent the side length of the square base and h represent the height in inches. Then the formula for the volume V, in cubic inches, of these prisms is $V = w^2 h$ and the equation for the surface area S, in square inches, is $S = 2w^2 + 4wh$.

$$V = \omega^2 h = 125$$

$$h = \frac{125}{\omega^2}$$



Write an equation for S as a function of w when the volume of the prism is 125 in 3 . Explain or show your work.

$$S(\omega) = 2\omega^2 + 4\omega \left(\frac{125}{\omega^2}\right)$$
$$= 2\omega^2 + \frac{500}{\omega}$$