

**5.12 Classwork: Trigonometric Transformations & Derivatives (no calculator)**

1. A Ferris wheel has a diameter of 80 metres. The lowest point of the wheel is 5 metres above the ground. The wheel completes one full revolution every 16 minutes. A seat starts at the lowest point at time  $t = 0$ .

The height,  $h$  metres, of the seat above the ground after  $t$  minutes is modelled by

$$h(t) = a \cos(bt) + d$$

where  $a, b, d \in \mathbb{R}$ .

- (a) Find the value of  $a$ ,  $b$ , and  $d$ . [4]

- (b) Find the height of the seat after 4 minutes. [2]

- (c) Find  $h'(t)$  and state what it represents in context. [2]

2. The graph of  $y = \sin x$  is transformed to give the graph of  $y = 3 \sin\left(2\left(x - \frac{\pi}{4}\right)\right) + 1$ .

(a) Describe the following transformations applied to  $y = \sin x$ :

(i) the vertical stretch, [1]

(ii) the horizontal compression, [1]

(iii) the horizontal translation, [1]


(iv) the vertical translation. [1]

(b) For the transformed function, write down:

(i) the amplitude, [1]

(ii) the period. [1]

(c) On the axes below, sketch the graph of  $y = 3 \sin\left(2\left(x - \frac{\pi}{4}\right)\right) + 1$  for  $0 \leq x \leq 2\pi$ .  
Clearly label any intercepts and maximum/minimum points. [4]



3. Find the derivative of each function.

(a)  $f(x) = 3 \sin x + 2 \cos x$  [2]

(b)  $g(x) = 4 \cos(3x)$  [2]

(c)  $h(x) = \sin^2 x$  [3]

*Hint: Write  $\sin^2 x = (\sin x)^2$  and use the chain rule.*

4. The function  $f$  is defined by  $f(x) = 2 \sin(3x) - 1$  for  $0 \leq x \leq \pi$ .

(a) Find  $f'(x)$ . [2]

(b) Find the  $x$ -coordinates of the points where  $f$  has a maximum and a minimum value. [3]

(c) Find the maximum and minimum values of  $f$ . [2]

5. The function  $g$  is defined by  $g(x) = a \sin(bx) + c$  for  $0 \leq x \leq 2\pi$ , where  $a, b, c \in \mathbb{Z}^+$ . The graph of  $g$  has a maximum value of 7, a minimum value of 1, and a period of  $\frac{2\pi}{3}$ .

(a) Find the values of  $a$ ,  $b$ , and  $c$ . [3]

(b) Find  $g'\left(\frac{\pi}{6}\right)$ . [3]

6. Consider the function  $f(x) = \cos 2x + 2 \sin x$  for  $0 \leq x \leq 2\pi$ .

(a) Show that  $f'(x) = 2 \cos x(1 - 2 \sin x)$ . [3]

(b) Hence find the values of  $x$  for which  $f'(x) = 0$ . [3]