

Lecture 2: predictive modeling @ NYT

desc/pred/pres


descriptive:	specify x ; learn $z(x)$ or $p(z x)$ where z is “simpler” than x
predictive:	specify x and y ; learn to predict y from x
prescriptive:	specify x , y , and a ; learn to prescribe a given x to maximize y

Figure 2: desc/pred/pres

- caveat: difference between observation and experiment. why?

blossom example

prescriptive modeling, e.g.,




leverage methods which are predictive yet performant

Figure 3: Reminder: Blossom

blossom + boosting ('exponential')

Margin-Based Surrogate Loss Functions



from "are you a bayesian or a frequentist"
-michael jordan

Figure 4: Reminder: Surrogate Loss Functions

tangent: logistic function as surrogate loss function

- ▶ define $f(x) \equiv \log p(y = 1|x)/p(y = -1|x) \in R$

tangent: logistic function as surrogate loss function

- ▶ define $f(x) \equiv \log p(y = 1|x)/p(y = -1|x) \in R$
- ▶ $p(y = 1|x) + p(y = -1|x) = 1 \rightarrow p(y|x) = 1/(1 + \exp(-yf))$

tangent: logistic function as surrogate loss function

- ▶ define $f(x) \equiv \log p(y = 1|x)/p(y = -1|x) \in R$
- ▶ $p(y = 1|x) + p(y = -1|x) = 1 \rightarrow p(y|x) = 1/(1 + \exp(-yf))$
- ▶ $-\log_2 p(\{y\}_1^N) = \sum_i \log_2 (1 + e^{-y_i f(x_i)}) \equiv \sum_i \ell(y_i f(x_i))$

tangent: logistic function as surrogate loss function

- ▶ define $f(x) \equiv \log p(y = 1|x)/p(y = -1|x) \in R$
- ▶ $p(y = 1|x) + p(y = -1|x) = 1 \rightarrow p(y|x) = 1/(1 + \exp(-yf))$
- ▶ $-\log_2 p(\{y\}_1^N) = \sum_i \log_2 (1 + e^{-y_i f(x_i)}) \equiv \sum_i \ell(y_i f(x_i))$
- ▶ $\ell'' > 0$, $\ell(\mu) > 1[\mu < 0] \forall \mu \in R$.

tangent: logistic function as surrogate loss function

- ▶ define $f(x) \equiv \log p(y = 1|x)/p(y = -1|x) \in R$
- ▶ $p(y = 1|x) + p(y = -1|x) = 1 \rightarrow p(y|x) = 1/(1 + \exp(-yf))$
- ▶ $-\log_2 p(\{y\}_1^N) = \sum_i \log_2 (1 + e^{-y_i f(x_i)}) \equiv \sum_i \ell(y_i f(x_i))$
- ▶ $\ell'' > 0$, $\ell(\mu) > 1[\mu < 0] \forall \mu \in R$.
- ▶ \therefore maximizing log-likelihood is minimizing a surrogate convex loss function for classification (though not strongly convex, cf. Yoram's talk)

tangent: logistic function as surrogate loss function

- ▶ define $f(x) \equiv \log p(y = 1|x)/p(y = -1|x) \in R$
- ▶ $p(y = 1|x) + p(y = -1|x) = 1 \rightarrow p(y|x) = 1/(1 + \exp(-yf))$
- ▶ $-\log_2 p(\{y\}_1^N) = \sum_i \log_2 (1 + e^{-y_i f(x_i)}) \equiv \sum_i \ell(y_i f(x_i))$
- ▶ $\ell'' > 0, \ell(\mu) > 1[\mu < 0] \forall \mu \in R$.
- ▶ \therefore maximizing log-likelihood is minimizing a surrogate convex loss function for classification (though not strongly convex, cf. Yoram's talk)
- ▶ but $\sum_i \log_2 (1 + e^{-y_i w^T h(x_i)})$ not as easy as $\sum_i e^{-y_i w^T h(x_i)}$

boosting 1

L exponential surrogate loss function, summed over examples:

- ▶ $L[F] = \sum_i \exp(-y_i F(x_i))$

boosting 1

L exponential surrogate loss function, summed over examples:

- ▶ $L[F] = \sum_i \exp(-y_i F(x_i))$
- ▶ $= \sum_i \exp(-y_i \sum_{t'}^t w_{t'} h_{t'}(x_i)) \equiv L_t(\mathbf{w}_t)$

boosting 1

L exponential surrogate loss function, summed over examples:

- ▶ $L[F] = \sum_i \exp(-y_i F(x_i))$
- ▶ $= \sum_i \exp(-y_i \sum_{t'}^t w_{t'} h_{t'}(x_i)) \equiv L_t(\mathbf{w}_t)$
- ▶ Draw $h_t \in \mathcal{H}$ large space of rules s.t. $h(x) \in \{-1, +1\}$

boosting 1

L exponential surrogate loss function, summed over examples:

- ▶ $L[F] = \sum_i \exp(-y_i F(x_i))$
- ▶ $= \sum_i \exp(-y_i \sum_{t'}^t w_{t'} h_{t'}(x_i)) \equiv L_t(\mathbf{w}_t)$
- ▶ Draw $h_t \in \mathcal{H}$ large space of rules s.t. $h(x) \in \{-1, +1\}$
- ▶ label $y \in \{-1, +1\}$

boosting 1

L exponential surrogate loss function, summed over examples:

$$\blacktriangleright L_{t+1}(\mathbf{w}_t; w) \equiv \sum_i d_i^t \exp(-y_i w h_{t+1}(x_i))$$

Punchlines: sparse, predictive, interpretable, fast (to execute), and easy to extend, e.g., trees, flexible hypotheses spaces, L_1, L_∞^1, \dots

boosting 1

L exponential surrogate loss function, summed over examples:

- ▶ $L_{t+1}(\mathbf{w}_t; w) \equiv \sum_i d_i^t \exp(-y_i w h_{t+1}(x_i))$
- ▶ $= \sum_{y=h'} d_i^t e^{-w} + \sum_{y \neq h'} d_i^t e^{+w} \equiv e^{-w} D_+ + e^{+w} D_-$

Punchlines: sparse, predictive, interpretable, fast (to execute), and easy to extend, e.g., trees, flexible hypotheses spaces, L_1, L_∞^1, \dots

boosting 1

L exponential surrogate loss function, summed over examples:

- ▶ $L_{t+1}(\mathbf{w}_t; w) \equiv \sum_i d_i^t \exp(-y_i w h_{t+1}(x_i))$
- ▶ $= \sum_{y=h'} d_i^t e^{-w} + \sum_{y \neq h'} d_i^t e^{+w} \equiv e^{-w} D_+ + e^{+w} D_-$
- ▶ $\therefore w_{t+1} = \operatorname{argmin}_w L_{t+1}(w) = (1/2) \log D_+ / D_-$

Punchlines: sparse, predictive, interpretable, fast (to execute), and easy to extend, e.g., trees, flexible hypotheses spaces, L_1, L_∞^1, \dots

boosting 1

L exponential surrogate loss function, summed over examples:

- ▶ $L_{t+1}(\mathbf{w}_t; w) \equiv \sum_i d_i^t \exp(-y_i w h_{t+1}(x_i))$
- ▶ $= \sum_{y=h'} d_i^t e^{-w} + \sum_{y \neq h'} d_i^t e^{+w} \equiv e^{-w} D_+ + e^{+w} D_-$
- ▶ $\therefore w_{t+1} = \operatorname{argmin}_w L_{t+1}(w) = (1/2) \log D_+ / D_-$
- ▶ $L_{t+1}(\mathbf{w}_{t+1}) = 2\sqrt{D_+ D_-} = 2\sqrt{\nu_+(1 - \nu_+)}/D$, where $0 \leq \nu_+ \equiv D_+/D = D_+/L_t \leq 1$

Punchlines: sparse, predictive, interpretable, fast (to execute), and easy to extend, e.g., trees, flexible hypotheses spaces, L_1, L_∞^1, \dots

boosting 1

L exponential surrogate loss function, summed over examples:

- ▶ $L_{t+1}(\mathbf{w}_t; w) \equiv \sum_i d_i^t \exp(-y_i w h_{t+1}(x_i))$
- ▶ $= \sum_{y=h'} d_i^t e^{-w} + \sum_{y \neq h'} d_i^t e^{+w} \equiv e^{-w} D_+ + e^{+w} D_-$
- ▶ $\therefore w_{t+1} = \operatorname{argmin}_w L_{t+1}(w) = (1/2) \log D_+/D_-$
- ▶ $L_{t+1}(\mathbf{w}_{t+1}) = 2\sqrt{D_+ D_-} = 2\sqrt{\nu_+(1-\nu_+)}/D$, where $0 \leq \nu_+ \equiv D_+/D = D_+/L_t \leq 1$
- ▶ update example weights $d_i^{t+1} = d_i^t e^{\mp w}$

Punchlines: sparse, predictive, interpretable, fast (to execute), and easy to extend, e.g., trees, flexible hypotheses spaces, L_1, L_∞^1, \dots

¹Duchi + Singer “Boosting with structural sparsity” ICML '09

predicting people

- ▶ “customer journey” prediction

predicting people

- ▶ “customer journey” prediction
 - ▶ fun covariates

predicting people

- ▶ “customer journey” prediction
 - ▶ fun covariates
 - ▶ observational complication v structural models

predicting people (reminder)








TFNAME	DB-MOTIF	MOTIF	DBNAME	$d(p,q)$
CBF1	CACGTG		YPD	0.032635
CGG everted repeat	CGGN*CCG		YPD	0.032821
MSN2			TRANSFAC	0.085626
HSF1	TTCNNNGAA		SCPD	0.102410
XBP1			TRANSFAC	0.140561

Figure 5: both in science and in real world, feature analysis guides future experiments

single copy (reminder)




Figure 6: from Lecture 1

example in CAR (computer assisted reporting)



Figure 7: Tabuchi article

example in CAR (computer assisted reporting)

- ▶ cf. Friedman's "Statistical models and Shoe Leather"²

²Freedman, David A. "Statistical models and shoe leather." Sociological methodology 21.2 (1991): 291-313.

example in CAR (computer assisted reporting)

- ▶ cf. Friedman's "Statistical models and Shoe Leather"²
- ▶ Takata airbag fatalities

²Freedman, David A. "Statistical models and shoe leather." Sociological methodology 21.2 (1991): 291-313.

example in CAR (computer assisted reporting)

- ▶ cf. Friedman's "Statistical models and Shoe Leather"²
- ▶ Takata airbag fatalities
- ▶ 2219 labeled³ examples from 33,204 comments

²Freedman, David A. "Statistical models and shoe leather." Sociological methodology 21.2 (1991): 291-313.

³By Hiroko Tabuchi, a Pulitzer winner

example in CAR (computer assisted reporting)

- ▶ cf. Friedman's "Statistical models and Shoe Leather"²
- ▶ Takata airbag fatalities
- ▶ 2219 labeled³ examples from 33,204 comments
- ▶ cf. Box's "Science and Statistics"⁴

²Freedman, David A. "Statistical models and shoe leather." Sociological methodology 21.2 (1991): 291-313.

³By Hiroko Tabuchi, a Pulitzer winner

⁴Science and Statistics, George E. P. Box Journal of the American Statistical Association, Vol. 71, No. 356. (Dec., 1976), pp. 791-799.

computer assisted reporting

► Impact




Figure 8: impact

Lecture 3: prescriptive modeling @ NYT

the natural abstraction

- ▶ operators⁵ make decisions

⁵In the sense of business deciders; that said, doctors, including those who operate, also have to make decisions, cf., personalized medicines

the natural abstraction

- ▶ operators⁵ make decisions
- ▶ faster horses v. cars

⁵In the sense of business deciders; that said, doctors, including those who operate, also have to make decisions, cf., personalized medicines

the natural abstraction

- ▶ operators⁵ make decisions
- ▶ faster horses v. cars
- ▶ general insights v. optimal policies

⁵In the sense of business deciders; that said, doctors, including those who operate, also have to make decisions, cf., personalized medicines

maximizing outcome

- ▶ the problem: maximizing an outcome over policies. . .

maximizing outcome

- ▶ the problem: maximizing an outcome over policies. . .
- ▶ . . . while inferring causality from observation

maximizing outcome

- ▶ the problem: maximizing an outcome over policies. . .
- ▶ . . . while inferring causality from observation
- ▶ different from predicting outcome in absence of action/policy

examples

- ▶ observation is not experiment

examples

- ▶ observation is not experiment
 - ▶ e.g., (Med.) smoking hurts vs unhealthy people smoke

examples

- ▶ observation is not experiment
 - ▶ e.g., (Med.) smoking hurts vs unhealthy people smoke
 - ▶ e.g., (Med.) affluent get prescribed different meds/treatment
-

examples

- ▶ observation is not experiment
 - ▶ e.g., (Med.) smoking hurts vs unhealthy people smoke
 - ▶ e.g., (Med.) affluent get prescribed different meds/treatment
 - ▶ e.g., (life) veterans earn less vs the rich serve less⁶

⁶Angrist, Joshua D. (1990). "Lifetime Earnings and the Vietnam Draft Lottery: Evidence from Social Security Administrative Records". American Economic Review 80 (3): 313–336.

examples

- ▶ observation is not experiment
 - ▶ e.g., (Med.) smoking hurts vs unhealthy people smoke
 - ▶ e.g., (Med.) affluent get prescribed different meds/treatment
 - ▶ e.g., (life) veterans earn less vs the rich serve less⁶
 - ▶ e.g., (life) admitted to school vs learn at school?

⁶Angrist, Joshua D. (1990). "Lifetime Earnings and the Vietnam Draft Lottery: Evidence from Social Security Administrative Records". American Economic Review 80 (3): 313–336.

reinforcement/machine learning/graphical models

- ▶ key idea: model joint $p(y, a, x)$

reinforcement/machine learning/graphical models

- ▶ key idea: model joint $p(y, a, x)$
- ▶ explore/exploit: family of joints $p_{\alpha}(y, a, x)$

reinforcement/machine learning/graphical models

- ▶ key idea: model joint $p(y, a, x)$
- ▶ explore/exploit: family of joints $p_{\alpha}(y, a, x)$
- ▶ “causality”: $p_{\alpha}(y, a, x) = p(y|a, x)p_{\alpha}(a|x)p(x)$ “a causes y”

reinforcement/machine learning/graphical models

- ▶ key idea: model joint $p(y, a, x)$
- ▶ explore/exploit: family of joints $p_{\alpha}(y, a, x)$
- ▶ “causality”: $p_{\alpha}(y, a, x) = p(y|a, x)p_{\alpha}(a|x)p(x)$ “a causes y”
- ▶ nomenclature: ‘response’, ‘policy’/‘bias’, ‘prior’ above

in general




Figure 9: policy/bias, response, and prior define the distribution

also describes both the 'exploration' and 'exploitation' distributions

randomized controlled trial




Figure 10: RCT: 'bias' removed, random 'policy' (response and prior unaffected)

also Pearl's 'do' distribution: a distribution with "no arrows" pointing to the action variable.

POISE: calculation, estimation, optimization

- ▶ POISE: “policy optimization via importance sample estimation”

POISE: calculation, estimation, optimization

- ▶ POISE: “policy optimization via importance sample estimation”
- ▶ Monte Carlo importance sampling estimation

POISE: calculation, estimation, optimization

- ▶ POISE: “policy optimization via importance sample estimation”
- ▶ Monte Carlo importance sampling estimation
 - ▶ aka “off policy estimation”

POISE: calculation, estimation, optimization

- ▶ POISE: “policy optimization via importance sample estimation”
- ▶ Monte Carlo importance sampling estimation
 - ▶ aka “off policy estimation”
 - ▶ role of “IPW”

POISE: calculation, estimation, optimization

- ▶ POISE: “policy optimization via importance sample estimation”
- ▶ Monte Carlo importance sampling estimation
 - ▶ aka “off policy estimation”
 - ▶ role of “IPW”
- ▶ reduction

POISE: calculation, estimation, optimization

- ▶ POISE: “policy optimization via importance sample estimation”
- ▶ Monte Carlo importance sampling estimation
 - ▶ aka “off policy estimation”
 - ▶ role of “IPW”
- ▶ reduction
- ▶ normalization

POISE: calculation, estimation, optimization

- ▶ POISE: “policy optimization via importance sample estimation”
- ▶ Monte Carlo importance sampling estimation
 - ▶ aka “off policy estimation”
 - ▶ role of “IPW”
- ▶ reduction
- ▶ normalization
- ▶ hyper-parameter searching

POISE: calculation, estimation, optimization

- ▶ POISE: “policy optimization via importance sample estimation”
- ▶ Monte Carlo importance sampling estimation
 - ▶ aka “off policy estimation”
 - ▶ role of “IPW”
- ▶ reduction
- ▶ normalization
- ▶ hyper-parameter searching
- ▶ unexpected connection: personalized medicine

POISE setup and Goal

- ▶ “a causes y” $\iff \exists$ family $p_\alpha(y, a, x) = p(y|a, x)p_\alpha(a|x)p(x)$

POISE setup and Goal

- ▶ “a causes y” $\iff \exists$ family $p_{\alpha}(y, a, x) = p(y|a, x)p_{\alpha}(a|x)p(x)$
- ▶ define off-policy/exploration distribution
 $p_{-}(y, a, x) = p(y|a, x)p_{-}(a|x)p(x)$

POISE setup and Goal

- ▶ “a causes y” $\iff \exists$ family $p_{\alpha}(y, a, x) = p(y|a, x)p_{\alpha}(a|x)p(x)$
- ▶ define off-policy/exploration distribution
$$p_{-}(y, a, x) = p(y|a, x)p_{-}(a|x)p(x)$$
- ▶ define exploitation distribution
$$p_{+}(y, a, x) = p(y|a, x)p_{+}(a|x)p(x)$$

POISE setup and Goal

- ▶ “a causes y” $\iff \exists$ family $p_{\alpha}(y, a, x) = p(y|a, x)p_{\alpha}(a|x)p(x)$
- ▶ define off-policy/exploration distribution
$$p_{-}(y, a, x) = p(y|a, x)p_{-}(a|x)p(x)$$
- ▶ define exploitation distribution
$$p_{+}(y, a, x) = p(y|a, x)p_{+}(a|x)p(x)$$
- ▶ Goal: Maximize $E_{+}(Y)$ over $p_{+}(a|x)$ using data drawn from $p_{-}(y, a, x)$.

POISE setup and Goal

- ▶ “a causes y” $\iff \exists$ family $p_{\alpha}(y, a, x) = p(y|a, x)p_{\alpha}(a|x)p(x)$
- ▶ define off-policy/exploration distribution
$$p_{-}(y, a, x) = p(y|a, x)p_{-}(a|x)p(x)$$
- ▶ define exploitation distribution
$$p_{+}(y, a, x) = p(y|a, x)p_{+}(a|x)p(x)$$
- ▶ Goal: Maximize $E_{+}(Y)$ over $p_{+}(a|x)$ using data drawn from $p_{-}(y, a, x)$.

POISE setup and Goal

- ▶ “a causes y” $\iff \exists$ family $p_{\alpha}(y, a, x) = p(y|a, x)p_{\alpha}(a|x)p(x)$
- ▶ define off-policy/exploration distribution
$$p_{-}(y, a, x) = p(y|a, x)p_{-}(a|x)p(x)$$
- ▶ define exploitation distribution
$$p_{+}(y, a, x) = p(y|a, x)p_{+}(a|x)p(x)$$
- ▶ Goal: Maximize $E_{+}(Y)$ over $p_{+}(a|x)$ using data drawn from $p_{-}(y, a, x)$.

notation: $\{x, a, y\} \in \{X, A, Y\}$ i.e., $E_{\alpha}(Y)$ is not a function of y

POISE math: IS+Monte Carlo estimation=ISE

i.e, “importance sampling estimation”

- ▶ $E_+(Y) \equiv \sum_{yax} y p_+(y, a, x)$

POISE math: IS+Monte Carlo estimation=ISE

i.e, “importance sampling estimation”

- ▶ $E_+(Y) \equiv \sum_{yax} yp_+(y, a, x)$
- ▶ $E_+(Y) = \sum_{yax} yp_-(y, a, x)(p_+(y, a, x)/p_-(y, a, x))$

POISE math: IS+Monte Carlo estimation=ISE

i.e, “importance sampling estimation”

- ▶ $E_+(Y) \equiv \sum_{yax} y p_+(y, a, x)$
- ▶ $E_+(Y) = \sum_{yax} y p_-(y, a, x) (p_+(y, a, x) / p_-(y, a, x))$
- ▶ $E_+(Y) = \sum_{yax} y p_-(y, a, x) (p_+(a|x) / p_-(a|x))$

POISE math: IS+Monte Carlo estimation=ISE

i.e, “importance sampling estimation”

- ▶ $E_+(Y) \equiv \sum_{yax} yp_+(y, a, x)$
- ▶ $E_+(Y) = \sum_{yax} yp_-(y, a, x)(p_+(y, a, x)/p_-(y, a, x))$
- ▶ $E_+(Y) = \sum_{yax} yp_-(y, a, x)(p_+(a|x)/p_-(a|x))$
- ▶ $E_+(Y) \approx N^{-1} \sum_i y_i(p_+(a_i|x_i)/p_-(a_i|x_i))$

POISE math: IS+Monte Carlo estimation=ISE

i.e, “importance sampling estimation”

- ▶ $E_+(Y) \equiv \sum_{yax} yp_+(y, a, x)$
- ▶ $E_+(Y) = \sum_{yax} yp_-(y, a, x)(p_+(y, a, x)/p_-(y, a, x))$
- ▶ $E_+(Y) = \sum_{yax} yp_-(y, a, x)(p_+(a|x)/p_-(a|x))$
- ▶ $E_+(Y) \approx N^{-1} \sum_i y_i(p_+(a_i|x_i)/p_-(a_i|x_i))$

POISE math: IS+Monte Carlo estimation=ISE

i.e, “importance sampling estimation”

- ▶ $E_+(Y) \equiv \sum_{yax} yp_+(y, a, x)$
- ▶ $E_+(Y) = \sum_{yax} yp_-(y, a, x)(p_+(y, a, x)/p_-(y, a, x))$
- ▶ $E_+(Y) = \sum_{yax} yp_-(y, a, x)(p_+(a|x)/p_-(a|x))$
- ▶ $E_+(Y) \approx N^{-1} \sum_i y_i(p_+(a_i|x_i)/p_-(a_i|x_i))$

let's spend some time getting to know this last equation, the importance sampling estimate of outcome in a “causal model” (“a causes y”) among $\{y, a, x\}$

Observation (cf. Bottou⁷)

- ▶ factorizing $P_{\pm}(x)$: $\frac{P_{+}(x)}{P_{-}(x)} = \prod_{\text{factors}} \frac{P_{+\text{but not}-}(x)}{P_{-\text{but not}+}(x)}$

Observation (cf. Bottou⁷)

- ▶ factorizing $P_{\pm}(x)$: $\frac{P_{+}(x)}{P_{-}(x)} = \prod_{\text{factors}} \frac{P_{+\text{but not}-}(x)}{P_{-\text{but not}+}(x)}$
- ▶ origin: importance sampling $E_q(f) = E_p(fq/p)$ (as in variational methods)

Observation (cf. Bottou⁷)

- ▶ factorizing $P_{\pm}(x)$: $\frac{P_{+}(x)}{P_{-}(x)} = \prod_{\text{factors}} \frac{P_{+\text{but not}-}(x)}{P_{-\text{but not}+}(x)}$
 - ▶ origin: importance sampling $E_q(f) = E_p(fq/p)$ (as in variational methods)
 - ▶ the “causal” model $p_{\alpha}(y, a, x) = p(y|a, x)p_{\alpha}(a|x)p(x)$ helps here
-

Observation (cf. Bottou⁷)

- ▶ factorizing $P_{\pm}(x)$: $\frac{P_{+}(x)}{P_{-}(x)} = \prod_{\text{factors}} \frac{P_{+\text{but not}-}(x)}{P_{-\text{but not}+}(x)}$
 - ▶ origin: importance sampling $E_q(f) = E_p(fq/p)$ (as in variational methods)
 - ▶ the “causal” model $p_{\alpha}(y, a, x) = p(y|a, x)p_{\alpha}(a|x)p(x)$ helps here
 - ▶ factors left over are numerator ($p_{+}(a|x)$, to optimize) and denominator ($p_{-}(a|x)$, to infer if not a RCT)
-

Observation (cf. Bottou⁷)

- ▶ factorizing $P_{\pm}(x)$: $\frac{P_{+}(x)}{P_{-}(x)} = \prod_{\text{factors}} \frac{P_{+\text{but not}-}(x)}{P_{-\text{but not}+}(x)}$
- ▶ origin: importance sampling $E_q(f) = E_p(fq/p)$ (as in variational methods)
- ▶ the “causal” model $p_{\alpha}(y, a, x) = p(y|a, x)p_{\alpha}(a|x)p(x)$ helps here
- ▶ factors left over are numerator ($p_{+}(a|x)$, to optimize) and denominator ($p_{-}(a|x)$, to infer if not a RCT)
- ▶ unobserved confounders will confound us (later)

⁷Counterfactual Reasoning and Learning Systems, arXiv:1209.2355

Reduction (cf. Langford^{8,9,10} ('05, '08, '09))

- ▶ consider numerator for deterministic policy:

$$p_+(a|x) = 1[a = h(x)]$$

Reduction (cf. Langford^{8,9,10} ('05, '08, '09))

- ▶ consider numerator for deterministic policy:

$$p_+(a|x) = 1[a = h(x)]$$

- ▶ $E_+(Y) \propto \sum_i (y_i / p_-(a|x)) 1[a = h(x)] \equiv \sum_i w_i 1[a = h(x)]$

Reduction (cf. Langford^{8,9,10} ('05, '08, '09))

- ▶ consider numerator for deterministic policy:

$$p_+(a|x) = 1[a = h(x)]$$

- ▶ $E_+(Y) \propto \sum_i (y_i / p_-(a|x)) 1[a = h(x)] \equiv \sum_i w_i 1[a = h(x)]$
 - ▶ Note: $1[c = d] = 1 - 1[c \neq d]$
-

Reduction (cf. Langford^{8,9,10} ('05, '08, '09))

- ▶ consider numerator for deterministic policy:

$$p_+(a|x) = 1[a = h(x)]$$

- ▶ $E_+(Y) \propto \sum_i (y_i / p_-(a|x)) 1[a = h(x)] \equiv \sum_i w_i 1[a = h(x)]$
 - ▶ Note: $1[c = d] = 1 - 1[c \neq d]$
 - ▶ $\therefore E_+(Y) \propto \text{constant} - \sum_i w_i 1[a \neq h(x)]$
-

Reduction (cf. Langford^{8,9,10} ('05, '08, '09))

- ▶ consider numerator for deterministic policy:
 $p_+(a|x) = 1[a = h(x)]$
- ▶ $E_+(Y) \propto \sum_i (y_i / p_-(a|x)) 1[a = h(x)] \equiv \sum_i w_i 1[a = h(x)]$
- ▶ Note: $1[c = d] = 1 - 1[c \neq d]$
- ▶ $\therefore E_+(Y) \propto \text{constant} - \sum_i w_i 1[a \neq h(x)]$
- ▶ \therefore reduces policy optimization to (weighted) classification

⁸Langford & Zadrozny “Relating Reinforcement Learning Performance to Classification Performance” ICML 2005

⁹Beygelzimer & Langford “The offset tree for learning with partial labels” (KDD 2009)

¹⁰Tutorial on “Reductions” (including at ICML 2009)

Reduction w/optimistic complication

- ▶ Prescription \iff classification $L = \sum_i w_i 1[a_i \neq h(x_i)]$

Reduction w/optimistic complication

- ▶ Prescription \iff classification $L = \sum_i w_i 1[a_i \neq h(x_i)]$
 - ▶ weight $w_i = y_i/p_-(a_i|x_i)$, inferred or RCT
-

Reduction w/optimistic complication

- ▶ Prescription \iff classification $L = \sum_i w_i 1[a_i \neq h(x_i)]$
 - ▶ weight $w_i = y_i/p_-(a_i|x_i)$, inferred or RCT
 - ▶ destroys measure by treating $p_-(a|x)$ differently than $1/p_-(a|x)$
-

Reduction w/optimistic complication

- ▶ Prescription \iff classification $L = \sum_i w_i 1[a_i \neq h(x_i)]$
 - ▶ weight $w_i = y_i/p_-(a_i|x_i)$, inferred or RCT
 - ▶ destroys measure by treating $p_-(a|x)$ differently than $1/p_-(a|x)$
 - ▶ normalize as $\tilde{L} \equiv \frac{\sum_i y_i 1[a_i \neq h(x_i)]/p_-(a_i|x_i)}{\sum_i 1[a_i \neq h(x_i)]/p_-(a_i|x_i)}$
-

Reduction w/optimistic complication

- ▶ Prescription \iff classification $L = \sum_i w_i 1[a_i \neq h(x_i)]$
 - ▶ weight $w_i = y_i/p_-(a_i|x_i)$, inferred or RCT
 - ▶ destroys measure by treating $p_-(a|x)$ differently than $1/p_-(a|x)$
 - ▶ normalize as $\tilde{L} \equiv \frac{\sum_i y_i 1[a_i \neq h(x_i)]/p_-(a_i|x_i)}{\sum_i 1[a_i \neq h(x_i)]/p_-(a_i|x_i)}$
 - ▶ destroys lovely reduction
-

Reduction w/optimistic complication

- ▶ Prescription \iff classification $L = \sum_i w_i 1[a_i \neq h(x_i)]$
- ▶ weight $w_i = y_i/p_-(a_i|x_i)$, inferred or RCT
- ▶ destroys measure by treating $p_-(a|x)$ differently than $1/p_-(a|x)$
- ▶ normalize as $\tilde{L} \equiv \frac{\sum_i y_i 1[a_i \neq h(x_i)]/p_-(a_i|x_i)}{\sum_i 1[a_i \neq h(x_i)]/p_-(a_i|x_i)}$
- ▶ destroys lovely reduction
- ▶ simply¹¹ $L(\lambda) = \sum_i (y_i - \lambda) 1[a_i \neq h(x_i)]/p_-(a_i|x_i)$

¹¹Suggestion by Dan Hsu

Reduction w/optimistic complication

- ▶ Prescription \iff classification $L = \sum_i w_i 1[a_i \neq h(x_i)]$
- ▶ weight $w_i = y_i/p_-(a_i|x_i)$, inferred or RCT
- ▶ destroys measure by treating $p_-(a|x)$ differently than $1/p_-(a|x)$
- ▶ normalize as $\tilde{L} \equiv \frac{\sum_i y_i 1[a_i \neq h(x_i)]/p_-(a_i|x_i)}{\sum_i 1[a_i \neq h(x_i)]/p_-(a_i|x_i)}$
- ▶ destroys lovely reduction
- ▶ simply¹¹ $L(\lambda) = \sum_i (y_i - \lambda) 1[a_i \neq h(x_i)]/p_-(a_i|x_i)$
- ▶ hidden here is a 2nd parameter, in classification, \therefore harder search

¹¹Suggestion by Dan Hsu

POISE punchlines

- ▶ allows policy planning even with implicit logged exploration data¹²

¹²Strehl, Alex, et al. "Learning from logged implicit exploration data." Advances in Neural Information Processing Systems. 2010.

POISE punchlines

- ▶ allows policy planning even with implicit logged exploration data¹²
- ▶ e.g., two hospital story

¹²Strehl, Alex, et al. "Learning from logged implicit exploration data." Advances in Neural Information Processing Systems. 2010.

POISE punchlines

- ▶ allows policy planning even with implicit logged exploration data¹²
- ▶ e.g., two hospital story
- ▶ “personalized medicine” is also a policy

¹²Strehl, Alex, et al. “Learning from logged implicit exploration data.” Advances in Neural Information Processing Systems. 2010.

POISE punchlines

- ▶ allows policy planning even with implicit logged exploration data¹²
- ▶ e.g., two hospital story
- ▶ “personalized medicine” is also a policy
- ▶ abundant data available, under-explored IMHO

¹²Strehl, Alex, et al. “Learning from logged implicit exploration data.” Advances in Neural Information Processing Systems. 2010.

tangent: causality as told by an economist

different, related goal

- ▶ they think in terms of ATE/ITE instead of policy

tangent: causality as told by an economist

different, related goal

- ▶ they think in terms of ATE/ITE instead of policy
 - ▶ ATE

tangent: causality as told by an economist

different, related goal

- ▶ they think in terms of ATE/ITE instead of policy

- ▶ ATE

- ▶ $\tau \equiv E_0(Y|a=1) - E_0(Y|a=0) \equiv Q(a=1) - Q(a=0)$

tangent: causality as told by an economist

different, related goal

- ▶ they think in terms of ATE/ITE instead of policy
 - ▶ ATE
 - ▶ $\tau \equiv E_0(Y|a=1) - E_0(Y|a=0) \equiv Q(a=1) - Q(a=0)$
 - ▶ CATE aka Individualized Treatment Effect (ITE)

tangent: causality as told by an economist

different, related goal

- ▶ they think in terms of ATE/ITE instead of policy
 - ▶ ATE
 - ▶ $\tau \equiv E_0(Y|a=1) - E_0(Y|a=0) \equiv Q(a=1) - Q(a=0)$
 - ▶ CATE aka Individualized Treatment Effect (ITE)
 - ▶ $\tau(x) \equiv E_0(Y|a=1, x) - E_0(Y|a=0, x)$

tangent: causality as told by an economist

different, related goal

- ▶ they think in terms of ATE/ITE instead of policy
 - ▶ ATE
 - ▶ $\tau \equiv E_0(Y|a=1) - E_0(Y|a=0) \equiv Q(a=1) - Q(a=0)$
 - ▶ CATE aka Individualized Treatment Effect (ITE)
 - ▶ $\tau(x) \equiv E_0(Y|a=1, x) - E_0(Y|a=0, x)$
 - ▶ $\equiv Q(a=1, x) - Q(a=0, x)$

Q-note: “generalizing” Monte Carlo w/kernels

► MC: $E_p(f) = \sum_x p(x)f(x) \approx N^{-1} \sum_{i \sim p} f(x_i)$

Q-note: “generalizing” Monte Carlo w/kernels

- ▶ MC: $E_p(f) = \sum_x p(x)f(x) \approx N^{-1} \sum_{i \sim p} f(x_i)$
- ▶ K: $p \approx N^{-1} \sum_i K(x|x_i)$

Q-note: “generalizing” Monte Carlo w/kernels

- ▶ MC: $E_p(f) = \sum_x p(x)f(x) \approx N^{-1} \sum_{i \sim p} f(x_i)$
- ▶ K: $p \approx N^{-1} \sum_i K(x|x_i)$
- ▶ $\Rightarrow \sum_x p(x)f(x) \approx N^{-1} \sum_i \sum_x f(x)K(x|x_i)$

Q-note: “generalizing” Monte Carlo w/kernels

- ▶ MC: $E_p(f) = \sum_x p(x)f(x) \approx N^{-1} \sum_{i \sim p} f(x_i)$
- ▶ K : $p \approx N^{-1} \sum_i K(x|x_i)$
- ▶ $\Rightarrow \sum_x p(x)f(x) \approx N^{-1} \sum_i \sum_x f(x)K(x|x_i)$
- ▶ K can be any normalized function, e.g., $K(x|x_i) = \delta_{x,x_i}$, which yields MC.

Q-note: “generalizing” Monte Carlo w/kernels

- ▶ MC: $E_p(f) = \sum_x p(x)f(x) \approx N^{-1} \sum_{i \sim p} f(x_i)$
- ▶ K : $p \approx N^{-1} \sum_i K(x|x_i)$
- ▶ $\Rightarrow \sum_x p(x)f(x) \approx N^{-1} \sum_i \sum_x f(x)K(x|x_i)$
- ▶ K can be any normalized function, e.g., $K(x|x_i) = \delta_{x,x_i}$, which yields MC.
- ▶ multivariate
$$E_p(f) \approx N^{-1} \sum_i \sum_{yax} f(y, a, x) K_1(y|y_i) K_2(a|a_i) K_3(x|x_i)$$

Q-note: application w/strata+matching, setup

Helps think about economists' approach:

$$\blacktriangleright Q(a, x) \equiv E(Y|a, x) = \sum_y y p(y|a, x) = \sum_y y \frac{p_-(y, a, x)}{p_-(a|x)p(x)}$$

Q-note: application w/strata+matching, setup

Helps think about economists' approach:

- ▶ $Q(a, x) \equiv E(Y|a, x) = \sum_y y p(y|a, x) = \sum_y y \frac{p_-(y, a, x)}{p_-(a|x)p(x)}$
- ▶ $= \frac{1}{p_-(a|x)p(x)} \sum_y y p_-(y, a, x)$

Q-note: application w/strata+matching, setup

Helps think about economists' approach:

- ▶ $Q(a, x) \equiv E(Y|a, x) = \sum_y y p(y|a, x) = \sum_y y \frac{p_-(y, a, x)}{p_-(a|x)p(x)}$
- ▶ $= \frac{1}{p_-(a|x)p(x)} \sum_y y p_-(y, a, x)$
- ▶ stratify x using $z(x)$ such that $\cup z = X$, and $\cap z, z' =$

Q-note: application w/strata+matching, setup

Helps think about economists' approach:

- ▶ $Q(a, x) \equiv E(Y|a, x) = \sum_y y p(y|a, x) = \sum_y y \frac{p_-(y, a, x)}{p_-(a|x)p(x)}$
- ▶ $= \frac{1}{p_-(a|x)p(x)} \sum_y y p_-(y, a, x)$
- ▶ stratify x using $z(x)$ such that $\cup z = X$, and $\cap z, z' =$
- ▶ $n(x) = \sum_i 1[z(x_i) = z(x)]$ = number of points in x 's stratum

Q-note: application w/strata+matching, setup

Helps think about economists' approach:

- ▶ $Q(a, x) \equiv E(Y|a, x) = \sum_y y p(y|a, x) = \sum_y y \frac{p_-(y, a, x)}{p_-(a|x)p(x)}$
- ▶ $= \frac{1}{p_-(a|x)p(x)} \sum_y y p_-(y, a, x)$
- ▶ stratify x using $z(x)$ such that $\cup z = X$, and $\cap z, z' =$
- ▶ $n(x) = \sum_i 1[z(x_i) = z(x)]$ = number of points in x 's stratum
- ▶ $\Omega(x) = \sum_{x'} 1[z(x') = z(x)]$ = area of x 's stratum

Q-note: application w/strata+matching, setup

Helps think about economists' approach:

- ▶ $Q(a, x) \equiv E(Y|a, x) = \sum_y y p(y|a, x) = \sum_y y \frac{p_-(y, a, x)}{p_-(a|x)p(x)}$
- ▶ $= \frac{1}{p_-(a|x)p(x)} \sum_y y p_-(y, a, x)$
- ▶ stratify x using $z(x)$ such that $\cup z = X$, and $\cap z, z' =$
- ▶ $n(x) = \sum_i 1[z(x_i) = z(x)]$ = number of points in x 's stratum
- ▶ $\Omega(x) = \sum_{x'} 1[z(x') = z(x)]$ = area of x 's stratum
- ▶ $\therefore K_3(x|x_i) = 1[z(x) = z(x_i)]/\Omega(x)$

Q-note: application w/strata+matching, setup

Helps think about economists' approach:

- ▶ $Q(a, x) \equiv E(Y|a, x) = \sum_y y p(y|a, x) = \sum_y y \frac{p_-(y, a, x)}{p_-(a|x)p(x)}$
- ▶ $= \frac{1}{p_-(a|x)p(x)} \sum_y y p_-(y, a, x)$
- ▶ stratify x using $z(x)$ such that $\cup z = X$, and $\cap z, z' =$
- ▶ $n(x) = \sum_i 1[z(x_i) = z(x)]$ = number of points in x 's stratum
- ▶ $\Omega(x) = \sum_{x'} 1[z(x') = z(x)]$ = area of x 's stratum
- ▶ $\therefore K_3(x|x_i) = 1[z(x) = z(x_i)]/\Omega(x)$
- ▶ as in MC, $K_1(y|y_i) = \delta_{y, y_i}$, $K_2(a|a_i) = \delta_{a, a_i}$

Q-note: application w/strata+matching, payoff

$$\blacktriangleright \sum_y y p_-(y, a, x) \approx N^{-1} \Omega(x)^{-1} \sum_{a_i=a, z(x_i)=z(x)} y_i$$

Q-note: application w/strata+matching, payoff

- ▶ $\sum_y y p_-(y, a, x) \approx N^{-1} \Omega(x)^{-1} \sum_{a_i=a, z(x_i)=z(x)} y_i$
- ▶ $p(x) \approx (n(x)/N) \Omega(x)^{-1}$

Q-note: application w/strata+matching, payoff

- ▶ $\sum_y y p_-(y, a, x) \approx N^{-1} \Omega(x)^{-1} \sum_{a_i=a, z(x_i)=z(x)} y_i$
- ▶ $p(x) \approx (n(x)/N) \Omega(x)^{-1}$
- ▶ $\therefore Q(a, x) \approx p_-(a|x)^{-1} n(x)^{-1} \sum_{a_i=a, z(x_i)=z(x)} y_i$

Q-note: application w/strata+matching, payoff

- ▶ $\sum_y y p_-(y, a, x) \approx N^{-1} \Omega(x)^{-1} \sum_{a_i=a, z(x_i)=z(x)} y_i$
- ▶ $p(x) \approx (n(x)/N) \Omega(x)^{-1}$
- ▶ $\therefore Q(a, x) \approx p_-(a|x)^{-1} n(x)^{-1} \sum_{a_i=a, z(x_i)=z(x)} y_i$

Q-note: application w/strata+matching, payoff

- ▶ $\sum_y y p_-(y, a, x) \approx N^{-1} \Omega(x)^{-1} \sum_{a_i=a, z(x_i)=z(x)} y_i$
- ▶ $p(x) \approx (n(x)/N) \Omega(x)^{-1}$
- ▶ $\therefore Q(a, x) \approx p_-(a|x)^{-1} n(x)^{-1} \sum_{a_i=a, z(x_i)=z(x)} y_i$

“matching” means: choose each z to contain 1 positive example & 1 negative example,

- ▶ $p_-(a|x) \approx 1/2, n(x) = 2$

Q-note: application w/strata+matching, payoff

- ▶ $\sum_y y p_-(y, a, x) \approx N^{-1} \Omega(x)^{-1} \sum_{a_i=a, z(x_i)=z(x)} y_i$
- ▶ $p(x) \approx (n(x)/N) \Omega(x)^{-1}$
- ▶ $\therefore Q(a, x) \approx p_-(a|x)^{-1} n(x)^{-1} \sum_{a_i=a, z(x_i)=z(x)} y_i$

“matching” means: choose each z to contain 1 positive example & 1 negative example,

- ▶ $p_-(a|x) \approx 1/2, n(x) = 2$
- ▶ $\therefore \tau(a, x) = Q(a = 1, x) - Q(a = 0, x) = y_1(x) - y_0(x)$

Q-note: application w/strata+matching, payoff

- ▶ $\sum_y y p_-(y, a, x) \approx N^{-1} \Omega(x)^{-1} \sum_{a_i=a, z(x_i)=z(x)} y_i$
- ▶ $p(x) \approx (n(x)/N) \Omega(x)^{-1}$
- ▶ $\therefore Q(a, x) \approx p_-(a|x)^{-1} n(x)^{-1} \sum_{a_i=a, z(x_i)=z(x)} y_i$

“matching” means: choose each z to contain 1 positive example & 1 negative example,

- ▶ $p_-(a|x) \approx 1/2, n(x) = 2$
- ▶ $\therefore \tau(a, x) = Q(a = 1, x) - Q(a = 0, x) = y_1(x) - y_0(x)$
- ▶ z -generalizations: graphs, digraphs, k -NN, “matching”

Q-note: application w/strata+matching, payoff

- ▶ $\sum_y y p_-(y, a, x) \approx N^{-1} \Omega(x)^{-1} \sum_{a_i=a, z(x_i)=z(x)} y_i$
- ▶ $p(x) \approx (n(x)/N) \Omega(x)^{-1}$
- ▶ $\therefore Q(a, x) \approx p_-(a|x)^{-1} n(x)^{-1} \sum_{a_i=a, z(x_i)=z(x)} y_i$

“matching” means: choose each z to contain 1 positive example & 1 negative example,

- ▶ $p_-(a|x) \approx 1/2, n(x) = 2$
- ▶ $\therefore \tau(a, x) = Q(a = 1, x) - Q(a = 0, x) = y_1(x) - y_0(x)$
- ▶ z -generalizations: graphs, digraphs, k -NN, “matching”
- ▶ K -generalizations: continuous a , any metric or similarity you like, . . .

Q-note: application w/strata+matching, payoff

- ▶ $\sum_y y p_-(y, a, x) \approx N^{-1} \Omega(x)^{-1} \sum_{a_i=a, z(x_i)=z(x)} y_i$
- ▶ $p(x) \approx (n(x)/N) \Omega(x)^{-1}$
- ▶ $\therefore Q(a, x) \approx p_-(a|x)^{-1} n(x)^{-1} \sum_{a_i=a, z(x_i)=z(x)} y_i$

“matching” means: choose each z to contain 1 positive example & 1 negative example,

- ▶ $p_-(a|x) \approx 1/2, n(x) = 2$
- ▶ $\therefore \tau(a, x) = Q(a = 1, x) - Q(a = 0, x) = y_1(x) - y_0(x)$
- ▶ z -generalizations: graphs, digraphs, k -NN, “matching”
- ▶ K -generalizations: continuous a , any metric or similarity you like, . . .

Q-note: application w/strata+matching, payoff

- ▶ $\sum_y y p_-(y, a, x) \approx N^{-1} \Omega(x)^{-1} \sum_{a_i=a, z(x_i)=z(x)} y_i$
- ▶ $p(x) \approx (n(x)/N) \Omega(x)^{-1}$
- ▶ $\therefore Q(a, x) \approx p_-(a|x)^{-1} n(x)^{-1} \sum_{a_i=a, z(x_i)=z(x)} y_i$

“matching” means: choose each z to contain 1 positive example & 1 negative example,

- ▶ $p_-(a|x) \approx 1/2, n(x) = 2$
- ▶ $\therefore \tau(a, x) = Q(a = 1, x) - Q(a = 0, x) = y_1(x) - y_0(x)$
- ▶ z -generalizations: graphs, digraphs, k -NN, “matching”
- ▶ K -generalizations: continuous a , any metric or similarity you like, . . .

IMHO underexplored

causality, as understood in marketing

- ▶ a/b testing and RCT




Figure 11: Blattberg, Robert C., Byung-Do Kim, and Scott A. Neslin. Database Marketing, Springer New York, 2008

causality, as understood in marketing

- ▶ a/b testing and RCT
- ▶ yield optimization




Figure 11: Blattberg, Robert C., Byung-Do Kim, and Scott A. Neslin. Database Marketing, Springer New York, 2008

causality, as understood in marketing

- ▶ a/b testing and RCT
- ▶ yield optimization
- ▶ Lorenz curve (vs ROC plots)




Figure 11: Blattberg, Robert C., Byung-Do Kim, and Scott A. Neslin. Database Marketing, Springer New York, 2008

unobserved confounders vs. “causality” modeling

- ▶ truth: $p_{\alpha}(y, a, x, u) = p(y|a, x, u)p_{\alpha}(a|x, u)p(x, u)$

unobserved confounders vs. “causality” modeling

- ▶ truth: $p_{\alpha}(y, a, x, u) = p(y|a, x, u)p_{\alpha}(a|x, u)p(x, u)$
- ▶ but: $p_{+}(y, a, x, u) = p(y|a, x, u)p_{-}(a|x)p(x, u)$

unobserved confounders vs. “causality” modeling

- ▶ truth: $p_{\alpha}(y, a, x, u) = p(y|a, x, u)p_{\alpha}(a|x, u)p(x, u)$
- ▶ but: $p_{+}(y, a, x, u) = p(y|a, x, u)p_{-}(a|x)p(x, u)$
- ▶ $E_{+}(Y) \equiv \sum_{yaxu} yp_{+}(yaxu) \approx$
 $N^{-1} \sum_{i \sim p_{-}} y_i p_{+}(a|x) / p_{-}(a|x, u)$

unobserved confounders vs. “causality” modeling

- ▶ truth: $p_{\alpha}(y, a, x, u) = p(y|a, x, u)p_{\alpha}(a|x, u)p(x, u)$
- ▶ but: $p_{+}(y, a, x, u) = p(y|a, x, u)p_{-}(a|x)p(x, u)$
- ▶ $E_{+}(Y) \equiv \sum_{yaxu} yp_{+}(yaxu) \approx$
 $N^{-1} \sum_{i \sim p_{-}} y_i p_{+}(a|x) / p_{-}(a|x, u)$
- ▶ denominator can not be inferred, ignore at your peril

cautionary tale problem: Simpson's paradox

- ▶ a : admissions ($a=1$: admitted, $a=0$: declined)

cautionary tale problem: Simpson's paradox

- ▶ a : admissions ($a=1$: admitted, $a=0$: declined)
- ▶ x : gender ($x=1$: female, $x=0$: male)

cautionary tale problem: Simpson's paradox

- ▶ a : admissions ($a=1$: admitted, $a=0$: declined)
 - ▶ x : gender ($x=1$: female, $x=0$: male)
 - ▶ lawsuit (1973): $.44 = p(a = 1|x = 0) > p(a = 1|x = 1) = .35$
-

cautionary tale problem: Simpson's paradox

- ▶ a : admissions ($a=1$: admitted, $a=0$: declined)
- ▶ x : gender ($x=1$: female, $x=0$: male)
- ▶ lawsuit (1973): $.44 = p(a = 1|x = 0) > p(a = 1|x = 1) = .35$
- ▶ 'resolved' by Bickel (1975)¹³ (See also Pearl¹⁴)

¹³P.J. Bickel, E.A. Hammel and J.W. O'Connell (1975). "Sex Bias in Graduate Admissions: Data From Berkeley". Science 187 (4175): 398–404

¹⁴Pearl, Judea (December 2013). "Understanding Simpson's paradox". UCLA Cognitive Systems Laboratory, Technical Report R-414.

cautionary tale problem: Simpson's paradox

- ▶ a : admissions ($a=1$: admitted, $a=0$: declined)
- ▶ x : gender ($x=1$: female, $x=0$: male)
- ▶ lawsuit (1973): $.44 = p(a = 1|x = 0) > p(a = 1|x = 1) = .35$
- ▶ 'resolved' by Bickel (1975)¹³ (See also Pearl¹⁴)
- ▶ u : unobserved department they applied to

¹³P.J. Bickel, E.A. Hammel and J.W. O'Connell (1975). "Sex Bias in Graduate Admissions: Data From Berkeley". Science 187 (4175): 398–404

¹⁴Pearl, Judea (December 2013). "Understanding Simpson's paradox". UCLA Cognitive Systems Laboratory, Technical Report R-414.

cautionary tale problem: Simpson's paradox

- ▶ a : admissions ($a=1$: admitted, $a=0$: declined)
- ▶ x : gender ($x=1$: female, $x=0$: male)
- ▶ lawsuit (1973): $.44 = p(a = 1|x = 0) > p(a = 1|x = 1) = .35$
- ▶ 'resolved' by Bickel (1975)¹³ (See also Pearl¹⁴)
- ▶ u : unobserved department they applied to
- ▶ $p(a|x) = \sum_{u=1}^{u=6} p(a|x, u)p(u|x)$

¹³P.J. Bickel, E.A. Hammel and J.W. O'Connell (1975). "Sex Bias in Graduate Admissions: Data From Berkeley". Science 187 (4175): 398–404

¹⁴Pearl, Judea (December 2013). "Understanding Simpson's paradox". UCLA Cognitive Systems Laboratory, Technical Report R-414.

cautionary tale problem: Simpson's paradox

- ▶ a : admissions ($a=1$: admitted, $a=0$: declined)
- ▶ x : gender ($x=1$: female, $x=0$: male)
- ▶ lawsuit (1973): $.44 = p(a = 1|x = 0) > p(a = 1|x = 1) = .35$
- ▶ 'resolved' by Bickel (1975)¹³ (See also Pearl¹⁴)
- ▶ u : unobserved department they applied to
- ▶ $p(a|x) = \sum_{u=1}^{u=6} p(a|x, u)p(u|x)$
- ▶ e.g., gender-blind: $p(a|1) - p(a|0) = p(a|u) \cdot (p(u|1) - p(u|0))$

¹³P.J. Bickel, E.A. Hammel and J.W. O'Connell (1975). "Sex Bias in Graduate Admissions: Data From Berkeley". Science 187 (4175): 398–404

¹⁴Pearl, Judea (December 2013). "Understanding Simpson's paradox". UCLA Cognitive Systems Laboratory, Technical Report R-414.

confounded approach: quasi-experiments + instruments ¹⁷

- ▶ Q: does engagement drive retention? (NYT, NFLX, ...)

confounded approach: quasi-experiments + instruments ¹⁷

- ▶ Q: does engagement drive retention? (NYT, NFLX, ...)
 - ▶ we don't directly control engagement

confounded approach: quasi-experiments + instruments ¹⁷

- ▶ Q: does engagement drive retention? (NYT, NFLX, ...)
 - ▶ we don't directly control engagement
 - ▶ nonetheless useful since many things can influence it
-

confounded approach: quasi-experiments + instruments ¹⁷

- ▶ Q: does engagement drive retention? (NYT, NFLX, ...)
 - ▶ we don't directly control engagement
 - ▶ nonetheless useful since many things can influence it
- ▶ Q: does serving in Vietnam war decrease earnings¹⁵?

¹⁵Angrist, Joshua D. "Lifetime earnings and the Vietnam era draft lottery: evidence from social security administrative records." The American Economic Review (1990): 313-336.

confounded approach: quasi-experiments + instruments ¹⁷

- ▶ Q: does engagement drive retention? (NYT, NFLX, ...)
 - ▶ we don't directly control engagement
 - ▶ nonetheless useful since many things can influence it
- ▶ Q: does serving in Vietnam war decrease earnings¹⁵?
 - ▶ US didn't directly control serving in Vietnam, either¹⁶

¹⁵Angrist, Joshua D. "Lifetime earnings and the Vietnam era draft lottery: evidence from social security administrative records." *The American Economic Review* (1990): 313-336.

¹⁶cf., George Bush, Donald Trump, Bill Clinton, Dick Cheney...

confounded approach: quasi-experiments + instruments ¹⁷

- ▶ Q: does engagement drive retention? (NYT, NFLX, ...)
 - ▶ we don't directly control engagement
 - ▶ nonetheless useful since many things can influence it
- ▶ Q: does serving in Vietnam war decrease earnings¹⁵?
 - ▶ US didn't directly control serving in Vietnam, either¹⁶
- ▶ requires **strong assumptions**, including linear model

¹⁵Angrist, Joshua D. "Lifetime earnings and the Vietnam era draft lottery: evidence from social security administrative records." The American Economic Review (1990): 313-336.

¹⁶cf., George Bush, Donald Trump, Bill Clinton, Dick Cheney...

¹⁷I thank Sinan Aral, MIT Sloan, for bringing this to my attention

IV: graphical model assumption




Figure 12: independence assumption

IV: graphical model assumption (sideways)




Figure 13: independence assumption

IV: review s/OLS/MOM/ (E is empirical average)

- ▶ *a endogenous*

IV: review s/OLS/MOM/ (E is empirical average)

- ▶ a endogenous

- ▶ e.g., $\exists u$ s.t. $p(y|a, x, u), p(a|x, u)$

IV: review s/OLS/MOM/ (E is empirical average)

- ▶ a endogenous
 - ▶ e.g., $\exists u$ s.t. $p(y|a, x, u), p(a|x, u)$
- ▶ linear ansatz: $y = \beta^T a + \epsilon$

IV: review s/OLS/MOM/ (E is empirical average)

- ▶ *a* endogenous

- ▶ e.g., $\exists u$ s.t. $p(y|a, x, u), p(a|x, u)$

- ▶ linear ansatz: $y = \beta^T a + \epsilon$

- ▶ if *a* exogenous (e.g., OLS), use $E[YA_j] = E[\beta^T AA_j] + E[\epsilon A_j]$
(note that $E[A_j A_k]$ gives square matrix; invert for β)

IV: review s/OLS/MOM/ (E is empirical average)

- ▶ *a* endogenous
 - ▶ e.g., $\exists u$ s.t. $p(y|a, x, u), p(a|x, u)$
- ▶ linear ansatz: $y = \beta^T a + \epsilon$
- ▶ if *a* exogenous (e.g., OLS), use $E[YA_j] = E[\beta^T AA_j] + E[\epsilon A_j]$
(note that $E[A_j A_k]$ gives square matrix; invert for β)
- ▶ add *instrument* x uncorrelated with ϵ

IV: review s/OLS/MOM/ (E is empirical average)

- ▶ *a* endogenous
 - ▶ e.g., $\exists u$ s.t. $p(y|a, x, u), p(a|x, u)$
- ▶ linear ansatz: $y = \beta^T a + \epsilon$
- ▶ if *a* exogenous (e.g., OLS), use $E[YA_j] = E[\beta^T AA_j] + E[\epsilon A_j]$
(note that $E[A_j A_k]$ gives square matrix; invert for β)
- ▶ add *instrument* x uncorrelated with ϵ
- ▶ $E[YX_k] = E[\beta^T AX_k] + E[\epsilon]E[X_k]$

IV: review s/OLS/MOM/ (E is empirical average)

- ▶ *a* endogenous

- ▶ e.g., $\exists u$ s.t. $p(y|a, x, u), p(a|x, u)$

- ▶ linear ansatz: $y = \beta^T a + \epsilon$

- ▶ if *a* exogenous (e.g., OLS), use $E[YA_j] = E[\beta^T AA_j] + E[\epsilon A_j]$
(note that $E[A_j A_k]$ gives square matrix; invert for β)

- ▶ add *instrument* x uncorrelated with ϵ

- ▶ $E[YX_k] = E[\beta^T AX_k] + E[\epsilon]E[X_k]$

- ▶ $E[Y] = E[\beta^T A] + E[\epsilon]$ (from ansatz)

IV: review s/OLS/MOM/ (E is empirical average)

- ▶ *a* endogenous
 - ▶ e.g., $\exists u$ s.t. $p(y|a, x, u), p(a|x, u)$
- ▶ linear ansatz: $y = \beta^T a + \epsilon$
- ▶ if *a* exogenous (e.g., OLS), use $E[YA_j] = E[\beta^T AA_j] + E[\epsilon A_j]$
(note that $E[A_j A_k]$ gives square matrix; invert for β)
- ▶ add *instrument* x uncorrelated with ϵ
- ▶ $E[YX_k] = E[\beta^T AX_k] + E[\epsilon]E[X_k]$
- ▶ $E[Y] = E[\beta^T A] + E[\epsilon]$ (from ansatz)
- ▶ $C(Y, X_k) = \beta^T C(A, X_k)$, not an “inversion” problem, requires “two stage regression”

IV: binary, binary case (aka “Wald estimator”)

- ▶ $y = \beta \mathbf{a} + \epsilon$

IV: binary, binary case (aka “Wald estimator”)

- ▶ $y = \beta a + \epsilon$
- ▶ $E(Y|x) = \beta E(A|x) + E(\epsilon)$, evaluate at $x = \{0, 1\}$

IV: binary, binary case (aka “Wald estimator”)

- ▶ $y = \beta a + \epsilon$
- ▶ $E(Y|x) = \beta E(A|x) + E(\epsilon)$, evaluate at $x = \{0, 1\}$
- ▶ $\beta = (E(Y|x = 1) - E(Y|x = 0)) / (E(A|x = 1) - E(A|x = 0))$.

bandits: obligatory slide




Figure 14: almost all the talks I've gone to on bandits have this image

bandits

- ▶ wide applicability: humane clinical trials, targeting, ...

bandits

- ▶ wide applicability: humane clinical trials, targeting, ...
- ▶ replace meetings with code

bandits

- ▶ wide applicability: humane clinical trials, targeting, ...
 - ▶ replace meetings with code
 - ▶ requires software engineering to replace decisions with, e.g., Javascript
-

bandits

- ▶ wide applicability: humane clinical trials, targeting, ...
 - ▶ replace meetings with code
 - ▶ requires software engineering to replace decisions with, e.g., Javascript
 - ▶ most useful if decisions or items get “stale” quickly
-

bandits

- ▶ wide applicability: humane clinical trials, targeting, ...
 - ▶ replace meetings with code
 - ▶ requires software engineering to replace decisions with, e.g., Javascript
 - ▶ most useful if decisions or items get “stale” quickly
 - ▶ less useful for one-off, major decisions to be “interpreted”
-

bandits

- ▶ wide applicability: humane clinical trials, targeting, ...
 - ▶ replace meetings with code
 - ▶ requires software engineering to replace decisions with, e.g., Javascript
 - ▶ most useful if decisions or items get “stale” quickly
 - ▶ less useful for one-off, major decisions to be “interpreted”
-

bandits

- ▶ wide applicability: humane clinical trials, targeting, ...
- ▶ replace meetings with code
- ▶ requires software engineering to replace decisions with, e.g., Javascript
- ▶ most useful if decisions or items get “stale” quickly
- ▶ less useful for one-off, major decisions to be “interpreted”

examples

- ▶ ϵ -greedy (no context, aka ‘vanilla’, aka ‘context-free’)
-

bandits

- ▶ wide applicability: humane clinical trials, targeting, ...
- ▶ replace meetings with code
- ▶ requires software engineering to replace decisions with, e.g., Javascript
- ▶ most useful if decisions or items get “stale” quickly
- ▶ less useful for one-off, major decisions to be “interpreted”

examples

- ▶ ϵ -greedy (no context, aka ‘vanilla’, aka ‘context-free’)
 - ▶ UCB1 (2002) (no context) + LinUCB (with context)
-

bandits

- ▶ wide applicability: humane clinical trials, targeting, ...
- ▶ replace meetings with code
- ▶ requires software engineering to replace decisions with, e.g., Javascript
- ▶ most useful if decisions or items get “stale” quickly
- ▶ less useful for one-off, major decisions to be “interpreted”

examples

- ▶ ϵ -greedy (no context, aka ‘vanilla’, aka ‘context-free’)
- ▶ UCB1 (2002) (no context) + LinUCB (with context)
- ▶ Thompson Sampling (1933)^{18,19,20} (general, with or without context)

¹⁸Thompson, William R. “On the likelihood that one unknown probability exceeds another in view of the evidence of two samples”. *Biometrika*, 25(3–4):285–294, 1933.

¹⁹AKA “probability matching”, “posterior sampling”

²⁰cf., “Bayesian Bandit Explorer” ([link](#))

TS: connecting w/ “generative causal modeling” 0

- ▶ WAS $p(y, x, a) = p(y|x, a)p_{\alpha}(a|x)p(x)$

TS: connecting w/ “generative causal modeling” 0

- ▶ WAS $p(y, x, a) = p(y|x, a)p_{\alpha}(a|x)p(x)$
- ▶ These 3 terms were treated by

TS: connecting w/ “generative causal modeling” 0

- ▶ WAS $p(y, x, a) = p(y|x, a)p_{\alpha}(a|x)p(x)$
- ▶ These 3 terms were treated by
 - ▶ response $p(y|a, x)$: avoid regression/infering using importance sampling

TS: connecting w/ “generative causal modeling” 0

- ▶ WAS $p(y, x, a) = p(y|x, a)p_{\alpha}(a|x)p(x)$
- ▶ These 3 terms were treated by
 - ▶ response $p(y|a, x)$: avoid regression/infering using importance sampling
 - ▶ policy $p_{\alpha}(a|x)$: optimize ours, infer theirs

TS: connecting w/ “generative causal modeling” 0

- ▶ WAS $p(y, x, a) = p(y|x, a)p_\alpha(a|x)p(x)$
- ▶ These 3 terms were treated by
 - ▶ response $p(y|a, x)$: avoid regression/infering using importance sampling
 - ▶ policy $p_\alpha(a|x)$: optimize ours, infer theirs
 - ▶ (NB: ours was deterministic: $p(a|x) = 1[a = h(x)]$)

TS: connecting w/“generative causal modeling” 0

- ▶ WAS $p(y, x, a) = p(y|x, a)p_{\alpha}(a|x)p(x)$
- ▶ These 3 terms were treated by
 - ▶ response $p(y|a, x)$: avoid regression/infering using importance sampling
 - ▶ policy $p_{\alpha}(a|x)$: optimize ours, infer theirs
 - ▶ (NB: ours was deterministic: $p(a|x) = 1[a = h(x)]$)
 - ▶ prior $p(x)$: either avoid by importance sampling or estimate via kernel methods

TS: connecting w/“generative causal modeling” 0

- ▶ WAS $p(y, x, a) = p(y|x, a)p_{\alpha}(a|x)p(x)$
- ▶ These 3 terms were treated by
 - ▶ response $p(y|a, x)$: avoid regression/infering using importance sampling
 - ▶ policy $p_{\alpha}(a|x)$: optimize ours, infer theirs
 - ▶ (NB: ours was deterministic: $p(a|x) = 1[a = h(x)]$)
 - ▶ prior $p(x)$: either avoid by importance sampling or estimate via kernel methods
- ▶ In the economics approach we focus on

TS: connecting w/ “generative causal modeling” 0

- ▶ WAS $p(y, x, a) = p(y|x, a)p_\alpha(a|x)p(x)$
- ▶ These 3 terms were treated by
 - ▶ response $p(y|a, x)$: avoid regression/infering using importance sampling
 - ▶ policy $p_\alpha(a|x)$: optimize ours, infer theirs
 - ▶ (NB: ours was deterministic: $p(a|x) = 1[a = h(x)]$)
 - ▶ prior $p(x)$: either avoid by importance sampling or estimate via kernel methods
- ▶ In the economics approach we focus on
- ▶ $\tau(\dots) \equiv Q(a = 1, \dots) - Q(a = 0, \dots)$ “treatment effect”

TS: connecting w/ “generative causal modeling” 0

- ▶ WAS $p(y, x, a) = p(y|x, a)p_\alpha(a|x)p(x)$
- ▶ These 3 terms were treated by
 - ▶ response $p(y|a, x)$: avoid regression/infering using importance sampling
 - ▶ policy $p_\alpha(a|x)$: optimize ours, infer theirs
 - ▶ (NB: ours was deterministic: $p(a|x) = 1[a = h(x)]$)
 - ▶ prior $p(x)$: either avoid by importance sampling or estimate via kernel methods
- ▶ In the economics approach we focus on
- ▶ $\tau(\dots) \equiv Q(a = 1, \dots) - Q(a = 0, \dots)$ “treatment effect”
- ▶ where $Q(a, \dots) = \sum_y yp(y|\dots)$

TS: connecting w/ “generative causal modeling” 0

- ▶ WAS $p(y, x, a) = p(y|x, a)p_\alpha(a|x)p(x)$
- ▶ These 3 terms were treated by
 - ▶ response $p(y|a, x)$: avoid regression/infering using importance sampling
 - ▶ policy $p_\alpha(a|x)$: optimize ours, infer theirs
 - ▶ (NB: ours was deterministic: $p(a|x) = 1[a = h(x)]$)
 - ▶ prior $p(x)$: either avoid by importance sampling or estimate via kernel methods
- ▶ In the economics approach we focus on
- ▶ $\tau(\dots) \equiv Q(a = 1, \dots) - Q(a = 0, \dots)$ “treatment effect”
- ▶ where $Q(a, \dots) = \sum_y yp(y|\dots)$

TS: connecting w/“generative causal modeling” 0

- ▶ WAS $p(y, x, a) = p(y|x, a)p_\alpha(a|x)p(x)$
- ▶ These 3 terms were treated by
 - ▶ response $p(y|a, x)$: avoid regression/infering using importance sampling
 - ▶ policy $p_\alpha(a|x)$: optimize ours, infer theirs
 - ▶ (NB: ours was deterministic: $p(a|x) = 1[a = h(x)]$)
 - ▶ prior $p(x)$: either avoid by importance sampling or estimate via kernel methods
- ▶ In the economics approach we focus on
- ▶ $\tau(\dots) \equiv Q(a = 1, \dots) - Q(a = 0, \dots)$ “treatment effect”
- ▶ where $Q(a, \dots) = \sum_y yp(y|\dots)$

In Thompson sampling we will generate 1 datum at a time, by

- ▶ asserting a parameterized generative model for $p(y|a, x, \theta)$

TS: connecting w/“generative causal modeling” 0

- ▶ WAS $p(y, x, a) = p(y|x, a)p_\alpha(a|x)p(x)$
- ▶ These 3 terms were treated by
 - ▶ response $p(y|a, x)$: avoid regression/infering using importance sampling
 - ▶ policy $p_\alpha(a|x)$: optimize ours, infer theirs
 - ▶ (NB: ours was deterministic: $p(a|x) = 1[a = h(x)]$)
 - ▶ prior $p(x)$: either avoid by importance sampling or estimate via kernel methods
- ▶ In the economics approach we focus on
- ▶ $\tau(\dots) \equiv Q(a = 1, \dots) - Q(a = 0, \dots)$ “treatment effect”
- ▶ where $Q(a, \dots) = \sum_y yp(y|\dots)$

In Thompson sampling we will generate 1 datum at a time, by

- ▶ asserting a parameterized generative model for $p(y|a, x, \theta)$
- ▶ using a deterministic but averaged policy

TS: connecting w/“generative causal modeling” 1

- ▶ model true world response function $p(y|a, x)$ parametrically as $p(y|a, x, \theta^*)$

TS: connecting w/“generative causal modeling” 1

- ▶ model true world response function $p(y|a, x)$ parametrically as $p(y|a, x, \theta^*)$
- ▶ (i.e., θ^* is the true value of the parameter)²¹

²¹Note that θ is a vector, with components for each action.

TS: connecting w/“generative causal modeling” 1

- ▶ model true world response function $p(y|a, x)$ parametrically as $p(y|a, x, \theta^*)$
- ▶ (i.e., θ^* is the true value of the parameter)²¹
- ▶ if you knew θ :

²¹Note that θ is a vector, with components for each action.

TS: connecting w/“generative causal modeling” 1

- ▶ model true world response function $p(y|a, x)$ parametrically as $p(y|a, x, \theta^*)$
- ▶ (i.e., θ^* is the true value of the parameter)²¹
- ▶ if you knew θ :
 - ▶ could compute $Q(a, x, \theta) \equiv \sum_y yp(y|x, a, \theta^*)$ directly

²¹Note that θ is a vector, with components for each action.

TS: connecting w/“generative causal modeling” 1

- ▶ model true world response function $p(y|a, x)$ parametrically as $p(y|a, x, \theta^*)$
- ▶ (i.e., θ^* is the true value of the parameter)²¹
- ▶ if you knew θ :
 - ▶ could compute $Q(a, x, \theta) \equiv \sum_y yp(y|x, a, \theta^*)$ directly
 - ▶ then choose $h(x; \theta) = \operatorname{argmax}_a Q(a, x, \theta)$

²¹Note that θ is a vector, with components for each action.

TS: connecting w/“generative causal modeling” 1

- ▶ model true world response function $p(y|a, x)$ parametrically as $p(y|a, x, \theta^*)$
- ▶ (i.e., θ^* is the true value of the parameter)²¹
- ▶ if you knew θ :
 - ▶ could compute $Q(a, x, \theta) \equiv \sum_y yp(y|x, a, \theta^*)$ directly
 - ▶ then choose $h(x; \theta) = \operatorname{argmax}_a Q(a, x, \theta)$
 - ▶ inducing policy $p(a|x, \theta) = 1[a = h(x; \theta) = \operatorname{argmax}_a Q(a, x, \theta)]$

²¹Note that θ is a vector, with components for each action.

TS: connecting w/“generative causal modeling” 1

- ▶ model true world response function $p(y|a, x)$ parametrically as $p(y|a, x, \theta^*)$
- ▶ (i.e., θ^* is the true value of the parameter)²¹
- ▶ if you knew θ :
 - ▶ could compute $Q(a, x, \theta) \equiv \sum_y yp(y|x, a, \theta^*)$ directly
 - ▶ then choose $h(x; \theta) = \operatorname{argmax}_a Q(a, x, \theta)$
 - ▶ inducing policy $p(a|x, \theta) = 1[a = h(x; \theta) = \operatorname{argmax}_a Q(a, x, \theta)]$
- ▶ idea: use prior data $D = \{y, a, x\}_1^t$ to define *non-deterministic* policy:

²¹Note that θ is a vector, with components for each action.

TS: connecting w/ “generative causal modeling” 1

- ▶ model true world response function $p(y|a, x)$ parametrically as $p(y|a, x, \theta^*)$
- ▶ (i.e., θ^* is the true value of the parameter)²¹
- ▶ if you knew θ :
 - ▶ could compute $Q(a, x, \theta) \equiv \sum_y y p(y|x, a, \theta^*)$ directly
 - ▶ then choose $h(x; \theta) = \operatorname{argmax}_a Q(a, x, \theta)$
 - ▶ inducing policy $p(a|x, \theta) = 1[a = h(x; \theta) = \operatorname{argmax}_a Q(a, x, \theta)]$
- ▶ idea: use prior data $D = \{y, a, x\}_1^t$ to define *non-deterministic* policy:
 - ▶ $p(a|x) = \int d\theta p(a|x, \theta) p(\theta|D)$

²¹Note that θ is a vector, with components for each action.

TS: connecting w/ “generative causal modeling” 1

- ▶ model true world response function $p(y|a, x)$ parametrically as $p(y|a, x, \theta^*)$
- ▶ (i.e., θ^* is the true value of the parameter)²¹
- ▶ if you knew θ :
 - ▶ could compute $Q(a, x, \theta) \equiv \sum_y y p(y|x, a, \theta^*)$ directly
 - ▶ then choose $h(x; \theta) = \operatorname{argmax}_a Q(a, x, \theta)$
 - ▶ inducing policy $p(a|x, \theta) = 1[a = h(x; \theta) = \operatorname{argmax}_a Q(a, x, \theta)]$
- ▶ idea: use prior data $D = \{y, a, x\}_1^t$ to define *non-deterministic* policy:
 - ▶ $p(a|x) = \int d\theta p(a|x, \theta) p(\theta|D)$
 - ▶ $p(a|x) = \int d\theta 1[a = \operatorname{argmax}_{a'} Q(a', x, \theta)] p(\theta|D)$

²¹Note that θ is a vector, with components for each action.

TS: connecting w/ “generative causal modeling” 1

- ▶ model true world response function $p(y|a, x)$ parametrically as $p(y|a, x, \theta^*)$
- ▶ (i.e., θ^* is the true value of the parameter)²¹
- ▶ if you knew θ :
 - ▶ could compute $Q(a, x, \theta) \equiv \sum_y y p(y|x, a, \theta)$ directly
 - ▶ then choose $h(x; \theta) = \operatorname{argmax}_a Q(a, x, \theta)$
 - ▶ inducing policy $p(a|x, \theta) = 1[a = h(x; \theta) = \operatorname{argmax}_a Q(a, x, \theta)]$
- ▶ idea: use prior data $D = \{y, a, x\}_1^t$ to define *non-deterministic* policy:
 - ▶ $p(a|x) = \int d\theta p(a|x, \theta) p(\theta|D)$
 - ▶ $p(a|x) = \int d\theta 1[a = \operatorname{argmax}_{a'} Q(a', x, \theta)] p(\theta|D)$
- ▶ hold up:

²¹Note that θ is a vector, with components for each action.

TS: connecting w/ “generative causal modeling” 1

- ▶ model true world response function $p(y|a, x)$ parametrically as $p(y|a, x, \theta^*)$
- ▶ (i.e., θ^* is the true value of the parameter)²¹
- ▶ if you knew θ :
 - ▶ could compute $Q(a, x, \theta) \equiv \sum_y y p(y|x, a, \theta^*)$ directly
 - ▶ then choose $h(x; \theta) = \operatorname{argmax}_a Q(a, x, \theta)$
 - ▶ inducing policy $p(a|x, \theta) = 1[a = h(x; \theta) = \operatorname{argmax}_a Q(a, x, \theta)]$
- ▶ idea: use prior data $D = \{y, a, x\}_1^t$ to define *non-deterministic* policy:
 - ▶ $p(a|x) = \int d\theta p(a|x, \theta) p(\theta|D)$
 - ▶ $p(a|x) = \int d\theta 1[a = \operatorname{argmax}_{a'} Q(a', x, \theta)] p(\theta|D)$
- ▶ hold up:
 - ▶ Q1: what's $p(\theta|D)$?

²¹Note that θ is a vector, with components for each action.

TS: connecting w/ “generative causal modeling” 1

- ▶ model true world response function $p(y|a, x)$ parametrically as $p(y|a, x, \theta^*)$
- ▶ (i.e., θ^* is the true value of the parameter)²¹
- ▶ if you knew θ :
 - ▶ could compute $Q(a, x, \theta) \equiv \sum_y y p(y|x, a, \theta^*)$ directly
 - ▶ then choose $h(x; \theta) = \operatorname{argmax}_a Q(a, x, \theta)$
 - ▶ inducing policy $p(a|x, \theta) = 1[a = h(x; \theta) = \operatorname{argmax}_a Q(a, x, \theta)]$
- ▶ idea: use prior data $D = \{y, a, x\}_1^t$ to define *non-deterministic* policy:
 - ▶ $p(a|x) = \int d\theta p(a|x, \theta) p(\theta|D)$
 - ▶ $p(a|x) = \int d\theta 1[a = \operatorname{argmax}_{a'} Q(a', x, \theta)] p(\theta|D)$
- ▶ hold up:
 - ▶ Q1: what's $p(\theta|D)$?
 - ▶ Q2: how am I going to evaluate this integral?

²¹Note that θ is a vector, with components for each action.

TS: connecting w/ “generative causal modeling” 2

- ▶ Q1: what's $p(\theta|D)$?

TS: connecting w/ “generative causal modeling” 2

- ▶ Q1: what's $p(\theta|D)$?
- ▶ Q2: how am I going to evaluate this integral?

TS: connecting w/“generative causal modeling” 2

- ▶ Q1: what's $p(\theta|D)$?
- ▶ Q2: how am I going to evaluate this integral?
- ▶ A1: $p(\theta|D)$ definable by choosing prior $p(\theta|\alpha)$ and likelihood on y given by the (modeled, parameterized) response $p(y|a, x, \theta)$.

TS: connecting w/“generative causal modeling” 2

- ▶ Q1: what's $p(\theta|D)$?
- ▶ Q2: how am I going to evaluate this integral?
- ▶ A1: $p(\theta|D)$ definable by choosing prior $p(\theta|\alpha)$ and likelihood on y given by the (modeled, parameterized) response $p(y|a, x, \theta)$.
 - ▶ (now you're not only generative, you're Bayesian.)

TS: connecting w/“generative causal modeling” 2

- ▶ Q1: what's $p(\theta|D)$?
- ▶ Q2: how am I going to evaluate this integral?
- ▶ A1: $p(\theta|D)$ definable by choosing prior $p(\theta|\alpha)$ and likelihood on y given by the (modeled, parameterized) response $p(y|a, x, \theta)$.
 - ▶ (now you're not only generative, you're Bayesian.)
 - ▶ $p(\theta|D) = p(\theta|\{y\}_1^t, \{a\}_1^t, \{x\}_1^t, \alpha)$

TS: connecting w/“generative causal modeling” 2

- ▶ Q1: what's $p(\theta|D)$?
- ▶ Q2: how am I going to evaluate this integral?
- ▶ A1: $p(\theta|D)$ definable by choosing prior $p(\theta|\alpha)$ and likelihood on y given by the (modeled, parameterized) response $p(y|a, x, \theta)$.
 - ▶ (now you're not only generative, you're Bayesian.)
 - ▶ $p(\theta|D) = p(\theta|\{y\}_1^t, \{a\}_1^t, \{x\}_1^t, \alpha)$
 - ▶ $\propto p(\{y\}_1^t|\{a\}_1^t, \{x\}_1^t, \theta)p(\theta|\alpha)$

TS: connecting w/“generative causal modeling” 2

- ▶ Q1: what's $p(\theta|D)$?
- ▶ Q2: how am I going to evaluate this integral?
- ▶ A1: $p(\theta|D)$ definable by choosing prior $p(\theta|\alpha)$ and likelihood on y given by the (modeled, parameterized) response $p(y|a, x, \theta)$.
 - ▶ (now you're not only generative, you're Bayesian.)
 - ▶ $p(\theta|D) = p(\theta|\{y\}_1^t, \{a\}_1^t, \{x\}_1^t, \alpha)$
 - ▶ $\propto p(\{y\}_1^t|\{a\}_1^t, \{x\}_1^t, \theta)p(\theta|\alpha)$
 - ▶ $= p(\theta|\alpha)\prod_t p(y_t|a_t, x_t, \theta)$

TS: connecting w/“generative causal modeling” 2

- ▶ Q1: what's $p(\theta|D)$?
- ▶ Q2: how am I going to evaluate this integral?
- ▶ A1: $p(\theta|D)$ definable by choosing prior $p(\theta|\alpha)$ and likelihood on y given by the (modeled, parameterized) response $p(y|a, x, \theta)$.
 - ▶ (now you're not only generative, you're Bayesian.)
 - ▶ $p(\theta|D) = p(\theta|\{y\}_1^t, \{a\}_1^t, \{x\}_1^t, \alpha)$
 - ▶ $\propto p(\{y\}_1^t|\{a\}_1^t, \{x\}_1^t, \theta)p(\theta|\alpha)$
 - ▶ $= p(\theta|\alpha)\prod_t p(y_t|a_t, x_t, \theta)$
 - ▶ *warning 1*: sometimes people write “ $p(D|\theta)$ ” but we don't need $p(a|\theta)$ or $p(x|\theta)$ here

TS: connecting w/“generative causal modeling” 2

- ▶ Q1: what's $p(\theta|D)$?
- ▶ Q2: how am I going to evaluate this integral?
- ▶ A1: $p(\theta|D)$ definable by choosing prior $p(\theta|\alpha)$ and likelihood on y given by the (modeled, parameterized) response $p(y|a, x, \theta)$.
 - ▶ (now you're not only generative, you're Bayesian.)
 - ▶ $p(\theta|D) = p(\theta|\{y\}_1^t, \{a\}_1^t, \{x\}_1^t, \alpha)$
 - ▶ $\propto p(\{y\}_1^t|\{a\}_1^t, \{x\}_1^t, \theta)p(\theta|\alpha)$
 - ▶ $= p(\theta|\alpha)\prod_t p(y_t|a_t, x_t, \theta)$
 - ▶ *warning 1*: sometimes people write “ $p(D|\theta)$ ” but we don't need $p(a|\theta)$ or $p(x|\theta)$ here
 - ▶ *warning 2*: don't need historical record of θ_t .

TS: connecting w/“generative causal modeling” 2

- ▶ Q1: what's $p(\theta|D)$?
- ▶ Q2: how am I going to evaluate this integral?
- ▶ A1: $p(\theta|D)$ definable by choosing prior $p(\theta|\alpha)$ and likelihood on y given by the (modeled, parameterized) response $p(y|a, x, \theta)$.
 - ▶ (now you're not only generative, you're Bayesian.)
 - ▶ $p(\theta|D) = p(\theta|\{y\}_1^t, \{a\}_1^t, \{x\}_1^t, \alpha)$
 - ▶ $\propto p(\{y\}_1^t|\{a\}_1^t, \{x\}_1^t, \theta)p(\theta|\alpha)$
 - ▶ $= p(\theta|\alpha)\prod_t p(y_t|a_t, x_t, \theta)$
 - ▶ *warning 1*: sometimes people write “ $p(D|\theta)$ ” but we don't need $p(a|\theta)$ or $p(x|\theta)$ here
 - ▶ *warning 2*: don't need historical record of θ_t .
 - ▶ (we used Bayes rule, but only in θ and y .)

TS: connecting w/“generative causal modeling” 2

- ▶ Q1: what's $p(\theta|D)$?
- ▶ Q2: how am I going to evaluate this integral?
- ▶ A1: $p(\theta|D)$ definable by choosing prior $p(\theta|\alpha)$ and likelihood on y given by the (modeled, parameterized) response $p(y|a, x, \theta)$.
 - ▶ (now you're not only generative, you're Bayesian.)
 - ▶ $p(\theta|D) = p(\theta|\{y\}_1^t, \{a\}_1^t, \{x\}_1^t, \alpha)$
 - ▶ $\propto p(\{y\}_1^t|\{a\}_1^t, \{x\}_1^t, \theta)p(\theta|\alpha)$
 - ▶ $= p(\theta|\alpha)\prod_t p(y_t|a_t, x_t, \theta)$
 - ▶ *warning 1*: sometimes people write “ $p(D|\theta)$ ” but we don't need $p(a|\theta)$ or $p(x|\theta)$ here
 - ▶ *warning 2*: don't need historical record of θ_t .
 - ▶ (we used Bayes rule, but only in θ and y .)
- ▶ A2: evaluate integral by $N = 1$ Monte Carlo

TS: connecting w/“generative causal modeling” 2

- ▶ Q1: what's $p(\theta|D)$?
- ▶ Q2: how am I going to evaluate this integral?
- ▶ A1: $p(\theta|D)$ definable by choosing prior $p(\theta|\alpha)$ and likelihood on y given by the (modeled, parameterized) response $p(y|a, x, \theta)$.
 - ▶ (now you're not only generative, you're Bayesian.)
 - ▶ $p(\theta|D) = p(\theta|\{y\}_1^t, \{a\}_1^t, \{x\}_1^t, \alpha)$
 - ▶ $\propto p(\{y\}_1^t|\{a\}_1^t, \{x\}_1^t, \theta)p(\theta|\alpha)$
 - ▶ $= p(\theta|\alpha)\prod_t p(y_t|a_t, x_t, \theta)$
 - ▶ *warning 1*: sometimes people write “ $p(D|\theta)$ ” but we don't need $p(a|\theta)$ or $p(x|\theta)$ here
 - ▶ *warning 2*: don't need historical record of θ_t .
 - ▶ (we used Bayes rule, but only in θ and y .)
- ▶ A2: evaluate integral by $N = 1$ Monte Carlo
 - ▶ take 1 sample “ θ_t ” of θ from $p(\theta|D)$

TS: connecting w/“generative causal modeling” 2

- ▶ Q1: what's $p(\theta|D)$?
- ▶ Q2: how am I going to evaluate this integral?
- ▶ A1: $p(\theta|D)$ definable by choosing prior $p(\theta|\alpha)$ and likelihood on y given by the (modeled, parameterized) response $p(y|a, x, \theta)$.
 - ▶ (now you're not only generative, you're Bayesian.)
 - ▶ $p(\theta|D) = p(\theta|\{y\}_1^t, \{a\}_1^t, \{x\}_1^t, \alpha)$
 - ▶ $\propto p(\{y\}_1^t|\{a\}_1^t, \{x\}_1^t, \theta)p(\theta|\alpha)$
 - ▶ $= p(\theta|\alpha)\prod_t p(y_t|a_t, x_t, \theta)$
 - ▶ *warning 1*: sometimes people write “ $p(D|\theta)$ ” but we don't need $p(a|\theta)$ or $p(x|\theta)$ here
 - ▶ *warning 2*: don't need historical record of θ_t .
 - ▶ (we used Bayes rule, but only in θ and y .)
- ▶ A2: evaluate integral by $N = 1$ Monte Carlo
 - ▶ take 1 sample “ θ_t ” of θ from $p(\theta|D)$
 - ▶ $a_t = h(x_t; \theta_t) = \operatorname{argmax}_a Q(a, x, \theta_t)$

That sounds hard.

No, just general. Let's do toy case:

- ▶ $y \in \{0, 1\}$,

That sounds hard.

No, just general. Let's do toy case:

- ▶ $y \in \{0, 1\}$,
- ▶ no context x ,

That sounds hard.

No, just general. Let's do toy case:

- ▶ $y \in \{0, 1\}$,
- ▶ no context x ,
- ▶ Bernoulli (coin flipping), keep track of

That sounds hard.

No, just general. Let's do toy case:

- ▶ $y \in \{0, 1\}$,
- ▶ no context x ,
- ▶ Bernoulli (coin flipping), keep track of
 - ▶ $S_a \equiv$ number of successes flipping coin a

That sounds hard.

No, just general. Let's do toy case:

- ▶ $y \in \{0, 1\}$,
- ▶ no context x ,
- ▶ Bernoulli (coin flipping), keep track of
 - ▶ $S_a \equiv$ number of successes flipping coin a
 - ▶ $F_a \equiv$ number of failures flipping coin a

That sounds hard.

No, just general. Let's do toy case:

- ▶ $y \in \{0, 1\}$,
- ▶ no context x ,
- ▶ Bernoulli (coin flipping), keep track of
 - ▶ $S_a \equiv$ number of successes flipping coin a
 - ▶ $F_a \equiv$ number of failures flipping coin a

That sounds hard.

No, just general. Let's do toy case:

- ▶ $y \in \{0, 1\}$,
- ▶ no context x ,
- ▶ Bernoulli (coin flipping), keep track of
 - ▶ $S_a \equiv$ number of successes flipping coin a
 - ▶ $F_a \equiv$ number of failures flipping coin a

Then

- ▶ $p(\theta|D) \propto p(\theta|\alpha) \prod_t p(y_t|a_t, \theta)$

That sounds hard.

No, just general. Let's do toy case:

- ▶ $y \in \{0, 1\}$,
- ▶ no context x ,
- ▶ Bernoulli (coin flipping), keep track of
 - ▶ $S_a \equiv$ number of successes flipping coin a
 - ▶ $F_a \equiv$ number of failures flipping coin a

Then

- ▶ $p(\theta|D) \propto p(\theta|\alpha) \prod_t p(y_t|a_t, \theta)$
- ▶ $= \left(\prod_a \theta_a^{\alpha-1} (1 - \theta_a)^{\beta-1} \right) \left(\prod_{t,a_t} \theta_{a_t}^{y_t} (1 - \theta_{a_t})^{1-y_t} \right)$

That sounds hard.

No, just general. Let's do toy case:

- ▶ $y \in \{0, 1\}$,
- ▶ no context x ,
- ▶ Bernoulli (coin flipping), keep track of
 - ▶ $S_a \equiv$ number of successes flipping coin a
 - ▶ $F_a \equiv$ number of failures flipping coin a

Then

- ▶ $p(\theta|D) \propto p(\theta|\alpha) \prod_t p(y_t|a_t, \theta)$
- ▶ $= \left(\prod_a \theta_a^{\alpha-1} (1 - \theta_a)^{\beta-1} \right) \left(\prod_{t,a_t} \theta_{a_t}^{y_t} (1 - \theta_{a_t})^{1-y_t} \right)$
- ▶ $= \prod_a \theta_a^{\alpha+S_a-1} (1 - \theta_a)^{\beta+F_a-1}$

That sounds hard.

No, just general. Let's do toy case:

- ▶ $y \in \{0, 1\}$,
- ▶ no context x ,
- ▶ Bernoulli (coin flipping), keep track of
 - ▶ $S_a \equiv$ number of successes flipping coin a
 - ▶ $F_a \equiv$ number of failures flipping coin a

Then

- ▶ $p(\theta|D) \propto p(\theta|\alpha) \prod_t p(y_t|a_t, \theta)$
- ▶ $= \left(\prod_a \theta_a^{\alpha-1} (1 - \theta_a)^{\beta-1} \right) \left(\prod_{t,a_t} \theta_{a_t}^{y_t} (1 - \theta_{a_t})^{1-y_t} \right)$
- ▶ $= \prod_a \theta_a^{\alpha+S_a-1} (1 - \theta_a)^{\beta+F_a-1}$
- ▶ $\therefore \theta_a \sim \text{Beta}(\alpha + S_a, \beta + F_a)$

Thompson sampling: results (2011)

An Empirical Evaluation of Thompson Sampling

Olivier Chapelle
Yahoo! Research
Santa Clara, CA
chap@yahoo-inc.com

Lihong Li
Yahoo! Research
Santa Clara, CA
lihong@yahoo-inc.com

Figure 15: Chaleppe and Li 2011

TS: words

In the realizable case, the reward is a stochastic function of the action, context and the unknown, true parameter θ^* . Ideally, we would like to choose the action maximizing the expected reward, $\max_a \mathbb{E}(r|a, x, \theta^*)$.

Of course, θ^* is unknown. If we are just interested in maximizing the immediate reward (exploitation), then one should choose the action that maximizes $\mathbb{E}(r|a, x) = \int \mathbb{E}(r|a, x, \theta) P(\theta|D) d\theta$.

But in an exploration / exploitation setting, the probability matching heuristic consists in randomly selecting an action a according to its probability of being optimal. That is, action a is chosen with probability

$$\int \mathbb{I} \left[\mathbb{E}(r|a, x, \theta) = \max_{a'} \mathbb{E}(r|a', x, \theta) \right] P(\theta|D) d\theta,$$

where \mathbb{I} is the indicator function. Note that the integral does not have to be computed explicitly: it suffices to draw a random parameter θ at each round as explained in Algorithm 1. Implementation of the algorithm is thus efficient and straightforward in most applications.

Figure 16: from Chaleppe and Li 2011

Algorithm 1 Thompson sampling

```
 $D = \emptyset$   
for  $t = 1, \dots, T$  do  
    Receive context  $x_t$   
    Draw  $\theta^t$  according to  $P(\theta|D)$   
    Select  $a_t = \arg \max_a \mathbb{E}_r(r|x_t, a, \theta^t)$   
    Observe reward  $r_t$   
     $D = D \cup (x_t, a_t, r_t)$   
end for
```

Algorithm 2 Thompson sampling for the Bernoulli bandit

Require: α, β prior parameters of a Beta distribution
 $S_i = 0, F_i = 0, \forall i$. {Success and failure counters}
for $t = 1, \dots, T$ **do**
 for $i = 1, \dots, K$ **do**
 Draw θ_i according to $\text{Beta}(S_i + \alpha, F_i + \beta)$.
 end for
 Draw arm $\hat{i} = \arg \max_i \theta_i$ and observe reward r
 if $r = 1$ **then**
 $S_{\hat{i}} = S_{\hat{i}} + 1$
 else
 $F_{\hat{i}} = F_{\hat{i}} + 1$
 end if
end for

TS: Bernoulli bandit p-code (results)




Figure 19: from Chaleppe and Li 2011

UCB1 (2002), p-code

Deterministic policy: UCB1.

Initialization: Play each machine once.

Loop:

- Play machine j that maximizes $\bar{x}_j + \sqrt{\frac{2 \ln n}{n_j}}$, where \bar{x}_j is the average reward obtained from machine j , n_j is the number of times machine j has been played so far, and n is the overall number of plays done so far.

Figure 20: UCB1

from Auer, Peter, Nicolo Cesa-Bianchi, and Paul Fischer.

“Finite-time analysis of the multiarmed bandit problem.” Machine learning 47.2-3 (2002): 235-256.

TS: with context

Algorithm 3 Regularized logistic regression with batch updates

Require: Regularization parameter $\lambda > 0$.

$m_i = 0, q_i = \lambda$. {Each weight w_i has an independent prior $\mathcal{N}(m_i, q_i^{-1})$ }

for $t = 1, \dots, T$ **do**

 Get a new batch of training data $(\mathbf{x}_j, y_j), j = 1, \dots, n$.

 Find \mathbf{w} as the minimizer of: $\frac{1}{2} \sum_{i=1}^d q_i (w_i - m_i)^2 + \sum_{j=1}^n \log(1 + \exp(-y_j \mathbf{w}^\top \mathbf{x}_j))$.

$m_i = w_i$

$q_i = q_i + \sum_{j=1}^n x_{ij}^2 p_j (1 - p_j), p_j = (1 + \exp(-\mathbf{w}^\top \mathbf{x}_j))^{-1}$ {Laplace approximation}

end for

Figure 21: from Chaleppe and Li 2011

LinUCB: UCB with context

Algorithm 1 LinUCB with disjoint linear models.

```
0: Inputs:  $\alpha \in \mathbb{R}_+$ 
1: for  $t = 1, 2, 3, \dots, T$  do
2:   Observe features of all arms  $a \in \mathcal{A}_t$ :  $\mathbf{x}_{t,a} \in \mathbb{R}^d$ 
3:   for all  $a \in \mathcal{A}_t$  do
4:     if  $a$  is new then
5:        $\mathbf{A}_a \leftarrow \mathbf{I}_d$  ( $d$ -dimensional identity matrix)
6:        $\mathbf{b}_a \leftarrow \mathbf{0}_{d \times 1}$  ( $d$ -dimensional zero vector)
7:     end if
8:      $\hat{\boldsymbol{\theta}}_a \leftarrow \mathbf{A}_a^{-1} \mathbf{b}_a$ 
9:      $p_{t,a} \leftarrow \hat{\boldsymbol{\theta}}_a^\top \mathbf{x}_{t,a} + \alpha \sqrt{\mathbf{x}_{t,a}^\top \mathbf{A}_a^{-1} \mathbf{x}_{t,a}}$ 
10:   end for
11:   Choose arm  $a_t = \arg \max_{a \in \mathcal{A}_t} p_{t,a}$  with ties broken arbitrarily, and observe a real-valued payoff  $r_t$ 
12:    $\mathbf{A}_{a_t} \leftarrow \mathbf{A}_{a_t} + \mathbf{x}_{t,a_t} \mathbf{x}_{t,a_t}^\top$ 
13:    $\mathbf{b}_{a_t} \leftarrow \mathbf{b}_{a_t} + r_t \mathbf{x}_{t,a_t}$ 
14: end for
```

Figure 22: LinUCB

TS: with context (results)

Table 2: CTR regrets on the display advertising data.

Method	TS			LinUCB			ϵ -greedy			Exploit	Randor
Parameter	0.25	0.5	1	0.5	1	2	0.005	0.01	0.02		
Regret (%)	4.45	3.72	3.81	4.99	4.22	4.14	5.05	4.98	5.22	5.00	31.95




Figure 4: CTR regret over the 4 days test period for 3 algorithms: Thompson sampling with $\alpha = 0.5$, LinUCB with $\alpha = 2$, Exploit-only. The regret in the first hour is large, around 0.3, because the algorithms predict randomly (no initial model provided).

Bandits: Regret via Lai and Robbins (1985)

THEOREM 2. Assume that $I(\theta, \lambda)$ satisfies (1.6) and (1.7) and that Θ satisfies (1.9). Fix $j \in \{1, \dots, k\}$, and define Θ_j and Θ_j^* by (2.1). Let φ be any rule such that for every $\theta \in \Theta_j^*$, as $n \rightarrow \infty$

$$\sum_{i \neq j} E_{\theta} T_n(i) = o(n^a) \quad \text{for every } a > 0, \quad (2.2)$$

where $T_n(i)$, defined in (1.2), is the number of times that the rule φ samples from Π_i up to stage n . Then for every $\theta \in \Theta_j$ and every $\epsilon > 0$,

$$\lim_{n \rightarrow \infty} P_{\theta} \{ T_n(j) \geq (1 - \epsilon)(\log n)/I(\theta_j, \theta^*) \} = 1, \quad (2.3)$$

where θ^* is defined in (1.4), and hence

$$\liminf_{n \rightarrow \infty} E_{\theta} T_n(j) / \log n \geq 1/I(\theta_j, \theta^*).$$

Figure 24: Lai Robbins

Thompson sampling (1933) and optimality (2013)

Theorem 2. For any instance $\Theta = \{\mu_1, \dots, \mu_N\}$ of Bernoulli MAB,

$$R(T, \Theta) \leq (1 + \epsilon) \sum_{i \neq I^*} \frac{\ln(T) \Delta_i}{KL(\mu_i, \mu^*)} + O(N/\epsilon^2)$$

Recall that we have $\lim_{T \rightarrow \infty} \frac{R(T, \Theta)}{\ln(T)} \geq \sum_{i \neq I^*} \frac{\Delta_i}{KL(\mu_i, \mu^*)}$. Above theorem says that Thompson Sampling matches this lower bound. We also have the following problem independent regret bound for this algorithm.

Theorem 3. For all Θ ,

$$R(T) = \max_{\Theta} R(T, \Theta) \leq O(\sqrt{NT \log T} + N)$$

For proofs of above theorems, refer to [2].

Figure 25: TS result

from S. Agrawal, N. Goyal, "Further optimal regret bounds for Thompson Sampling", AISTATS 2013.; see also Agrawal, Shipra, and Navin Goyal. "Analysis of Thompson Sampling for the Multi-armed Bandit Problem." COLT. 2012 and Emilie Kaufmann, Nathaniel Korda, and R´emi Munos. Thompson sampling: An asymptotically optimal finite-time analysis. In Algorithmic Learning Theory, pages 199–213. Springer, 2012.

other 'Causalities': structure learning




Figure 26: (a) Causal graphs showing for explanations for an observed dependence between *Ad* and *Buy*. The node *H* corresponds to a hidden common cause of *Ad* and *Buy*. The shaded node *S* indicates that the case has been included in the database. (b) A Bayesian network for which *A* causes *B* is the only causal explanation, given the causal Markov condition.

Figure 26: from heckerman 1995

D. Heckerman. A Tutorial on Learning with Bayesian Networks.
Technical Report MSR-TR-95-06, Microsoft Research, March, 1995.

other 'Causalities': potential outcomes

- ▶ model distribution of $p(y_i(1), y_i(0), a_i, x_i)$

other 'Causalities': potential outcomes

- ▶ model distribution of $p(y_i(1), y_i(0), a_i, x_i)$
- ▶ “action” replaced by “observed outcome”

other 'Causalities': potential outcomes

- ▶ model distribution of $p(y_i(1), y_i(0), a_i, x_i)$
- ▶ “action” replaced by “observed outcome”
- ▶ aka Neyman-Rubin causal model: Neyman ('23); Rubin ('74)

other ‘Causalities’: potential outcomes

- ▶ model distribution of $p(y_i(1), y_i(0), a_i, x_i)$
- ▶ “action” replaced by “observed outcome”
- ▶ aka Neyman-Rubin causal model: Neyman ('23); Rubin ('74)
- ▶ see Morgan + Winship²³ for connections between frameworks

²³Morgan, Stephen L., and Christopher Winship. *Counterfactuals and causal inference* Cambridge University Press, 2014.

Lecture 4: descriptive modeling @ NYT

review: (latent) inference and clustering

- ▶ what does kmeans mean?

review: (latent) inference and clustering

- ▶ what does kmeans mean?
 - ▶ given $x_i \in R^D$

review: (latent) inference and clustering

- ▶ what does kmeans mean?
 - ▶ given $x_i \in R^D$
 - ▶ given $d : R^D \rightarrow R^1$

review: (latent) inference and clustering

- ▶ what does kmeans mean?
 - ▶ given $x_i \in R^D$
 - ▶ given $d : R^D \rightarrow R^1$
 - ▶ assign z_i

review: (latent) inference and clustering

- ▶ what does kmeans mean?
 - ▶ given $x_i \in R^D$
 - ▶ given $d : R^D \rightarrow R^1$
 - ▶ assign z_i
- ▶ generative modeling gives meaning

review: (latent) inference and clustering

- ▶ what does kmeans mean?
 - ▶ given $x_i \in R^D$
 - ▶ given $d : R^D \rightarrow R^1$
 - ▶ assign z_i
- ▶ generative modeling gives meaning
 - ▶ given $p(x|z, \theta)$

review: (latent) inference and clustering

- ▶ what does kmeans mean?
 - ▶ given $x_i \in R^D$
 - ▶ given $d : R^D \rightarrow R^1$
 - ▶ assign z_i
- ▶ generative modeling gives meaning
 - ▶ given $p(x|z, \theta)$
 - ▶ maximize $p(x|\theta)$

review: (latent) inference and clustering

- ▶ what does kmeans mean?
 - ▶ given $x_i \in R^D$
 - ▶ given $d : R^D \rightarrow R^1$
 - ▶ assign z_i
- ▶ generative modeling gives meaning
 - ▶ given $p(x|z, \theta)$
 - ▶ maximize $p(x|\theta)$
 - ▶ output assignment $p(z|x, \theta)$

actual math

- ▶ define $P \equiv p(x, z|\theta)$

actual math

- ▶ define $P \equiv p(x, z|\theta)$
- ▶ log-likelihood $L \equiv \log p(x|\theta) = \log \sum_z P = \log E_q P / q$
(cf. importance sampling)

actual math

- ▶ define $P \equiv p(x, z|\theta)$
- ▶ log-likelihood $L \equiv \log p(x|\theta) = \log \sum_z P = \log E_q P/q$
(cf. importance sampling)
- ▶ Jensen's:
$$L \geq \tilde{L} \equiv E_q \log P/q = E_q \log P + H[q] = -(U - H) = -\mathcal{F}$$

actual math

- ▶ define $P \equiv p(x, z|\theta)$
- ▶ log-likelihood $L \equiv \log p(x|\theta) = \log \sum_z P = \log E_q P/q$
(cf. importance sampling)
- ▶ Jensen's:
$$L \geq \tilde{L} \equiv E_q \log P/q = E_q \log P + H[q] = -(U - H) = -\mathcal{F}$$
 - ▶ analogy to free energy in physics

actual math

- ▶ define $P \equiv p(x, z|\theta)$
- ▶ log-likelihood $L \equiv \log p(x|\theta) = \log \sum_z P = \log E_q P/q$
(cf. importance sampling)
- ▶ Jensen's:
$$L \geq \tilde{L} \equiv E_q \log P/q = E_q \log P + H[q] = -(U - H) = -\mathcal{F}$$
 - ▶ analogy to free energy in physics
- ▶ alternate optimization on θ and on q

actual math

- ▶ define $P \equiv p(x, z|\theta)$
- ▶ log-likelihood $L \equiv \log p(x|\theta) = \log \sum_z P = \log E_q P/q$
(cf. importance sampling)
- ▶ Jensen's:
$$L \geq \tilde{L} \equiv E_q \log P/q = E_q \log P + H[q] = -(U - H) = -\mathcal{F}$$
 - ▶ analogy to free energy in physics
- ▶ alternate optimization on θ and on q
 - ▶ NB: q step gives $q(z) = p(z|x, \theta)$

actual math

- ▶ define $P \equiv p(x, z|\theta)$
- ▶ log-likelihood $L \equiv \log p(x|\theta) = \log \sum_z P = \log E_q P/q$
(cf. importance sampling)
- ▶ Jensen's:
$$L \geq \tilde{L} \equiv E_q \log P/q = E_q \log P + H[q] = -(U - H) = -\mathcal{F}$$
 - ▶ analogy to free energy in physics
- ▶ alternate optimization on θ and on q
 - ▶ NB: q step gives $q(z) = p(z|x, \theta)$
 - ▶ NB: $\log P$ convenient for independent examples w/ exponential families

actual math

- ▶ define $P \equiv p(x, z|\theta)$
- ▶ log-likelihood $L \equiv \log p(x|\theta) = \log \sum_z P = \log E_q P/q$
(cf. importance sampling)
- ▶ Jensen's:
$$L \geq \tilde{L} \equiv E_q \log P/q = E_q \log P + H[q] = -(U - H) = -\mathcal{F}$$
 - ▶ analogy to free energy in physics
- ▶ alternate optimization on θ and on q
 - ▶ NB: q step gives $q(z) = p(z|x, \theta)$
 - ▶ NB: $\log P$ convenient for independent examples w/ exponential families
 - ▶ e.g., GMMs: $\mu_k \leftarrow E[x|z]$ and $\sigma_k^2 \leftarrow E[(x - \mu)^2|z]$ are sufficient statistics

actual math

- ▶ define $P \equiv p(x, z|\theta)$
- ▶ log-likelihood $L \equiv \log p(x|\theta) = \log \sum_z P = \log E_q P/q$
(cf. importance sampling)
- ▶ Jensen's:
$$L \geq \tilde{L} \equiv E_q \log P/q = E_q \log P + H[q] = -(U - H) = -\mathcal{F}$$
 - ▶ analogy to free energy in physics
- ▶ alternate optimization on θ and on q
 - ▶ NB: q step gives $q(z) = p(z|x, \theta)$
 - ▶ NB: $\log P$ convenient for independent examples w/ exponential families
 - ▶ e.g., GMMs: $\mu_k \leftarrow E[x|z]$ and $\sigma_k^2 \leftarrow E[(x - \mu)^2|z]$ are sufficient statistics
 - ▶ e.g., LDAs: word counts are sufficient statistics

tangent: more math on GMMs, part 1

Energy U (to be minimized):

$$\blacktriangleright -U \equiv E_q \log P = \sum_z \sum_i q_i(z) \log P(x_i, z_i) \equiv U_x + U_z$$

tangent: more math on GMMs, part 1

Energy U (to be minimized):

- ▶ $-U \equiv E_q \log P = \sum_z \sum_i q_i(z) \log P(x_i, z_i) \equiv U_x + U_z$
- ▶ $-U_x \equiv \sum_z \sum_i q_i(z) \log p(x_i|z_i)$

tangent: more math on GMMs, part 1

Energy U (to be minimized):

- ▶ $-U \equiv E_q \log P = \sum_z \sum_i q_i(z) \log P(x_i, z_i) \equiv U_x + U_z$
 - ▶ $-U_x \equiv \sum_z \sum_i q_i(z) \log p(x_i|z_i)$
 - ▶ $= \sum_i \sum_z q_i(z) \sum_k 1[z_i = k] \log p(x_i|z_i)$
-

tangent: more math on GMMs, part 1

Energy U (to be minimized):

- ▶ $-U \equiv E_q \log P = \sum_z \sum_i q_i(z) \log P(x_i, z_i) \equiv U_x + U_z$
 - ▶ $-U_x \equiv \sum_z \sum_i q_i(z) \log p(x_i|z_i)$
 - ▶ $= \sum_i \sum_z q_i(z) \sum_k 1[z_i = k] \log p(x_i|z_i)$
 - ▶ define $r_{ik} = \sum_z q_i(z) 1[z_i = k]$
-

tangent: more math on GMMs, part 1

Energy U (to be minimized):

- ▶ $-U \equiv E_q \log P = \sum_z \sum_i q_i(z) \log P(x_i, z_i) \equiv U_x + U_z$
 - ▶ $-U_x \equiv \sum_z \sum_i q_i(z) \log p(x_i|z_i)$
 - ▶ $= \sum_i \sum_z q_i(z) \sum_k 1[z_i = k] \log p(x_i|z_i)$
 - ▶ define $r_{ik} = \sum_z q_i(z) 1[z_i = k]$
 - ▶ $-U_x = \sum_i r_{ik} \log p(x_i|k)$.
-

tangent: more math on GMMs, part 1

Energy U (to be minimized):

- ▶ $-U \equiv E_q \log P = \sum_z \sum_i q_i(z) \log P(x_i, z_i) \equiv U_x + U_z$
- ▶ $-U_x \equiv \sum_z \sum_i q_i(z) \log p(x_i|z_i)$
- ▶ $= \sum_i \sum_z q_i(z) \sum_k 1[z_i = k] \log p(x_i|z_i)$
- ▶ define $r_{ik} = \sum_z q_i(z) 1[z_i = k]$
- ▶ $-U_x = \sum_i r_{ik} \log p(x_i|k)$.
- ▶ Gaussian²⁴
 $\Rightarrow -U_x = \sum_i r_{ik} \left(-\frac{1}{2}(x_i - \mu_k)^2 \lambda_k + \frac{1}{2} \ln \lambda_k - \frac{1}{2} \ln 2\pi \right)$

²⁴math is simpler if you work with $\lambda_k \equiv \sigma^{-2}$

tangent: more math on GMMs, part 1

Energy U (to be minimized):

- ▶ $-U \equiv E_q \log P = \sum_z \sum_i q_i(z) \log P(x_i, z_i) \equiv U_x + U_z$
- ▶ $-U_x \equiv \sum_z \sum_i q_i(z) \log p(x_i|z_i)$
- ▶ $= \sum_i \sum_z q_i(z) \sum_k 1[z_i = k] \log p(x_i|z_i)$
- ▶ define $r_{ik} = \sum_z q_i(z) 1[z_i = k]$
- ▶ $-U_x = \sum_i r_{ik} \log p(x_i|k)$.
- ▶ Gaussian²⁴
 $\Rightarrow -U_x = \sum_i r_{ik} \left(-\frac{1}{2}(x_i - \mu_k)^2 \lambda_k + \frac{1}{2} \ln \lambda_k - \frac{1}{2} \ln 2\pi \right)$

²⁴math is simpler if you work with $\lambda_k \equiv \sigma^{-2}$

tangent: more math on GMMs, part 1

Energy U (to be minimized):

- ▶ $-U \equiv E_q \log P = \sum_z \sum_i q_i(z) \log P(x_i, z_i) \equiv U_x + U_z$
- ▶ $-U_x \equiv \sum_z \sum_i q_i(z) \log p(x_i|z_i)$
- ▶ $= \sum_i \sum_z q_i(z) \sum_k 1[z_i = k] \log p(x_i|z_i)$
- ▶ define $r_{ik} = \sum_z q_i(z) 1[z_i = k]$
- ▶ $-U_x = \sum_i r_{ik} \log p(x_i|k)$.
- ▶ Gaussian²⁴
 $\Rightarrow -U_x = \sum_i r_{ik} \left(-\frac{1}{2}(x_i - \mu_k)^2 \lambda_k + \frac{1}{2} \ln \lambda_k - \frac{1}{2} \ln 2\pi \right)$

simple to minimize for parameters $\vartheta = \{\mu_k, \lambda_k\}$

²⁴math is simpler if you work with $\lambda_k \equiv \sigma^{-2}$

tangent: more math on GMMs, part 2

$$\blacktriangleright -U_x = \sum_i r_{ik} \left(-\frac{1}{2}(x_i - \mu_k)^2 \lambda_k + \frac{1}{2} \ln \lambda_k - \frac{1}{2} \ln 2\pi \right)$$

tangent: more math on GMMs, part 2

- ▶ $-U_x = \sum_i r_{ik} \left(-\frac{1}{2}(x_i - \mu_k)^2 \lambda_k + \frac{1}{2} \ln \lambda_k - \frac{1}{2} \ln 2\pi \right)$
- ▶ $\mu_k \leftarrow E[x|k]$ solves $\sum_i r_{ik} = \sum_i r_{ik} x_i$

tangent: more math on GMMs, part 2

- ▶ $-U_x = \sum_i r_{ik} \left(-\frac{1}{2}(x_i - \mu_k)^2 \lambda_k + \frac{1}{2} \ln \lambda_k - \frac{1}{2} \ln 2\pi \right)$
- ▶ $\mu_k \leftarrow E[x|k]$ solves $\sum_i r_{ik} = \sum_i r_{ik} x_i$
- ▶ $\lambda_k \leftarrow E[(x - \mu)^2|k]$ solves $\sum_i r_{ik} \frac{1}{2} (x_i - \mu_k)^2 = \lambda_k^{-1} \sum_i r_{ik}$

tangent: Gaussians \in exponential family²⁶

- ▶ as before, $-U = \sum_i r_{ik} \log p(x_i|k)$

tangent: Gaussians \in exponential family²⁶

- ▶ as before, $-U = \sum_i r_{ik} \log p(x_i|k)$
- ▶ define $p(x_i|k) = \exp(\eta(\theta) \cdot T(x) - A(\theta) + B(x))$

tangent: Gaussians \in exponential family²⁶

- ▶ as before, $-U = \sum_i r_{ik} \log p(x_i|k)$
- ▶ define $p(x_i|k) = \exp(\eta(\theta) \cdot T(x) - A(\theta) + B(x))$
- ▶ e.g., Gaussian case ²⁵,

²⁵Choosing $\eta(\theta) = \eta$ called 'canonical form'

tangent: Gaussians \in exponential family²⁶

- ▶ as before, $-U = \sum_i r_{ik} \log p(x_i|k)$
- ▶ define $p(x_i|k) = \exp(\eta(\theta) \cdot T(x) - A(\theta) + B(x))$
- ▶ e.g., Gaussian case ²⁵,
 - ▶ $T_1 = x$,

²⁵Choosing $\eta(\theta) = \eta$ called 'canonical form'

tangent: Gaussians \in exponential family²⁶

- ▶ as before, $-U = \sum_i r_{ik} \log p(x_i|k)$
- ▶ define $p(x_i|k) = \exp(\eta(\theta) \cdot T(x) - A(\theta) + B(x))$
- ▶ e.g., Gaussian case ²⁵,
 - ▶ $T_1 = x$,
 - ▶ $T_2 = x^2$

²⁵Choosing $\eta(\theta) = \eta$ called 'canonical form'

tangent: Gaussians \in exponential family²⁶

- ▶ as before, $-U = \sum_i r_{ik} \log p(x_i|k)$
- ▶ define $p(x_i|k) = \exp(\eta(\theta) \cdot T(x) - A(\theta) + B(x))$
- ▶ e.g., Gaussian case ²⁵,
 - ▶ $T_1 = x$,
 - ▶ $T_2 = x^2$
 - ▶ $\eta_1 = \mu/\sigma^2 = \mu\lambda$

²⁵Choosing $\eta(\theta) = \eta$ called 'canonical form'

tangent: Gaussians \in exponential family²⁶

- ▶ as before, $-U = \sum_i r_{ik} \log p(x_i|k)$
- ▶ define $p(x_i|k) = \exp(\eta(\theta) \cdot T(x) - A(\theta) + B(x))$
- ▶ e.g., Gaussian case ²⁵,
 - ▶ $T_1 = x$,
 - ▶ $T_2 = x^2$
 - ▶ $\eta_1 = \mu/\sigma^2 = \mu\lambda$
 - ▶ $\eta_2 = -\frac{1}{2}\lambda = -1/(2\sigma^2)$

²⁵Choosing $\eta(\theta) = \eta$ called 'canonical form'

tangent: Gaussians \in exponential family²⁶

- ▶ as before, $-U = \sum_i r_{ik} \log p(x_i|k)$
- ▶ define $p(x_i|k) = \exp(\eta(\theta) \cdot T(x) - A(\theta) + B(x))$
- ▶ e.g., Gaussian case ²⁵,
 - ▶ $T_1 = x$,
 - ▶ $T_2 = x^2$
 - ▶ $\eta_1 = \mu/\sigma^2 = \mu\lambda$
 - ▶ $\eta_2 = -\frac{1}{2}\lambda = -1/(2\sigma^2)$
 - ▶ $A = \lambda\mu^2/2 - \frac{1}{2} \ln \lambda$

²⁵Choosing $\eta(\theta) = \eta$ called 'canonical form'

tangent: Gaussians \in exponential family²⁶

- ▶ as before, $-U = \sum_i r_{ik} \log p(x_i|k)$
- ▶ define $p(x_i|k) = \exp(\eta(\theta) \cdot T(x) - A(\theta) + B(x))$
- ▶ e.g., Gaussian case ²⁵,
 - ▶ $T_1 = x$,
 - ▶ $T_2 = x^2$
 - ▶ $\eta_1 = \mu/\sigma^2 = \mu\lambda$
 - ▶ $\eta_2 = -\frac{1}{2}\lambda = -1/(2\sigma^2)$
 - ▶ $A = \lambda\mu^2/2 - \frac{1}{2} \ln \lambda$
 - ▶ $\exp(B(x)) = (2\pi)^{-1/2}$

²⁵Choosing $\eta(\theta) = \eta$ called 'canonical form'

tangent: Gaussians \in exponential family²⁶

- ▶ as before, $-U = \sum_i r_{ik} \log p(x_i|k)$
- ▶ define $p(x_i|k) = \exp(\eta(\theta) \cdot T(x) - A(\theta) + B(x))$
- ▶ e.g., Gaussian case ²⁵,
 - ▶ $T_1 = x$,
 - ▶ $T_2 = x^2$
 - ▶ $\eta_1 = \mu/\sigma^2 = \mu\lambda$
 - ▶ $\eta_2 = -\frac{1}{2}\lambda = -1/(2\sigma^2)$
 - ▶ $A = \lambda\mu^2/2 - \frac{1}{2} \ln \lambda$
 - ▶ $\exp(B(x)) = (2\pi)^{-1/2}$
- ▶ note that in a mixture model, there are separate η (and thus $A(\eta)$) for each value of z

²⁵Choosing $\eta(\theta) = \eta$ called 'canonical form'

²⁶NB: Gaussians \in exponential family, GMM \notin exponential family! (Thanks to Eszter Vértés for pointing out this error in earlier title.)

tangent: variational joy \in exponential family

- ▶ as before, $-U = \sum_i r_{ik} \left(\eta_k^T T(x_i) - A(\eta_k) + B(x_i) \right)$

tangent: variational joy \in exponential family

- ▶ as before, $-U = \sum_i r_{ik} \left(\eta_k^T T(x_i) - A(\eta_k) + B(x_i) \right)$
- ▶ $\eta_{k,\alpha}$ solves $\sum_i r_{ik} T_{k,\alpha}(x_i) = \frac{\partial A(\eta_k)}{\partial \eta_{k,\alpha}} \sum_i r_{ik}$ (canonical)

tangent: variational joy \in exponential family

- ▶ as before, $-U = \sum_i r_{ik} \left(\eta_k^T T(x_i) - A(\eta_k) + B(x_i) \right)$
 - ▶ $\eta_{k,\alpha}$ solves $\sum_i r_{ik} T_{k,\alpha}(x_i) = \frac{\partial A(\eta_k)}{\partial \eta_{k,\alpha}} \sum_i r_{ik}$ (canonical)
 - ▶ $\therefore \partial_{\eta_{k,\alpha}} A(\eta_k) \leftarrow E[T_{k,\alpha}|k]$ (canonical)
-

tangent: variational joy \in exponential family

- ▶ as before, $-U = \sum_i r_{ik} \left(\eta_k^T T(x_i) - A(\eta_k) + B(x_i) \right)$
- ▶ $\eta_{k,\alpha}$ solves $\sum_i r_{ik} T_{k,\alpha}(x_i) = \frac{\partial A(\eta_k)}{\partial \eta_{k,\alpha}} \sum_i r_{ik}$ (canonical)
- ▶ $\therefore \partial_{\eta_{k,\alpha}} A(\eta_k) \leftarrow E[T_{k,\alpha}|k]$ (canonical)
- ▶ nice connection w/physics, esp. mean field theory²⁷

²⁷read MacKay, David JC. *Information theory, inference and learning algorithms*, Cambridge university press, 2003 to learn more. Actually you should read it regardless.

clustering and inference: GMM/k-means case study

- ▶ generative model gives meaning and optimization

clustering and inference: GMM/k-means case study

- ▶ generative model gives meaning and optimization
- ▶ large freedom to choose different optimization approaches

clustering and inference: GMM/k-means case study

- ▶ generative model gives meaning and optimization
- ▶ large freedom to choose different optimization approaches
 - ▶ e.g., hard clustering limit

clustering and inference: GMM/k-means case study

- ▶ generative model gives meaning and optimization
- ▶ large freedom to choose different optimization approaches
 - ▶ e.g., hard clustering limit
 - ▶ e.g., streaming solutions

clustering and inference: GMM/k-means case study

- ▶ generative model gives meaning and optimization
- ▶ large freedom to choose different optimization approaches
 - ▶ e.g., hard clustering limit
 - ▶ e.g., streaming solutions
 - ▶ e.g., stochastic gradient methods

general framework: E+M/variational

- ▶ e.g., GMM+hard clustering gives kmeans

general framework: E+M/variational

- ▶ e.g., GMM+hard clustering gives kmeans
- ▶ e.g., some favorite applications:

general framework: E+M/variational

- ▶ e.g., GMM+hard clustering gives kmeans
- ▶ e.g., some favorite applications:
 - ▶ hmm

general framework: E+M/variational

- ▶ e.g., GMM+hard clustering gives kmeans
- ▶ e.g., some favorite applications:
 - ▶ hmm
 - ▶ vbmod: [arXiv:0709.3512](https://arxiv.org/abs/0709.3512)

general framework: E+M/variational

- ▶ e.g., GMM+hard clustering gives kmeans
- ▶ e.g., some favorite applications:
 - ▶ hmm
 - ▶ vbmod: [arXiv:0709.3512](https://arxiv.org/abs/0709.3512)
 - ▶ ebfret: [ebfret.github.io](https://github.com/ebfret)

general framework: E+M/variational

- ▶ e.g., GMM+hard clustering gives kmeans
- ▶ e.g., some favorite applications:
 - ▶ hmm
 - ▶ vbmod: [arXiv:0709.3512](https://arxiv.org/abs/0709.3512)
 - ▶ ebfret: [ebfret.github.io](https://github.com/ebfret)
 - ▶ EDHMM: [edhmm.github.io](https://github.com/edhmm)

example application: LDA+topics




Figure 27: From Blei 2003




Figure 28: From Blei 2011

recall: recommendation via factoring

$$\min_{U,V} \sum_{i,j} (r_{ij} - u_i^T v_j)^2 + \lambda_u \|u_i\|^2 + \lambda_v \|v_j\|^2,$$

Figure 29: From Blei 2011

CTM: combined loss function

Maximization of the posterior is equivalent to maximizing the complete log likelihood of U , V , $\theta_{1:J}$, and R given λ_u , λ_v and β ,

$$\begin{aligned} \mathcal{L} = & -\frac{\lambda_u}{2} \sum_i u_i^T u_i - \frac{\lambda_v}{2} \sum_j (v_j - \theta_j)^T (v_j - \theta_j) \\ & + \sum_j \sum_n \log (\sum_k \theta_{jk} \beta_{k, w_{jn}}) - \sum_{i,j} \frac{c_{ij}}{2} (r_{ij} - u_i^T v_j)^2. \end{aligned} \quad (7)$$

Figure 30: From Blei 2011

CTM: updates for factors

$$u_i \leftarrow (VC_iV^T + \lambda_u I_K)^{-1}VC_iR_i \quad (8)$$

$$v_j \leftarrow (UC_jU^T + \lambda_v I_K)^{-1}(UC_jR_j + \lambda_v \theta_j). \quad (9)$$

Figure 31: From Blei 2011

CTM: (via Jensen's, again) bound on loss

$$\begin{aligned}\mathcal{L}(\theta_j) &\geq -\frac{\lambda_n}{2}(v_j - \theta_j)^T(v_j - \theta_j) \\ &\quad + \sum_n \sum_k \phi_{jnk} (\log \theta_{jk} \beta_{k, w_{jn}} - \log \phi_{jnk}) \\ &= \mathcal{L}(\theta_j, \phi_j).\end{aligned}\tag{10}$$

Figure 32: From Blei 2011

Lecture 5 data product

data science and design thinking

- ▶ knowing customer

data science and design thinking

- ▶ knowing customer
- ▶ right tool for right job

data science and design thinking

- ▶ knowing customer
- ▶ right tool for right job
- ▶ practical matters:

data science and design thinking

- ▶ knowing customer
- ▶ right tool for right job
- ▶ practical matters:
 - ▶ munging

data science and design thinking

- ▶ knowing customer
- ▶ right tool for right job
- ▶ practical matters:
 - ▶ munging
 - ▶ data ops

data science and design thinking

- ▶ knowing customer
- ▶ right tool for right job
- ▶ practical matters:
 - ▶ munging
 - ▶ data ops
 - ▶ ML in prod

Thanks!

Thanks MLSS students for your great questions; please contact me @chrishwiggins or [chris.wiggins@{nytimes,gmail}.com](mailto:chris.wiggins@nytimes.com) with any questions, comments, or suggestions!