

Multiplicative Hitchin Systems and Supersymmetric Gauge Theory

Chris Elliott

Vasily Pestun

August 25, 2018

Abstract

1 Introduction

(Chris: There are a lot of notes in red: that just indicates that there are a lot of arguments that are only partially written. They're notes for myself.)

2 Multiplicative Higgs Bundles and q -Connections

We'll begin with an abstract definition of the moduli spaces we'll investigate in this paper using the language of derived algebraic geometry.

Let G be a reductive complex algebraic group, let C be a smooth complex algebraic curve and fix a finite set $D = \{z_1, \dots, z_k\}$ of closed points in C . We write $\mathrm{Bun}_G(C)$ for the moduli stack of G -bundles on C , which we view as a mapping stack $\underline{\mathrm{Map}}(C, BG)$ into the classifying stack of G .

Definition 2.1. The moduli stack of *multiplicative G -Higgs fields* on C with singularities at D is the fiber product

$$\mathrm{mHiggs}_G(C, D) = \mathrm{Bun}_G(C) \times_{\mathrm{Bun}_G(C \setminus D)} \underline{\mathrm{Map}}(C \setminus D, G/G)$$

where G/G is the adjoint quotient stack.

Remark 2.2. A closed point of $\mathrm{mHiggs}_G(C, D)$ consists of a principal G -bundle P on C along with an automorphism of the restriction $P|_{C \setminus D}$, i.e. a section of $\mathrm{ad}(P)$ away from D .

The adjoint quotient stack can also be described as the derived loop space $\underline{\mathrm{Map}}(S_B^1, BG)$ of the classifying stack, where S_B^1 is the “Betti stack” of S^1 , i.e. the constant derived stack at the simplicial set S^1 . We can therefore view $\underline{\mathrm{Map}}(C \setminus D, G/G)$ instead as the mapping stack $\underline{\mathrm{Map}}((C \setminus D) \times S_B^1, BG)$, and the moduli stack of multiplicative Higgs bundles instead as

$$\mathrm{mHiggs}_G(C, D) = \underline{\mathrm{Map}}((C \times S_B^1) \setminus (D \times \{0\}), BG).$$

The source of this mapping stack can be q -deformed. Indeed, let q denote an automorphism of the curve C . Write $C \times_q S_B^1$ for the *mapping torus* of q , i.e the derived fiber product

$$C \times_q S_B^1 = C \times_{C \times C} C$$

where the two maps $C \rightarrow C \times C$ are given by the diagonal and the q -twisted diagonal $(1, q)$ respectively.

Definition 2.3. The moduli stack of q difference connections for the group G on C with singularities at D is the mapping space

$$q\text{-Conn}_G(C, D) = \underline{\text{Map}}((C \times_q S_B^1) \setminus (D \times \{0\}), BG).$$

In particular when $q = 1$ this recovers the moduli stack of multiplicative Higgs bundles.

Remark 2.4. A closed point of $q\text{-Conn}_G(C, D)$ consists of a principal G -bundle P on C along with a q difference connection: an isomorphism of G -bundles $P|_{C \setminus D} \rightarrow q^*P|_{C \setminus D}$ away from the divisor D .

2.1 Residue Conditions

These moduli stacks are typically of infinite type. In order to obtain finite type stacks, and later in order to define symplectic rather than only Poisson structure, we can fix the behaviour of our multiplicative Higgs fields and q difference connections near the punctures $D \subseteq C$.

We'll write \mathbb{D} to denote the *formal disk* $\text{Spec } \mathbb{C}[[z]]$. Likewise we'll write \mathbb{D}^\times for the *formal punctured disk* $\text{Spec } \mathbb{C}((z))$. We'll then write \mathbb{B} for the derived pushout $\mathbb{D} \sqcup_{\mathbb{D}^\times} \mathbb{D}$. Let $LG = \underline{\text{Map}}(\mathbb{D}^\times, G)$ and let $L^+G = \underline{\text{Map}}(\mathbb{D}, G)$.

There is a canonical inclusion $\mathbb{B}^{\sqcup k} \rightarrow (C \times_q S_B^1) \setminus (D \times \{0\})$, the inclusion of the formal punctured neighbourhood of $D \times \{0\}$. This induces a restriction map on mapping spaces

$$\text{res}_D: q\text{-Conn}_G(C, D) \rightarrow \text{Bun}_G(\mathbb{B})^k.$$

The following is well-known (see e.g. the expository article [Zhu17]).

Lemma 2.5. The set of closed points of $\text{Bun}_G(\mathbb{B})$ is in canonical bijection with the set of dominant coweights of G .

Definition 2.6. Choose a map from D to the set of dominant coweights and denote it by $\omega^\vee: z_i \mapsto \omega_{z_i}^\vee$. Write Λ_i for the isotropy group of the point $\omega_{z_i}^\vee$ in $\text{Bun}_G(\mathbb{B})$. The moduli stack of q difference connections on C with singularities at D and fixed residues given by ω^\vee is defined to be the fiber product

$$q\text{-Conn}_G(C, D, \omega^\vee) = q\text{-Conn}_G(C, D) \times_{\text{Bun}_G(\mathbb{B})^k} (B\Lambda_1 \times \cdots \times B\Lambda_k).$$

Examples 2.7. The most important examples for our purposes are given by the following rational/trigonometric/elliptic trichotomy.

- **Rational:** We can enhance the definition of our moduli space by including a framing at a point $c \in C$ not contained in D . We always assume that such framed points are fixed by the automorphism q .

Definition 2.8. The moduli space of q -difference connections on C with a framing at c is defined to be the relative mapping space

$$q\text{-Conn}_G^{\text{fr}}(C) = \underline{\text{Map}}(C, G/G; f)$$

where $f: \{c\} \rightarrow G/G$ is a choice of adjoint orbit. We can define the framed mapping space with singularities and fixed residues in exactly the same way as above.

In this paper we'll be most interested in the following example. Let $C = \mathbb{CP}^1$ with framing point $c = \infty$, and consider automorphisms of the form $z \mapsto z + \varepsilon$ for $\varepsilon \in \mathbb{C}$. Choose a finite subset $D \subseteq \mathbb{A}^1$ and label the points $z_i \in D$ by dominant coweights $\omega_{z_i}^\vee$. We can then study the moduli space $\varepsilon\text{-Conn}_G^{\text{fr}}(\mathbb{CP}^1, D, \omega^\vee)$. The main object of study in this paper will be the holomorphic symplectic structure on this moduli space.

Note that the motivation for this definition comes in part from Spaide's formalism [Spa16] of AKSZ symplectic structures on relative mapping spaces – in this formalism \mathbb{CP}^1 with a single framing point is relatively 1-oriented, so mapping spaces out of it with 1-shifted symplectic targets have AKSZ 0-shifted symplectic structures.

- **Trigonometric:** Alternatively, we can enhance our definition by including a reduction of structure group at a point $c \in C$ not contained in D , again fixed by the automorphism q .

Definition 2.9. The moduli space of q -difference connections on C with an H -reduction at c for a subgroup $H \subseteq G$ is defined to be the fiber product

$$q\text{-Conn}_G^{H,c}(C) = \underline{\text{Map}}(C, G/G) \times_{G/G} H/H$$

associated to the evaluation at c map $\underline{\text{Map}}(C, G/G) \rightarrow G/G$. We can define the moduli space with H -reduction with singularities and fixed residues in the same way as above.

Again let $C = \mathbb{CP}^1$. Fix a pair of opposite Borel subgroups B_+ and $B_- \subseteq G$ with unipotent radicals N_\pm and consider the moduli space of q -connections with B_+ -reduction at 0 and N_- -reduction at ∞ . We'll now take q to be an automorphism of the form $z \mapsto qz$ for $q \in \mathbb{C}^\times$. We'll defer in depth analysis of this example to future work.

- **Elliptic:** Finally, let $C = E$ be a smooth curve of genus one. In this case we won't fix any additional boundary data, but just consider the moduli space $q\text{-Conn}_G(E, D, \omega^\vee)$. In the case $q = 1$ this space – or rather its polystable locus – was studied by Hurtubise and Markman [HM02], who proved that it can be given the structure of an algebraic integrable system with symplectic structure related to the elliptic R-matrix of Etingof and Varchenko [EV98].

(Vasily: What shall we say about higher genus? The definition of mHiggs as above makes sense. Will the resulting space be Poisson? I guess not because to define the Poisson structure we need to fix a section of canonical bundle on C , that is a one-form “dz”. However, in higher genus such 1-form necessarily has zeroes. May be here is a good place to comment or restrict attention to the trichotomy with an explanation why higher genus is on different grounds) (Chris: added a remark below.)

Remark 2.10. In the elliptic case it's natural to ask to what extent Hurtubise and Markman's integrable system structure can be extended from the variety of polystable multiplicative Higgs bundles to the full moduli stack. If D is empty then it's easy to see that we have a symplectic structure given by the AKSZ construction of Pantev-Toën-Vaquié-Vezzosi [PTVV13]. Indeed, E is compact 1-oriented and the quotient stack G/G is 1-shifted symplectic, so the mapping stack $\underline{\text{Map}}(E, G/G) = \text{mHiggs}_G(E)$ is equipped with a 0-shifted symplectic structure by [PTVV13, Theorem 2.5]. The role of the Hitchin fibration is played by the Chevalley map $\chi: G/G \rightarrow T/W$, and therefore

$$\underline{\text{Map}}(E, G/G) \rightarrow \underline{\text{Map}}(E, T/W).$$

The fibers of this map over regular points in T/W are given by moduli stacks of the form $\text{Bun}_T(\tilde{E})^W$ where \tilde{E} is a W -fold cover of E (the cameral cover). (Chris: is the base just T/W ? Come back to this, at least by dimension counting.) (Vasily: If there are no singularities, I think that's correct, but perhaps we can extend the comment above for the case of non-empty D ?)

Remark 2.11. While the moduli space of multiplicative Higgs bundles makes sense on a general curve it's only after restricting identity to this trichotomy of examples that we'll expect the existence of a Poisson structure. Such a structure arises by the AKSZ construction, i.e. by transgressing the 1-shifted symplectic structure on G/G to the mapping space using a fixed section of the canonical bundle on C (possibly with a boundary condition).

2.2 Stability Conditions

For comparison to results in the literature it is important that we briefly discuss the role of stability conditions for difference connections. In our main example of interest – the rational case – these conditions won't play a role, but they do appear in the comparison results between q -connections and monopoles in the literature for more general curves. For definitions for general G we refer to [Smi15], although see also [AB01] on polystable G -bundles. In what follows we fix a choice of $0 < t_0 < 2\pi R$.

Definition 2.12. Let (P, g) be a q -connection on a curve C , and let χ be a character of G . The χ -degree of (P, g) is defined to be

$$\deg_\chi(P, g) = \deg(P \times_\chi \mathbb{C}) - \frac{t_0}{2\pi R} \sum_{i=1}^k \deg(\chi \circ \omega_{z_i}^\vee).$$

(Vasily: Sorry, what that last zero “...(0)” denotes above?) (Chris: I changed the notation. The map $\chi \circ \omega_{z_i}^\vee$ is a map $\mathbb{G}_m \rightarrow \mathbb{G}_m$ and we want its degree.)

A q -connection (P, g) on C is *stable* if for every maximal parabolic subgroup $H \subseteq G$ with Levi decomposition $H = LN$ and every reduction of structure group (P_H, g) to H , we have

$$\deg_\chi(P_H, g) < 0$$

for the character $\chi = \det(\text{Ad}_L^n)$ defined to be the determinant of the adjoint representation of L on \mathfrak{n} .

The q -connection (P, g) is *polystable* if there exists a (not necessarily maximal) parabolic subgroup H with Levi factor L and a reduction of structure group (P_L, g) to L so that (P_L, g) is a stable q -connection and so that the associated H -bundle is admissible, meaning that for every character χ of H which is trivial on $Z(G)$ the associated line bundle $P_H \times_\chi \mathbb{C}$ has degree zero.

Below we'll write $q\text{-Conn}_G^{\text{ps}}(C, D, \omega^\vee)$ for the moduli space of polystable q -connections. This moduli space is a smooth algebraic variety of finite type. (Chris: Reference for this other than for q trivial?)

When $C = \mathbb{CP}^1$ every principal G -bundle on C admits an essentially unique (up to the action of the Weyl group) holomorphic reduction of structure group to a maximal torus [Gro57]. Since q -connections for an abelian group are automatically stable, polystability on the sphere is equivalent to admissibility of the torus reduction. As a consequence, for our main example of interest – the rational case – the moduli space of polystable q -connections is equivalent to the moduli space of all q -connections of admissible degree. For instance for $G = \text{SL}_n$ the moduli space of polystable q -connection is equivalent to the moduli space of q -connections on the trivial bundle.

3 Periodic Monopoles

Moduli spaces of q -connections on a Riemann surface C are closely related to moduli spaces of periodic monopoles, i.e. monopoles on 3-manifolds that fiber over the circle (more specifically, with fiber C and monodromy determined by q). Let $G_{\mathbb{R}}$ be a compact Lie group whose complexification is G . The discussion in this section will mostly follow that of [CH10, Smi15].

Write $M = C \times_q S_R^1$ for the C -bundle over S^1 with monodromy given by the automorphism q . More precisely, M is the Riemannian 3-manifold obtained by gluing the ends of the product $C \times [0, 2\pi R]$ of Riemannian manifolds by the isometry $(x, 2\pi R) \sim (q(x), 0)$.

Definition 3.1. A *monopole* on the Riemannian 3-manifold $M = C \times_q S_R^1$ is a smooth principal $G_{\mathbb{R}}$ -bundle P equipped with a connection A and a section Φ of the associated bundle \mathfrak{g}_P satisfying the Bogomolny equation

$$*F_A = d_A \Phi.$$

Remark 3.2. We should emphasise the difference between the Riemannian 3-manifold $M = C \times_q S_R^1$ appearing in this section and the derived stack $C \times_q S_B^1$ (the mapping torus) appearing in the previous section. These should be thought of as smooth and algebraic realizations of the same object (justified by the comparison Theorem 3.6) but they are a priori defined in different mathematical contexts.

(Vasily: We should specify the Riemannian metric on M in order to define the $*$ operation on forms. I guess our convention for the metric on M which leads to the symplectomorphism with mHiggs is

$$ds^2 = v \otimes \bar{v} + dt \otimes dt \tag{1}$$

where $v = "dz"$ is a chosen section of the canonical bundle on C ; I suggest to give some fixed coordinate-free name to this holomorphic volume form on C , like v or whatever, and may be even consider that that this volume form is fixed as an attribute of C through out the paper)) (Chris: I added a description of the Riemannian metric. I think we won't need to specify a section of the canonical bundle until Section 4.)

We can rephrase the data of a monopole on M as follows. Let $C_0 = C \times \{0\}$ be the fiber over 0 in S^1 , viewed as a Riemann surface. Let P be the restriction of the complexified bundle $\mathbf{P}_{\mathbb{C}}$ to C_0 . Consider first the restriction of the complexification of A to a connection A_0 on P over C_0 . The $(0, 1)$ part of A_0 automatically defines a holomorphic structure on P . We can introduce an additional piece of structure on this holomorphic G -bundle. In order to do so we can decompose the Bogomolny equation into one real and one complex equation as follows.

$$\begin{aligned} F_{A_0} - \nabla_t \Phi &= 0 \\ [\bar{\partial}_{A_0}, \nabla_t - i\Phi dt] &= 0 \end{aligned}$$

where ∇_t is the component of the covariant derivative d_A normal to C_0 . From now on we'll denote $\mathcal{A} = \nabla_t - i\Phi dt$.

Let's now introduce singularities into the story. We'll keep the description brief, referring the reader to [CH10, Smi15] for details.

Definition 3.3. Let $D \subseteq M$ be a finite subset. Let ω^\vee be a choice of coweight for G . A monopole on $M \setminus D$ has *Dirac singularity* at $z \in D$ with charge ω^\vee if locally on a neighbourhood of z in M it is obtained by pulling back under ω^\vee the standard Dirac monopole solution to the Bogomolny equation where Φ is spherically symmetric with a simple pole at z () and the restriction of a connection A to a two-sphere S^2 enclosing the singularity defines a $U(1)$ bundle on this S^2 of degree 1 so that

$$\frac{1}{2\pi} \int_{S^2} F = 1.$$

See e.g. [CH10, Section 2.2] for a more detailed description.

We can also introduce a framing (or a reduction of structure group as in the trigonometric example, though we won't consider the latter in this paper). As usual let $c \in C$ be a point fixed by the automorphism q .

Definition 3.4. A monopole on M with *framing* at the point $c \in C$ is a monopole (\mathbf{P}, A, Φ) on M (possibly with Dirac singularities at D) along with a trivialization of the restriction of \mathbf{P} to the circle $\{c\} \times S_R^1$, with the condition that the holonomy of \mathcal{A} around this circle lies in a fixed conjugacy class $f \in G/G$.

The moduli theory of monopoles on 3-manifolds of this form has been studied in the mathematics literature by Foscolo [Fos16], applying the analytic techniques of deformation theory to earlier work on periodic monopoles by Cherkis and Kapustin [CK98, CK01]. (Vasily: need to add refs to Kapustin Cherkis and others before Foscolo) (Chris: added something here. Are there other earlier references regarding deformation / moduli theory you had in mind?). In the cases of interest to us it can be obtained as a hyperkähler quotient. Let's focus initially on the rational case, so monopoles on $M = \mathbb{CP}^1 \times_\varepsilon S_R^1$ with Dirac singularities at $D \times \{t_0\}$ and a framing at ∞ . Consider the infinite-dimensional vector space \mathcal{V} consisting of pairs (A, Φ) where A is a connection on a fixed principal $G_{\mathbb{R}}$ -bundle \mathbf{P} on M , Φ is a section of $\mathfrak{g}_{\mathbf{P}}$, and (A, Φ) have a Dirac singularity with charge $\omega_{z_i}^\vee$ at each (z_i, t_0) in $D \times \{t_0\}$. Let \mathcal{G} be the group of gauge transformations of the bundle \mathbf{P} .

The hyperkähler moment map is given by the Bogomolny functional, namely

$$\begin{aligned} \mu: \mathcal{V} &\rightarrow \Omega^1(M; (\mathfrak{g}_{\mathbb{R}})_{\mathbf{P}}) \\ (A, \Phi) &\mapsto *F_A - d_A \Phi. \end{aligned}$$

Definition 3.5. Let D be a finite subset $\{(z_1, t_1), \dots, (z_k, t_k)\}$ of points in $M = \mathbb{CP}^1 \times_\varepsilon S_R^1$, and let ω_i^\vee be a choice of coweight for each point in D . The moduli space $\text{Mon}_G(M, D, \omega^\vee)$ is the hyperkähler quotient

$$\text{Mon}_G(M, D, \omega^\vee) = \mu^{-1}(0)/\mathcal{G}.$$

Now let's address the relationship between periodic monopoles and q -connections. Suppose from now on that q is in the identity component of the group of automorphisms of C (fixing the framing point c if present).

Theorem 3.6. There is an analytic isomorphism between the moduli space of polystable monopoles on $C \times_q S^1$ with Dirac singularities at $D \times \{t_0\}$ (and a possible framing on $\{c\} \times S^1$) and the moduli space of q -connections on C with singularities at D and framing at $\{c\}$. More precisely there is an analytic isomorphism

$$H: \text{Mon}_G^{(\text{fr})}(C \times_q S^1, D \times \{t_0\}, \omega^\vee) \rightarrow q\text{-Conn}_G^{\text{ps}, (\text{fr})}(C, D, \omega^\vee)$$

given by the holonomy map around S^1 , i.e. sending a monopole $(\mathbf{P}, \mathcal{A})$ to the holomorphic bundle $P = (\mathbf{P}_C)|_{C_0}$ with q -connection $g = \text{Hol}_{S^1}(\mathcal{A}): P \rightarrow q^*(P)$. (Vasily: Do we assume that $C_0 = C \times t_0$ or not necessarily the same t_0 where are the singularities?) (Chris: I think I want that $C_0 = C \times \{0\}$, a slice explicitly different from the slice where the singularities live.)

Proof. (Chris: check) This follows by the same argument as that given by Charbonneau-Hurtubise [CH10] and Smith [Smi15]. More explicitly, first let's think about injectivity, so let $(\mathbf{P}, \mathcal{A})$ and $(\mathbf{P}', \mathcal{A}')$ be a pair of periodic monopoles on $C \times_q S^1$ with images (P, g) and (P', g') respectively, and choose a bundle isomorphism $\tau: P \rightarrow P'$ intertwining the q -connections g and g' . One observes first that \mathbf{P} and \mathbf{P}' are also isomorphic G -bundles since, by intertwining with the q -connections, we have an isomorphism $\mathbf{P}|_{C \times \{t\}} \rightarrow \mathbf{P}'|_{C \times \{t\}}$ for every $t \in S^1$. That the monopole structures also match up follows by the same argument as in [CH10, Proposition 4.7].

For surjectivity, again we'll match the argument in the case where $q = \text{id}$. We begin by extending a holomorphic G -bundle P on C_0 with q -connection g to a G -bundle on $M \setminus (D \times \{t_0\}) = (C \times_q S_R^1) \setminus (D \times \{t_0\})$. Let $\gamma: [-2\pi R, 2\pi R] \rightarrow \text{Aut}(C)$ (Vasily: Sorry, what is the target space of map γ ? Perhaps not G but the group of automorphisms of the curve C ? Also, now we are assuming that there exists group homomorphism from real numbers to the automorphism group of C . It is true for the rational, and trigonometric case. However, what if automorphism group of the curve C includes some discrete factor, and q is that discrete automorphism. In this case we will not be able to smoothly interpolate. Perhaps we need to restrict explicitly to a subgroup of automorphism group of C which is covered by the group of real numbers or circle?) (Chris: Yes, I did try to make that explicit in the sentence before the theorem statement.)

be a geodesic with $\gamma(-2\pi R) = q^{-1}$, $\gamma(0) = 1$ and $\gamma(2\pi R) = q$. Let \widetilde{M} be the 3-manifold

$$\widetilde{M} = ((-2\pi R, 2\pi R) \times C) \setminus \bigcup_{j=1}^k (A_j^+ \cup A_j^-)$$

where A_j^+ is the arc $\{(t + t_0, \gamma(t)(z_j)): t \in (0, 2\pi R - t_0)\}$ and A_j^- is the arc $\{(t + t_0 - 4\pi R, \gamma(t)(z_j)): t \in [2\pi R - t_0, 2\pi R)\}$.

Let $\pi: \widetilde{M} \rightarrow C$ be the projection sending (t, z) to $\gamma(t)(z)$. The bundle P pulls back to a bundle $\pi^*(P)$ on \widetilde{M} . We obtain a bundle on $M \setminus (D \times t_0)$ by applying the identification $(t, z) \sim (t - 2\pi R, q(z))$. This bundle extends to an S^1 -invariant holomorphic G -bundle on $M \times S^1$. The remainder of the proof – verifying the existence of the monopole structure associated to an appropriate choice of hermitian structure – consists of local analysis which is independent of the value of the parameter q .

It remains to remark on the compatibility of framing data on the two sides. A trivialization of the bundle \mathbf{P} along the circle $\{c\} \times S^1$ yields a trivialization of the fiber of the bundle P at c . The condition that the holonomy around the circle at c is f fixes the value of the q -connection at c to be in the conjugacy class f .

□

Remark 3.7. Mochizuki [Moc17] proved a stronger result in the rational case for the group $G = \text{GL}_n$. He allows not just a framing at infinity in \mathbb{CP}^1 but also a singularity encoded in terms of a B -reduction of the bundle.

3.1 Deformation Theory

(Vasily: Do we assume in this section the case of rational C ? I think that some of our statements hold for generic C and some for rational case) (Chris: I believe everything in this section will make sense generally, and starting in the subsequent section we'll want to restrict to the rational case.)

In the next section we'll compare symplectic forms on these moduli spaces. In order to do so it will be important to understand the tangent spaces at a point of the source and target. There's a natural description of these tangent spaces in terms of the hypercohomology of certain cochain complexes.

Let's first describe the tangent complex to the moduli space of monopoles. For simplicity we'll restrict attention to the rational case where we can use the description as a hyperkähler reduction. For more general deformation theory calculations we refer to Foscolo [Fos16]. Recall that in this case we can write

$$\text{Mon}_G^{(\text{fr})}(C \times_q S^1, D \times \{t_0\}, \omega^\vee) \cong \mu^{-1}(0)/\mathcal{G}$$

where \mathcal{G} is the group of gauge transformations of \mathbf{P} and $\mu: \mathcal{V} \rightarrow \Omega^1(M; (\mathfrak{g}_R)_{\mathbf{P}})$ is the Bogomolny functional $(A, \Phi) \mapsto *F_A - d_A\Phi$. The tangent complex to this hyperkähler quotient at a point $(\mathbf{P}, \mathcal{A})$ can be written as $\Omega^0(M \setminus D; (\mathfrak{g}_R)_{\mathbf{P}})[1] \rightarrow \mathbb{T}_{\mu^{-1}(0)}$ where $\mathbb{T}_{\mu^{-1}(0)}$ is the tangent complex to the zero locus of the moment map, concentrated in non-negative degrees. Roughly speaking $\mathbb{T}_{\mu^{-1}(0)} = \mathbb{T}_{\mathcal{V}} \xrightarrow{d\mu} \Omega^1(M; (\mathfrak{g}_R)_{\mathbf{P}})[-1]$.

More explicitly, following [Fos16] let

$$\mathcal{F}_{\mathbf{P}, \mathcal{A}}^{\text{mon}} = \left(\Omega^0(M \setminus D; (\mathfrak{g}_R)_{\mathbf{P}}) \xrightarrow{d_1} \Omega^1(M \setminus D; (\mathfrak{g}_R)_{\mathbf{P}}) \oplus \Omega^0(M \setminus D; (\mathfrak{g}_R)_{\mathbf{P}}) \xrightarrow{d_2} \Omega^1(M \setminus D; (\mathfrak{g}_R)_{\mathbf{P}}) \right) \otimes_{\mathbb{R}} \mathbb{C}$$

placed in degrees $-1, 0$ and 1 where $d_1(g) = -(d_A(g), [\Phi, g])$ and $d_2(a, \psi) = *d_A(a) - d_A(\psi) + [\Phi, a]$. Write d_{mon} for the total differential.

Remark 3.8. Here we've chosen a point in the twistor sphere, forgetting the hyperkähler structure and retaining a holomorphic symplectic structure. In other words we've identified the target of the hyperkähler moment map with $\mathbb{R} \oplus \mathbb{C}$. (Vasily: Somewhere here we should clarify that the hyperKähler moment map takes value in the space imaginary quaternions which is isomorphic as real space to $\mathbb{R}_{\text{quat}}^3$. After we pick a complex structure on the twistor sphere, there is natural split $\mathbb{R}_{\text{quat}}^3 = \mathbb{C} \oplus \mathbb{R}$. If we care only about complex structure and holomorphic symplectic form on the tangent space to the moduli space of monopoles with respect to a certain fixed point on a twistor sphere, we trade the real component of the equation in the split $\mathbb{R}_{\text{quat}}^3 = \mathbb{C} \oplus \mathbb{R}$ in the third term with a complexification of the gauge group transformation in the first term in which we would replace $\mathfrak{g}_{\mathbb{R}}$ by $\mathfrak{g}_{\mathbb{C}}$.)

Remark 3.9. If we restrict $\mathcal{F}_{\mathbf{P}, \mathcal{A}}^{\text{mon}}$ to a slice $C_t = C \times \{t\}$ in the t -direction we can identify it with a complex of the form

$$\Omega^\bullet(C_t; \mathfrak{g}_P)[1] \xrightarrow{[\Phi, -]} \Omega^\bullet(C_t; \mathfrak{g}_P)$$

with total differential given by d_A on each of the two factors along with the differential $[\Phi, -]$ mixing the two factors. These two summands each split up into the sum of a Dolbeault complex on C with its dual. That is, there's a natural subcomplex of the form

$$\Omega^{0, \bullet}(C_t; \mathfrak{g}_P)[1] \xrightarrow{[\Phi, -]} i\Omega^{0, \bullet}(C_t; \mathfrak{g}_P)dt$$

where the internal differentials on the two factors are now given by $\bar{\partial}_{A_0}$. This complex is in turn quasi-isomorphic to the complex

$$\Omega^\bullet(S^1; \Omega^{0, \bullet}(C_t; \mathfrak{g}_P))[1]$$

with total differential $\bar{\partial}_{A_0} + d_A$.

(Vasily: Sorry, I am afraid I did not understand the notations of the remark and hence the statement. Anyways, I was expecting, in the line of our discussions, that the relevant differential, from which we would find the tangent bundle to Mon and then determine symplectic form, is given by

$$D = dtD_t + d\bar{z}D_{\bar{z}}, \quad D_t = \partial_t + A_t + i\Phi, \quad D_{\bar{z}} = \partial_{\bar{z}} + A_{\bar{z}} \quad (2)$$

that is a mixture of Dol differential along C and de Rham differential along S^1 . And we should have been referring here to the choice of the Riemmanian metric on M (1) to make the claims and the twistor sphere point corresponding to the split of $T_M \simeq \mathbb{R}_{quat}^3 \simeq \mathbb{C} \oplus \mathbb{R} = T_C \oplus T_{S^1}$) (Chris: I rewrote the remark a bit: I think the differential I'm describing and the differential you're describing are actually the same. Though note it needs editing to get the signs and factors of i correct.)

Remark 3.10. If one introduces a framing at a point $c \in C$ then we must correspondingly twist the complex \mathcal{F}^{mon} above by the line bundle $\mathcal{O}(-c)$ on C . So in that case we define (Chris: check)

$$\mathcal{F}_{(P,A)}^{\text{mon,fr}} = \mathcal{F}_{P,A}^{\text{mon}} \otimes (\mathbb{C}_{S^1} \boxtimes \mathcal{O}(-c)).$$

The following is proved in [Fos16].

Proposition 3.11. The tangent complex of $\text{Mon}_G(S^1 \times C, D \times \{t_0\}, \omega^\vee)$ at the point (P, A) is quasi-isomorphic to the hypercohomology $\mathbb{H}^\bullet(C \times S^1; \mathcal{F}'_{(P,A)})$ of a subsheaf $\mathcal{F}' \subseteq \mathcal{F}^{\text{mon}}$ where growth conditions are imposed on the degree 0 part of \mathcal{F}^{mon} near the singularities.

Now let's consider the tangent complex to the moduli space of q -connections. For the arguments in this article we'll only need to carefully consider the case $q = \text{id}$ of multiplicative Higgs bundles, but we'll include some remarks regarding the more general case. In this case the calculation was performed by Bottacin [Bot95], see also [HM02, Section 4]. Fix a multiplicative Higgs bundle (P, g) on C . We consider the sheaf of cochain complexes on C

$$\mathcal{F}_{(P,g)} = (\mathfrak{g}_P[1] \xrightarrow{\text{Ad}_g} \mathfrak{g}_P(D))$$

in degrees -1 and 0 with differential given by the adjoint action of g . More precisely let L_g and R_g be the bundle maps $\mathfrak{g}_P \rightarrow \mathfrak{g}_P$ obtained as the derivative of left- and right-multiplication. Then $\text{Ad}_g = L_g - R_g$. We can alternatively phrase this, as in [HM02, Section 4], as follows. Define $\text{ad}(g)$ to be the vector bundle

$$\text{ad}(g) = (\mathfrak{g}_P \oplus \mathfrak{g}_P) / \{(X, -gXg^{-1}) : X \in \mathfrak{g}_P\}.$$

Then we can write \mathcal{F} as the sheaf of complexes

$$\mathcal{F}_{(P,g)} = (\mathfrak{g}_P[1] \xrightarrow{\text{Ad}_g} \text{ad}(g))$$

where now Ad_g is just the map $X \mapsto [(X, -X)]$.

Remark 3.12. If one introduces a framing at a point $c \in C$ then we must correspondingly twist the complex \mathcal{F} above by the line bundle $\mathcal{O}(-c)$ on C , i.e. we restrict to deformations that preserve the framing and therefore are zero at the point c . So in that case we define

$$\mathcal{F}_{(P,g)}^{\text{fr}} = (\mathfrak{g}_P[1] \xrightarrow{\text{Ad}_g} \mathfrak{g}_P(D)) \otimes \mathcal{O}(-c).$$

Remark 3.13. For more general q we should modify this description by replacing g by a q -connection. Note that one can still define the (q -twisted) adjoint action $X \mapsto gXg^{-1}$ using a q -connection, and so we can still define the complex

$$\mathcal{F}_{(P,g)} = (\mathfrak{g}_P[1] \xrightarrow{\text{Ad}_g} \text{ad}(g))$$

just as in the untwisted case (Chris: check). (Vasily: I wonder what would replace the rational r -matrix that defines the Poisson structure on the Lie group of rational Higgs fields given the explicit formula for the symplectic structure on the moduli space of q -connections (5) for non-trivial q ? And if we quantize the space of functions on q -deformed big mHiggs what do we get instead of completed Yangian when q is trivial, say for the case of a single singularity at 0)

Proposition 3.14. The tangent complex of $\text{mHiggs}_G(C, D)$ at the point (P, g) is quasi-isomorphic to the hypercohomology $\mathbb{H}^\bullet(C; \mathcal{F}_{(P,g)})$ of the sheaf \mathcal{F} .

Proof. We'll explain the proof in different language to that used in [Bot95]. Recall that for a mapping stack $\text{Map}(X, Y)$ where X and Y are derived Artin stacks the tangent complex at a point f can be computed as the cohomology $\mathbb{T}_{\text{Map}(X, Y), f} \cong \mathbb{H}^*(X, f^* \mathbb{T}_Y)$. Let $X = C \setminus D$ and let $Y = G/G$, so $\mathbb{T}_{(P, g), \text{Map}(C \setminus D, G/G)} \cong \mathbb{H}^*(C \setminus D; \mathfrak{g}_P[1]|_{C \setminus D} \rightarrow \mathfrak{g}_P|_{C \setminus D})$ where the map $\mathfrak{g}_P|_{C \setminus D} \rightarrow \mathfrak{g}_P|_{C \setminus D}$ is given by the adjoint action of g . In order to compute the tangent complex to $\text{mHiggs}_G(C, D)$ we must impose the conditions that the deformation of the G -bundle P extends across the divisor D and that the deformation of the Higgs field has a simple pole at D . In other words we find

$$\mathbb{T}_{\text{mHiggs}_G(C, D), (P, g)} = \mathbb{H}^\bullet(C; \mathfrak{g}_P[1] \rightarrow \mathfrak{g}_P(D)) \subseteq \mathbb{H}^*(C \setminus D; \mathfrak{g}_P[1]|_{C \setminus D} \rightarrow \mathfrak{g}_P|_{C \setminus D}).$$

(Chris: This last step isn't rigorous, needs improving.) □

Remark 3.15. The embedding of the moduli space $\text{mHiggs}_G(C, D, \omega^\vee)$ into $\text{mHiggs}_G(C, D)$ induces an isomorphism of tangent complexes (in degree 0). In other words, all infinitesimal deformations of $(P, g) \in \text{mHiggs}_G(C, D, \omega^\vee)$ preserve the residue condition at D . (Chris: check and justify. To insert here: the fact that the tangent space is concentrated in degree 0.)

In order to calculate hypercohomology we'll use a Čech resolution. This will be straightforward for the multiplicative Higgs moduli space, and we'll use the isomorphism of Theorem 3.6 to give an analogous description on the monopole side. We define a cover $\mathcal{U} = \{U_0, U_1, \dots, U_k, U_\infty\}$ of C as follows. Let U_i be a contractible open neighbourhood of the point z_i and let U_∞ be a contractible analytic open neighbourhood of $c \in C$, all chosen to be pairwise disjoint. Let U_0 . Finally let $U_0 = C \setminus (D \cup \{c\})$. Since the U_i are contractible and the remaining subset $C \setminus (D \cup \{c\})$ is an affine algebraic curve, which means that for any quasi-coherent sheaf of cochain complexes the higher cohomology groups vanish. Likewise for the intersections: the punctured open sets U_i^\times are analytic open sets of an affine curve.

Specify a representative 0-cocycle $(\alpha_\infty, \{\alpha_i\}, \alpha_0, \beta_\infty, \{\beta_i\})$ for the Čech cohomology group with respect to our chosen cover \mathcal{U} . Explicitly a 0-cochain is given by the following data:

$$\begin{aligned} \alpha_\infty &\in \text{ad}(g)(-1)(U_\infty) \\ \alpha_i &\in \text{ad}(g)(1)(U_i) \text{ for } i = 1, \dots, k \\ \alpha_0 &\in \text{ad}(g)(C \setminus (D \cup \{c\})) \\ \beta_\infty &\in \mathfrak{g}_P(U_\infty^\times) \\ \beta_i &\in \mathfrak{g}_P(U_i^\times) \text{ for } i = 1, \dots, k \end{aligned}$$

where the notation (± 1) indicates tensoring by the line bundle $\mathcal{O}(\pm 1)$.

Being a 0-cocycle means that $(\alpha_\infty - \alpha_0)|_{U_\infty^\times} = \text{Ad}_g(\beta_\infty)$ and $(\alpha_i - \alpha_0)|_{U_i^\times} = \text{Ad}_g(\beta_i)$ for each i . We consider 0-cocycles modulo 0-coboundaries of the form $(\text{Ad}_g(f_\infty), \{\text{Ad}_g(f_i)\}, \text{Ad}_g(f_0), f_\infty - f_0, \{f_i - f_0\})$.

Now Ad_g is an isomorphism on U_0 for the sections α_0 of \mathfrak{g}_P that occur: those with no poles or zeroes in U_0 . That means that we can add a coboundary to force $\alpha_0 = 0$. We can likewise use the freedom in f_i to find a representative cocycle where the β_i are all zero. We now have a unique representative cocycle of the form $(\alpha_\infty, \{\alpha_i\}, 0, 0, 0)$.

Now, rather than describing the tangent complex to the moduli space of monopoles we'll define a complex that maps to it which we can define locally with respect to the cover \mathcal{U} . Consider the open cover $\mathcal{U} \times S^1 = \{U_i \times S^1\}$ of $M = C \times S^1$. For each element U_i of the cover we can define a map

$$\begin{aligned} \mathcal{F}'(U_i \times S^1) &\rightarrow (\Omega^0(S^1; \Omega^{0, \bullet}(U_i; \mathfrak{g}_P)) \rightarrow \Omega^1(S^1; \Omega^{0, \bullet}(U_i; \mathfrak{g}_P)(D_{U_i}))) [1] \\ &\cong \mathfrak{g}_P(U_i)[1] \rightarrow \mathfrak{g}_P(U_i)(D|_{U_i}) \\ &\cong \mathcal{F}(U_i) \end{aligned}$$

where we restrict the sheaf \mathcal{F}' whose hypercohomology calculated the monopole tangent complex to the holomorphic part of the slice at $\{t\} \in S^1$. Altogether this defines a map from the Čech cohomology with respect to this cover to the tangent complex of the moduli space of monopoles. That is, we have an explicit map

$$\check{\mathbb{H}}^\bullet(M, \mathcal{U} \times S^1, \mathcal{F}') \rightarrow \mathbb{H}^\bullet(M; \mathcal{F}')$$

that factors through the (isomorphic) tangent complex of the moduli space of multiplicative Higgs bundles. To verify this we need only note that these maps commute with the differentials in the Čech complex, i.e. with the restriction to the intersection of a pair of open sets. (Chris: todo)

(Chris: Summary of the argument I'm trying to make, I'm defining a map $\check{H}^\bullet(M, \mathcal{U} \times S^1, \mathcal{F}') \rightarrow \mathbb{H}^\bullet(M; \mathcal{F})$. Maybe it'll be surjective (the map on cohomology, obviously this doesn't make sense at the cochain level) but I don't think this will be important. Pairings on the source naturally induce pairings on the target, which means we can define a symplectic structure locally on open sets in $\mathcal{U} \times S^1$.)

Explicitly a 0-cochain in this Čech complex is given by $(\alpha_\infty, \{\alpha_i\}, \alpha_0, \beta_\infty, \{\beta_i\})$ where now

$$\begin{aligned} \alpha_\infty &\in \mathcal{F}_0^{\text{mon}}(U_\infty \times S^1) \\ \alpha_i &\in \mathcal{F}_0^{\text{mon}}(U_i \times S^1) \text{ for } i = 1, \dots, k \\ \alpha_0 &\in \mathcal{F}_0^{\text{mon}}(U_0 \times S^1) \\ \beta_\infty &\in \Omega_{\mathbb{C}}^0(U_\infty^\times \times S^1; \mathfrak{g}_{P_c}) \\ \beta_i &\in \Omega_{\mathbb{C}}^0(U_i^\times \times S^1; \mathfrak{g}_{P_c}) \text{ for } i = 1, \dots, k. \end{aligned}$$

Here we write $\mathcal{F}_0^{\text{mon}}$ to indicate the degree 0 term in the cochain complex. There's a similar condition for being a 0-cocycle involving the differential d_{mon} . Again we can add a 0-coboundary to find a representative cocycle of the form $(\alpha_\infty, \{\alpha_i\}, 0, 0, 0)$.

To conclude this subsection it will also be important to have an explicit description of the derivative of the holonomy map H as a map between tangent spaces. We can describe this map using our Čech resolutions on each contractible open set U_i individually.

Proposition 3.16. The derivative $dH: \mathbb{H}^\bullet(U_i \times S^1; \mathcal{F}'_{(P, \mathcal{A})}) \rightarrow \mathbb{H}^\bullet(\mathbb{D}_i; \mathcal{F}_{(P, g)})$ is given on an open patch $U_i \times (0, 2\pi)$ by the formula

$$dH(\alpha_i) = dH(d_{\text{mon}} b_i) = b_i(2\pi)H(\mathcal{A}) - H(\mathcal{A})b_i(0)$$

where $i = 1, \dots, k$ or ∞ . More precisely by $b_i(2\pi)$ and $b_i(0)$ we mean the limit of $b_i(t)$ as $t \rightarrow 2\pi$ or 0 respectively.

Proof. Note that the right-hand side is the derivative at \mathcal{A} of the map $B_i \mapsto B_i(2\pi)H(\mathcal{A})B_i(0)^{-1}$ where $B_i \in \Omega^0((U_i \times (0, 2\pi)) \setminus \{(z_i, t_0)\}; \mathfrak{g}_P)$. This is the definition of the action of the group of gauge transformations on the holonomy $H(\mathcal{A})$ from $t = 0$ to 2π . (Chris: say more?) \square

4 Symplectic Structures

From now on we'll exclusively study the rational situation where $C = \mathbb{CP}^1$ and we fix a framing point $c = \infty$.

4.1 The Example of GL_2

(Chris: Ultimately we might move this, but we should probably include a discussion of what happens for GL_2 before or in parallel to discussing the general story.)

(Vasily: Chris, is the idea to put here explicit expressions for Higgs field in terms of some chart Darboux coordinates (p, q) that we have discussed in the past drafts draft?. If all singularities are miniscule, then for GL_2 the leaves could be described quite explicitly as products of regular semi-simple co-adjoint orbits in \mathfrak{gl}_2 . Rouven is finishing a draft with explicit parametrization by Darboux coordinates of a certain family of symplectic leaves for GL_n . Perhaps here we can put a reference on it.) (Chris: My idea was to include the discussion of, at least, the simple example that you worked out for GL_2 with minimal singularities where you described the geometry of the moduli space and gave an expression for the symplectic form. I really only had in mind for GL_2 trying to give a more explicit

argument for what will appear below, i.e. that the moduli spaces after fixing residues were naturally symplectic. But maybe we could do what you suggest and described the symplectic leaves in some Darboux coordinates and refer to Rouven's paper for proof?)

4.2 Symplectic Structures for General G

We begin the more anstract general analysis by briefly discussing the holomorphic symplectic structure on the moduli space of periodic monopoles on \mathbb{CP}^1 following the analysis of Cherkis and Kapustin [CK98, CK02]. This structure arises from the description we gave as a hyperkähler quotient. To describe it specifically, let $\delta^{(1)}\mathcal{A}$ and $\delta^{(2)}\mathcal{A}$ be two tangent vectors at $(\mathbf{P}, \mathcal{A})$ to the moduli space of monopoles. Choose representatives for these two tangent vectors of the form α_i and α'_i respectively in the Čech resolution we described above. Then we can write the holomorphic symplectic form coming from the hyperkähler reduction in terms of the symplectic pairing on the infinite-dimensional vector space \mathcal{V} , which is given by the Killing form on \mathfrak{g} along with the wedge pairing of differential forms. So summing over the local patches in our Čech resolution we can write it as

$$\begin{aligned}\omega_{\text{mon}}(\delta^{(1)}\mathcal{A}, \delta^{(2)}\mathcal{A}) &= \int_M \kappa(\delta^{(1)}\mathcal{A} \wedge \delta^{(2)}\mathcal{A}) dz dt \\ &= \sum_{i=1}^k \int_{U_i \times S^1} \kappa(\alpha_i \wedge \alpha'_i) dz dt\end{aligned}$$

where the contributions to the integral away from the U_i vanish.

(Vasily: we should start from the expression like

$$\omega(\delta^{(1)}\mathcal{A}, \delta^{(2)}\mathcal{A}) = \int_M \kappa(\delta^{(1)}\mathcal{A} \wedge \delta^{(2)}\mathcal{A}) \wedge v \quad (3)$$

where $v = "dz"$ is the CY 1-form and where we should have had defined before (did we ?)

$$\mathcal{A} = A_{\bar{z}} d\bar{z} + A_z dz + (A_t + i\Phi) dt \quad (4)$$

) (Chris: altered what was written a bit here.)

Let's now consider the moduli space of framed q -connections on \mathbb{CP}^1 . This moduli space has an algebraic symplectic structure with a natural abstract origin, via *derived Poisson geometry*. We refer to [CPT⁺17] for the theory of derived Poisson structures and to [MS16, MS17, Spa16] for that of derived coisotropic structures.

Lemma 4.1. The restriction map $r: \text{Bun}_G^{\text{fr}}(\mathbb{CP}^1 \setminus D) \rightarrow \text{BLG}^k$ has a 2-shifted Lagrangian structure with respect to the residue pairing on BLG .

Remark 4.2. More precisely, BLG is a Tate stack in the sense of Hennion [Hen15]. Although we won't check that the residue pairing symplectic structure ω is compatible with the Tate structure, this problem doesn't arise when defining a Lagrangian structure as a potential for $r^*\omega$ in $\Omega^{2, \text{cl}}(\text{Bun}_G^{\text{fr}}(\mathbb{CP}^1 \setminus D), 2)$ since the stack in question is of ind-finite type.

Proof. Consider the inclusion $\text{dr}: \mathfrak{g}_- = \mathbb{T}_{\text{Bun}_G^{\text{fr}}(\mathbb{CP}^1 \setminus D)}[-1] \rightarrow r^*\mathbb{T}_{\text{BLG}^k}[-1] = \mathfrak{g}((z))^k$: a map of ind-pro Lie algebras concentrated in degree zero. The residue pairing vanishes after pulling back along r since elements of \mathfrak{g}_- are \mathfrak{g} -valued functions on \mathbb{CP}^1 with at least a simple pole at every puncture in D . So the map r is isotropic with 0 isotropic structure; this structure is unique for degree reasons. We must check that this structure is non-degenerate. It suffices to check that the sequence

$$\mathbb{T}_{\text{Bun}_G^{\text{fr}}(\mathbb{CP}^1 \setminus D)}[-1] \rightarrow r^*\mathbb{T}_{\text{BLG}^k}[-1] \rightarrow (\mathbb{T}_{\text{Bun}_G^{\text{fr}}(\mathbb{CP}^1 \setminus D)}[-1])^\vee$$

is an exact sequence of ind-pro vector spaces, and therefore an exact sequence of quasi-coherent sheaves on the stack $\text{Bun}_G^{\text{fr}}(\mathbb{CP}^1 \setminus D)$. To do this we identify the pair $(\mathfrak{g}_-, \mathfrak{g}((z))^k)$ as part of a Manin triple, where a complementary

isotropic subalgebra to \mathfrak{g}_- is given by $\mathfrak{g}_+ = \mathfrak{g}[[z]]^k$. Using the residue pairing we can identify \mathfrak{g}_+ with $(\mathfrak{g}_-)^{\vee}$ and therefore identify our sequence with the split exact sequence

$$0 \rightarrow \mathfrak{g}_- \rightarrow \mathfrak{g}((z))^k \rightarrow \mathfrak{g}_+ \rightarrow 0.$$

□

Construction 4.3. Recall that we can identify the moduli space of singular q -connections on a curve as a fiber product: $q\text{-Conn}_G^{\text{fr}}(\mathbb{CP}^1, D) \cong \text{Bun}_G^{\text{fr}}(C) \times_{\text{Bun}_G(\mathbb{CP}^1 \setminus D)^2} \text{Bun}_G(\mathbb{CP}^1 \setminus D)$ where the map $g: \text{Bun}_G^{\text{fr}}(\mathbb{CP}^1) \rightarrow \text{Bun}_G(\mathbb{CP}^1 \setminus D)$ is given by $P \mapsto (P|_{\mathbb{CP}^1 \setminus D}, q^*P|_{\mathbb{CP}^1 \setminus D})$. Consider the following commutative cube:

$$\begin{array}{ccccc}
 & q\text{-Conn}_G^{\text{fr}}(\mathbb{CP}^1, D) & \xrightarrow{f_1} & \text{Bun}_G^{\text{fr}}(\mathbb{CP}^1) & \\
 & \swarrow f_2 & \downarrow & \swarrow g_1 & \downarrow \\
 \text{Bun}_G^{\text{fr}}(\mathbb{CP}^1 \setminus D) & \xrightarrow{g_2} & \text{Bun}_G^{\text{fr}}(\mathbb{CP}^1 \setminus D)^2 & & \\
 \downarrow & \downarrow \text{res} & \downarrow r & & \downarrow \\
 & \text{Bun}_G(\mathbb{B})^k & \xrightarrow{r} & BL^+G^{2k} & \\
 & \swarrow & \downarrow & \swarrow & \\
 BLG^k & \xrightarrow{\quad} & BLG^{2k} & &
 \end{array}$$

Here the top and bottom faces are homotopy Cartesian. The front and the right faces are not homotopy Cartesian but there are canonical 1-shifted Lagrangian structures on the induced maps from the source of the face into the homotopy fiber product, i.e.

$$\begin{aligned}
 & \text{Bun}_G^{\text{fr}}(\mathbb{CP}^1) \rightarrow \text{Bun}_G^{\text{fr}}(\mathbb{CP}^1 \setminus D)^2 \times_{BLG^{2k}} BL^+G^{2k} \cong \text{Bun}_G^{\text{fr}}(\mathbb{CP}^1)^2 \\
 & \text{and } \text{Bun}_G^{\text{fr}}(\mathbb{CP}^1 \setminus D) \rightarrow \text{Bun}_G^{\text{fr}}(\mathbb{CP}^1 \setminus D)^2 \times_{BLG^{2k}} BLG^k.
 \end{aligned}$$

Call these Lagrangian structures μ_1 and μ_2 respectively. Denote the 2-shifted Lagrangian structure on the map r from Lemma 4.1 by ν . The 1-shifted Lagrangian structure that we want on the map res is then given by

$$\lambda = f_1^*(\mu_1) + f_2^*(\mu_2) + \eta$$

where η is any choice (Chris: todo: check this from e.g. Spaide: important) of potential for the difference of the pullbacks $(f_1 \circ g)^*\nu - (f_2 \circ g_2)^*\nu$. Here this difference is zero as the top square is genuinely commutative, not just up to homotopy.

This 1-shifted Lagrangian structure λ defines a quasi-isomorphism $\mathbb{T}_{q\text{-Conn}_G^{\text{fr}}(\mathbb{CP}^1, D)} \rightarrow \mathbb{L}_{\text{res}}$ where I'm writing \mathbb{L}_{res} for the dual of $\text{res}^*(\mathbb{T}_{q\text{-Conn}_G^{\text{fr}}(\mathbb{CP}^1, D)}) \rightarrow \mathbb{T}_{\text{Bun}_G(\mathbb{B})^k}[-1]$: the relative tangent complex to the map res . Write λ^{-1} for a quasi-inverse. This defines a Poisson bivector on $q\text{-Conn}_G^{\text{fr}}(\mathbb{CP}^1, D)$ where the associated map from the cotangent complex to the tangent complex is given by

$$\pi = \lambda^{-1} \circ \iota: \mathbb{L}_{q\text{-Conn}_G^{\text{fr}}(\mathbb{CP}^1, D)} \rightarrow \mathbb{L}_{\text{res}} \rightarrow \mathbb{T}_{q\text{-Conn}_G^{\text{fr}}(\mathbb{CP}^1, D)}$$

where ι is the canonical inclusion of the cotangent complex into the relative cotangent complex.

We obtain the 0-shifted symplectic structure on a Poisson leaf as the Lagrangian intersection

$$\begin{array}{ccc}
 q\text{-Conn}_G^{\text{fr}}(\mathbb{CP}^1, D, \omega^{\vee}) & \longrightarrow & q\text{-Conn}_G^{\text{fr}}(\mathbb{CP}^1, D) \\
 \downarrow & & \downarrow \\
 \prod B\Gamma_i & \longrightarrow & \text{Bun}_G(\mathbb{B})^k
 \end{array}$$

where $B\Gamma_i$ is the classifying space of the group of symmetries of the G -bundle on \mathbb{B} associated to the coweight ω_i^{\vee} . If we restrict the Poisson bivector to the leaf $q\text{-Conn}_G^{\text{fr}}(\mathbb{CP}^1, D, \omega^{\vee})$ it becomes non-degenerate and so defines a 0-shifted symplectic structure.

In order to work with this symplectic structure we'll need to give a local description as a pairing on the tangent complex. Using the description of the tangent complex in terms of hypercohomology from Section 3.1 it will suffice to describe the pairing on each summand in the Čech resolution.

Remark 4.4. From now on we'll write $\langle -, - \rangle$ to denote the residue pairing between elements of LG . That is,

$$\langle g_1, g_2 \rangle = \oint_{\mathbb{D}^\times} \kappa(g_1, g_2).$$

Lemma 4.5. As a pairing on the tangent complex at a point (P, g) the symplectic form on the moduli space of q -connections pairs α_i with β_i for $i = 1, \dots, k$. It is given by the formula

$$\omega((\{\alpha_i\}, \{\beta_i\}), (\{\alpha'_i\}, \{\beta'_i\})) = \sum_i \langle \rho_g^*(\alpha_i + \alpha_0)|_{U_i^\times}, \rho_g^*(\text{Ad}_g^*)^{-1}(\beta'_i) \rangle - \langle \rho_g^*(\alpha'_i + \alpha'_0)|_{U_i^\times}, \rho_g^*(\text{Ad}_g^*)^{-1}(\beta_i) \rangle,$$

where we use the Killing form to identify β'_i with a \mathfrak{g}^* -valued form, and where ρ_g^* is the pullback along the right multiplication by g . In this expression the sum is over $i = 1, \dots, k$ and $i = \infty$.

Proof. The formula given above is saying that the Poisson bivector – viewed as a map $\pi: \mathbb{L}_{(P,g)} \rightarrow \mathbb{T}_{(P,g)}$ – acts in degree -1 by identifying \mathfrak{g} with \mathfrak{g}^* then applying the map Ad_g^* . Recall that to describe the Poisson bivector we must locally on U_i describe the sum

$$\lambda = f_1^*(\mu_1) + f_2^*(\mu_2) + \eta,$$

where η is a 2-form of degree 0 which is closed for the internal differential. The Lagrangian structure μ_1 on the diagonal is 0, and the Lagrangian structure μ_2 is also 0 for degree reasons. Indeed, the tangent complex of $\text{Bun}_G^{\text{fr}}(\mathbb{CP}^1 \setminus D)$ is concentrated in degree -1 so there are no 2-forms of degree 1. We must therefore describe a pairing closed for the internal differential. This pairing defines a potential η as required, and therefore a Lagrangian structure.

Consider the image of our candidate pairing under the internal differential on the space of 2-forms (here modelled by the Čech differential). The result is the degree 1 pairing

$$\begin{aligned} \eta((\{\alpha_i\}, \{\beta_i\}), (\{\alpha'_i\}, \{\beta'_i\})) &= \sum_i \langle \rho_g^* \text{Ad}_g(\phi_i + \phi_0)|_{U_i^\times}, \rho_g^*(\text{Ad}_g^*)^{-1}(\beta'_i) \rangle - \langle \rho_g^* \text{Ad}_g(\phi'_i + \phi'_0)|_{U_i^\times}, \rho_g^*(\text{Ad}_g^*)^{-1}(\beta_i) \rangle \\ &= \sum_i \langle \rho_g^*(\phi_i + \phi_0)|_{U_i^\times}, \rho_g^*(\beta'_i) \rangle - \langle \rho_g^*(\phi'_i + \phi'_0)|_{U_i^\times}, \rho_g^*(\beta_i) \rangle, \end{aligned}$$

where ϕ_i is an element of $\mathfrak{g}_P(U_i)$: a degree -1 element of the Čech complex. This pairing vanishes on Čech cocycles, because if $\{\phi_i\}$ is a -1 -cocycle then $(\phi'_i + \phi'_0)|_{U_i^\times} = 0$. Our candidate pairing is therefore closed for the internal differential. \square

Proposition 4.6. The symplectic form on the moduli space of q -connections can be written in the form

$$\omega(\delta g, \delta g') = - \sum_{i=1}^d \langle b_i^L g^{-1}, b_i'^R g^{-1} \rangle - (b \leftrightarrow b') \quad (5)$$

(Vasily: I think some repairs are required here if we are to compare to the expression (7) below: 1) we should expect sandwiches between g and g^{-1} and not g^{-1} and g^{-1} 2) the summation should also include a contribution from ∞ 3) Pick either $\langle \rangle$ or κ notation for the Killing form) (Chris: other than making sure signs are correct I think the expression here works for what's below in Theorem 4.7 though I may of course be making an error. 1) We evaluate this pairing on a pair of expressions of the form $(b^L g, -gb^R)$, yielding $(b^L, -gb^R g^{-1})$, 2) doesn't the term at ∞ vanish because it can be assumed to land in $z\mathfrak{g}[[z]]$ where the residue pairing vanishes? 3) I added a remark to clarify, but by $\langle -, - \rangle$ I mean the combination of the Killing form pairing with the residue, so $\langle a, b \rangle = \oint \kappa(a, b)$.) where δg is represented on U_i by a pair $(b^L, b^R) \in \text{Ad}_g$. Here and throughout the notation “ $-(b \leftrightarrow b')$ ” denotes antisymmetrization.

Proof. Specify a representative cocycle $(\alpha_\infty, \{\alpha_i\}, \alpha_0, \beta_\infty, \{\beta_i\})$ for the first Čech cohomology group. By addition of a coboundary we can assume that $\alpha_0 = 0$, and to force α_∞ to land in $z\mathfrak{g}[[z]]$ and each α_i to land in $z^{-1}\mathfrak{g}[[z]]$. We've now fixed a representative cocycle – there's no further gauge freedom. Ignoring the factor at infinity – since after we add these coboundaries the residue pairing there vanishes – the pairing we end up with looks like

$$\begin{aligned}\omega(\delta g, \delta g') &= \sum_{i=1}^d \langle \rho_g^*(\alpha_i + \alpha_0)|_{U_i^\times}, \rho_g^*(\text{Ad}_g^*)^{-1}(\beta'_i) \rangle - ((\alpha, \beta) \leftrightarrow (\alpha', \beta')) \\ &= \sum_{i=1}^d \langle \rho_g^*(\alpha_i + \alpha_0)|_{U_i^\times}, \rho_g^*(\text{Ad}_g^*)^{-1}(\text{Ad}_g^{-1}(\alpha'_i - \alpha'_0)) \rangle - (\alpha \leftrightarrow \alpha') \\ &= \sum_{i=1}^d \langle \rho_g^*(\alpha_i)|_{U_i^\times}, \rho_g^*(\text{Ad}_g^*)^{-1}(\text{Ad}_g^{-1}(\alpha'_i)|_{U_i^\times}) \rangle - (\alpha \leftrightarrow \alpha').\end{aligned}$$

We can compute the composite operator $(\text{Ad}_g^*)^{-1}\text{Ad}_g^{-1}$ on $\text{ad}(P)(U_i^\times)$. It's given by the inverse of the operator Ad_gAd_g^* which sends a pair $(b^L, b^R) \in \text{ad}(P)(U_i^\times)$ to $(b^L - b^R, b^R - b^L) \sim (b^L + g^{-1}b^Lg, b^R - gb^Rg^{-1})$. Denote this by $((1 + A_g)b^L, (1 - A_{g^{-1}})b^R)$. We can describe the inverse using the expansion $(1 + A_g)^{-1} = 1 - A_g + A_g^2 - \dots$. After applying our gauge transformation above the remaining degree of freedom in α_i is its z^{-1} term. Each time we apply A it raises the order in z of this term by one, so only the linear summand $-A_g$ of $(1 + A_g)^{-1}$ contributes to the residue pairing. In the pairing we need to use the invariant pairing on $\mathfrak{g}((z)) \oplus \mathfrak{g}((z))$ that vanishes on the subalgebra spanned by $(X, A_g(X))$, so we take the difference of the residue / Killing form pairings on the two summands (Chris: scaled by 1/2: why this normalization?).

$$\begin{aligned}\omega(\delta g, \delta g') &= \frac{1}{2} \sum_{i=1}^d -\langle \rho_g^*A_gb_i^L, \rho_g^*b_i'^R \rangle - \langle \rho_g^*b_i^L, \rho_g^*A_{g^{-1}}b_i'^R \rangle - (b \leftrightarrow b') \\ &= -\sum_{i=1}^d \langle b_i^Lg^{-1}, b_i'^Rg^{-1} \rangle - (b \leftrightarrow b')\end{aligned}$$

using cyclic invariant to identify the two terms. □

We can now establish our main result.

Theorem 4.7. The symplectic structure on $\text{Mon}_G^{\text{fr}}(\mathbb{CP}^1 \times S^1, D \times \{0\}, \omega^\vee)$ and the pullback of the symplectic structure on $\text{mHiggs}_G^{\text{ps, fr}}(\mathbb{CP}^1, D, \omega^\vee)$ under H coincide.

Proof. This is straightforward now, by combining the two local descriptions of the symplectic structure on monopoles and multiplicative Higgs bundles on a neighbourhood of a puncture with the description in Proposition 3.16. So, let's begin by taking the symplectic form ω_{mHiggs} on the moduli space $\text{mHiggs}_G^{\text{ps, fr}}(\mathbb{CP}^1, D, \omega^\vee)$ and evaluating it at the image under dH of two elements $(\alpha_i, \alpha'_i) = (d_{\text{mon}}(b_i), d_{\text{mon}}(b'_i))$. Let us write $b_i(0) = b_i^L$ and $b_i(2\pi) = b_i^R$ for brevity. Likewise for consistency with the calculations above let us denote the image $H(\mathcal{A})$ under the holonomy map by g .

According to Proposition 3.16 we have

$$\begin{aligned}\omega_{\text{mHiggs}}(dH(\alpha_i), dH(\alpha'_i)) &= \omega_{\text{mHiggs}}(dH(d_{\text{mon}}(b_i)), dH(d_{\text{mon}}(b'_i))) \\ &= \omega_{\text{mHiggs}}(b_i^Rg - gb_i^L, b_i'^Rg - gb_i'^L).\end{aligned}$$

We can write this in terms of a pairing on the bundle $\text{ad}(g)$. Represent the class $b_i^Rg - gb_i^L$ by the pair $(b_i^Rg, -gb_i^L)$. Then applying the description of ω_{mHiggs} provided by Proposition 4.6 we have

$$\begin{aligned}\omega_{\text{mHiggs}}(dH(\alpha_i), dH(\alpha'_i)) &= \omega_{\text{mHiggs}}((b_i^Rg, -gb_i^L), (b_i'^Rg, -gb_i'^L)) \\ &= \langle b_i^R, gb_i'^Lg^{-1} \rangle - (b \leftrightarrow b')\end{aligned}$$

where the remaining terms are killed by the antisymmetrization.

On the other hand, we can evaluate the pairing $\omega_{\text{Mon}}(\alpha_i, \alpha'_i) = \omega_{\text{Mon}}(d_{\text{mon}}(b_i), d_{\text{mon}}(b'_i))$ via integration by parts. We'll compute the symplectic pairing for the full Čech complex, but it will split into a sum over open sets U_i . The result is that (Vasily: the expression (3) expands as) (Vasily: 1. let us unify the notations for the Higgs field in the equations above and the previous sections: somewhere holonomy or the Higgs field is called $H(\mathcal{A})$, somewhere ϕ , somewhere g . I would favour the notation g . 2. Let us introduce somewhere before the notation $b^L = b(0), b^R = b(2\pi)$ to make the equations more readable, and break the multiline computation into separate lines and insert between the lines the explanations of what we are doing) (Chris: made some notational changes following your suggestions.)

$$\begin{aligned}\omega_{\text{Mon}}(\{\alpha_i\}, \{\alpha'_i\}) &= \sum_{i=0, \dots, k, \infty} \int_{U_i} \kappa(d_{\text{mon}}(b_i) \wedge d_{\text{mon}}(b'_i)) dz - (b \leftrightarrow b') \\ &= \sum_{i=1, \dots, k, \infty} \int_{\partial \mathbb{D}_i \times [0, 2\pi]} \kappa(b_i - b_0, d_{\text{mon}}(b'_i)) dz - (b \leftrightarrow b')\end{aligned}$$

using here the fact that $d_{\text{mon}} b_i = d_{\text{mon}} b_0$ on $U_i \cap U_0$ and that $d_{\text{mon}} b_i(t) = 0$ when $t = 0$ or 2π . By Stokes' theorem we then have

$$\begin{aligned}\omega_{\text{Mon}}(\{\alpha_i\}, \{\alpha'_i\}) &= \sum_{i=1, \dots, k, \infty} \oint_{\partial \mathbb{D}_i} \kappa(b_i^R - b_0^R, b_i'^R) - \kappa(b_i^L - b_0^L, b_i'^L) - (b \leftrightarrow b') \\ &= \sum_{i=1, \dots, k, \infty} \oint_{\partial \mathbb{D}_i} -\kappa(b_0^R, b_i'^R) + \kappa(b_0^L, b_i'^L) - (b \leftrightarrow b').\end{aligned}$$

Pick out the summand corresponding to U_i . Choose our potentials so that on the boundary $\partial \mathbb{D}_i$ we have $b_0^L = b_0'^L = 0$. We can do this by setting $b^R = \delta g g^{-1}$ since g is non-singular on U_0 . This choice means that on $U_0 \cap U_i$, since $\delta g = b_i^R g - g b_i^L = b_0^R g$ we can make the identification $b_0^R = b_i^R - g b_i^L g^{-1}$. Therefore

$$\omega_{\text{Mon}}(\{\alpha_i\}, \{\alpha'_i\}) = \sum_{i=1, \dots, k, \infty} \langle (g b_i^L g^{-1}, b_i'^R) \rangle - (b \leftrightarrow b')$$

agreeing with the expression coming from mHiggs. (Chris: indeed there's a sign error somewhere which I haven't tracked down yet.)

(Vasily: Instead, I would suggest that here we choose the potentials such that $b_0^L = 0$ and $b_0'^L = 0$ everywhere on U_0 , and in particular on the boundary $\partial \mathbb{D}_i$. There is no obstruction to this choice because for any variation of the holonomy $\delta g = b^R g - g b^L$ with the choice $b^L = 0$ we set $b^R = \delta g g^{-1}$ which is a holomorphic section on U_0 for the reason that U_0 does not contain any singularities and because of the Bogomolny equation. Given this choice the last line becomes

$$\sum_{i=1, \dots, k, \infty} \oint_{\partial \mathbb{D}_i} -\kappa(b_0^R, b_i'^R) - (b \leftrightarrow b') \quad (6)$$

Next, from $\delta g = b_i^R g - g b_i^L = b_0^R g$ on $\partial \mathbb{D}_i = U_0 \cap U_i$ we find $b_0^R = b_i^R - g b_i^L g^{-1}$ and plug in the above line. Because of the antisymmetrization $b \leftrightarrow b'$ the term containing $b_i^R b_i'^R$ vanishes and we are left with

$$\sum_{i=1, \dots, k, \infty} \oint_{\partial \mathbb{D}_i} \kappa(g b_i^L g^{-1}, b_i'^R) - (b \leftrightarrow b') \quad (7)$$

) (Chris: modified)

(Vasily: I think the minus sign is missed if compared to (7)) □

Remark 4.8. (Chris: connection to Shapiro's symplectic leaves in [Sha16]. The claim is that taking the Taylor expansion at infinity defines a Poisson map to $\tilde{G} \subseteq G_1[[z^{-1}]]$ sending symplectic leaves to (discrete) unions of symplectic leaves. One can do the classification more completely in the type A case.)

5 Hyperkähler Structures

(Vasily: As of July 19, 2018 I have not yet passed this and following sections)

The results of the previous section imply that the symplectic structure on $\mathrm{mHiggs}_G^{\mathrm{ps},\mathrm{fr}}(\mathbb{CP}^1, D, \omega^\vee)$ extends canonically to a hyperkähler structure. In this section we'll compare the twistor rotation with the deformation to the moduli space of q -connections. (Chris: todo: expand on this)

Proposition 5.1. (Chris: Monopoles in the large R (Gaiotto) limit – twistor rotation and ε -deformation coincide.)

Proof. □

Corollary 5.2. If we equip $\mathrm{mHiggs}_G^{\mathrm{ps},\mathrm{fr}}(\mathbb{CP}^1, D, \omega^\vee)$ with the complex structure at the twistor parameter $\eta \in \mathbb{C}$, the resulting complex manifold is isomorphic to $\varepsilon\text{-Conn}_G^{\mathrm{ps},\mathrm{fr}}(\mathbb{CP}^1, D, \omega^\vee)$.

(Chris: comment on the holomorphic symplectic structure vs the one discussed above.)

(Chris: comment on the example of the Nahm transform.)

6 Quantization and Yangians

(Chris: We're still discussing how this story goes.)

7 Duality

(Chris: For a first draft I'm inserting some notes I prepared for a talk. Todo: talk about motivation and the content we discussed regarding non-simply-laced groups)

Pseudo-Conjecture 7.1 (Multiplicative Geometric Langlands). Let G be a Langlands self-dual group. There is an equivalence of categories

$$\mathrm{A}\text{-Branes}_{q^{-1}}(\mathrm{mHiggs}_G(C, D, \omega^\vee)) \cong \mathrm{B}\text{-Branes}(q\text{-Conn}_G(C, D, \omega^\vee))$$

where the category on the right-hand side depends on the value q .

What does this mean, and are there situations in which we can make it precise? We'll discuss three examples where we can say something more concrete. In each case, by “B-branes” we'll just mean the category $\mathrm{Coh}(q\text{-Conn}_G(C, D, \omega^\vee))$ of coherent sheaves. By “A-branes” we'll mean some version of q^\vee -difference connections on the stack $\mathrm{Bun}_G(C)$.

Remark 7.2. This equivalence is supposed to interchange objects corresponding to branes of opers on the two sides, and introduce an analogue of the Feigin-Frenkel isomorphism between deformed W-algebras (see [FRSTS98, STSS98]). This isomorphism only holds for self-dual groups, which motivates the restriction to the self-dual case here.

7.1 The Abelian Case

Suppose $G = \mathrm{GL}(1)$ (more generally we could consider a higher rank abelian gauge group). In general for an abelian group the moduli spaces we've defined are trivial – for instance the rational and trigonometric spaces are always

discrete. However there is one interesting non-trivial example: the elliptic case. For simplicity let's consider the abelian situation with $D = \emptyset$: the case with no punctures.

Definition 7.3. A q -difference module on a variety X with automorphism q is a module for the sheaf $\Delta_{q,X}$ of non-commutative rings generated by \mathcal{O}_X and an invertible generator Φ with the relation $\Phi \cdot f = q^*(f) \cdot \Phi$. Write $\text{Diff}_q(X)$ for the category of q -difference modules on X .

In the abelian case the space $q\text{-Conn}_{\text{GL}(1)}(E)$ is actually a stack, but one can split off the stacky part to define difference modules on it. Indeed, for any q one can write

$$\text{Bun}_{\text{GL}(1)}(E) \cong \text{BGL}(1) \times \mathbb{Z} \times E^\vee$$

and so

$$q\text{-Conn}_{\text{GL}(1)}(E) \cong \text{BGL}(1) \times \mathbb{Z} \times (E^\vee \times_q \mathbb{C}^\times)$$

which means one can define difference modules on these stacks associated to an automorphism of E^\vee or $E^\vee \times_q \mathbb{C}^\times$ respectively.

Conjecture 7.4. There is an equivalence of categories for any $q \in \mathbb{CP}^1$

$$\text{Diff}_q(\text{Bun}_{\text{GL}(1)}(E)) \cong \text{Coh}(q^{-1}\text{-Conn}_{\text{GL}(1)}(E)).$$

In this abelian case we can go even farther and make a more sensitive 2-parameter version of the conjecture.

Conjecture 7.5. There is an equivalence of categories for any $q_1, q_2 \in \mathbb{CP}^1$

$$\text{Diff}_{q_1}(q_2\text{-Conn}_{\text{GL}(1)}(E)) \cong \text{Diff}_{q_2^{-1}}(q_1^{-1}\text{-Conn}_{\text{GL}(1)}(E))$$

where q_1 is the automorphism of $E^\vee \times_{q_2} \mathbb{C}^\times$ acting fiberwise over each point of \mathbb{C}^\times .

This conjecture should be provable using the same techniques as the ordinary geometric Langlands correspondence in the abelian case, i.e. by a (quantum) twisted Fourier-Mukai transform (as constructed by Polishchuk and Rothstein [PR01]).

7.2 The Classical Case

Now, let's consider the limit $q \rightarrow 0$. This will give a conjectural statement involving coherent sheaves on both sides analogous to the classical limit of the geometric Langlands conjecture as conjectured by Donagi and Pantev [DP10]. The existence of an equivalence isn't so interesting in the self-dual case (where both sides are the same), but the classical multiplicative Langlands functor should be an *interesting* non-trivial equivalence. For example we can make the following conjecture

Conjecture 7.6. Let G be a Langlands self-dual group and let E be an elliptic curve. There is an automorphism of categories (for the rational, trigonometric and elliptic moduli spaces)

$$F: \text{Coh}(\text{mHiggs}_G(E)) \cong \text{Coh}(\text{mHiggs}_G(E))$$

so that the following square commutes:

$$\begin{array}{ccc} \text{Coh}(\text{mHiggs}_T(E)) & \xrightarrow{\text{FM}} & \text{Coh}(\text{mHiggs}_T(E)) \\ p_* q^! \downarrow & & \downarrow p_* q^! \\ \text{Coh}(\text{mHiggs}_G(E)) & \xrightarrow{F} & \text{Coh}(\text{mHiggs}_G(E)). \end{array}$$

Here we're using the natural morphisms $p: \text{mHiggs}_B(E) \rightarrow \text{mHiggs}_G(E)$ and $q: \text{mHiggs}_B(E) \rightarrow \text{mHiggs}_T(E)$, and FM is the Fourier-Mukai transform.

It ought to be possible to state something a bit stronger that includes singularities in this auto-duality.

7.3 The Rational Type A Case

There's one more example where we can say something precise, and even draw a connection to the ordinary geometric Langlands correspondence. We already mentioned the Nahm transform in the previous section: in the case where $G = \mathrm{GL}(n)$ and C is \mathbb{C} (where as usual we fix framing data at infinity) the Nahm transform identifies multiplicative Higgs bundles of degree k with *ordinary* Higgs bundles on \mathbb{CP}^1 for the group $\mathrm{GL}(k)$ with $n + 2$ tame singularities (with appropriate fixed locations and residues).

Claim. Under the Nahm transform, Pseudo-Conjecture 7.1 in the rational case for the group $\mathrm{GL}(n)$ becomes the ordinary geometric Langlands conjecture on \mathbb{CP}^1 with tame ramification.

References

- [AB01] Boudjemaa Anchouche and Indranil Biswas. Einstein-Hermitian connections on polystable principal bundles over a compact Kähler manifold. *Amer. J. Math.*, 123(2):207–228, 2001.
- [Bot95] Francesco Bottacin. Symplectic geometry on moduli spaces of stable pairs. *Ann. Sci. École Norm. Sup. (4)*, 28(4):391–433, 1995.
- [CH10] Benoît Charbonneau and Jacques Hurtubise. Singular Hermitian-Einstein monopoles on the product of a circle and a Riemann surface. *Int. Math. Res. Not.*, 2011(1):175–216, 2010.
- [CK98] Sergey Cherkis and Anton Kapustin. Singular monopoles and supersymmetric gauge theories in three dimensions. *Nuclear Physics B*, 525(1-2):215–234, 1998.
- [CK01] Sergey Cherkis and Anton Kapustin. Nahm transform for periodic monopoles and $\mathcal{N} = 2$ super Yang-Mills theory. *Communications in Mathematical Physics*, 218(2):333–371, 2001.
- [CK02] Sergey Cherkis and Anton Kapustin. Hyper-Kähler metrics from periodic monopoles. *Physical Review D*, 65(8):084015, 2002.
- [CPT⁺17] Damien Calaque, Tony Pantev, Bertrand Toën, Michel Vaquié, and Gabriele Vezzosi. Shifted Poisson structures and deformation quantization. *Journal of Topology*, 10(2):483–584, 2017.
- [DP10] Ron Donagi and Tony Pantev. Lectures on the geometric Langlands conjecture and non-abelian Hodge theory. *Second International School on Geometry and Physics Geometric Langlands and Gauge Theory*, page 129, 2010.
- [EV98] Pavel Etingof and Alexander Varchenko. Geometry and classification of solutions of the classical dynamical Yang-Baxter equation. *Comm. Math. Phys.*, 192(1):77–120, 1998.
- [Fos16] Lorenzo Foscolo. Deformation theory of periodic monopoles (with singularities). *Comm. Math. Phys.*, 341(1):351–390, 2016.
- [FRSTS98] Edward Frenkel, Nicolai Reshetikhin, and Michel Semenov-Tian-Shansky. Drinfeld-Sokolov reduction for difference operators and deformations of-Walgebras *i.* the case of Virasoro algebra. *Communications in mathematical physics*, 192(3):605–629, 1998.
- [Gro57] Alexander Grothendieck. Sur la classification des fibrés holomorphes sur la sphère de Riemann. *Amer. J. Math.*, 79:121–138, 1957.
- [Hen15] Benjamin Hennion. *Formal loops and tangent Lie algebras*. PhD thesis, Université de Montpellier, 2015.
- [HM02] Jacques Hurtubise and Eyal Markman. Elliptic Sklyanin integrable systems for arbitrary reductive groups. *Advances in Theoretical and Mathematical Physics*, 6(5):873–978, 2002.
- [Moc17] Takuro Mochizuki. Periodic monopoles and difference modules. *arXiv preprint arXiv:1712.08981*, 2017.
- [MS16] Valerio Melani and Pavel Safronov. Derived coisotropic structures. *arXiv preprint arXiv:1608.01482*, 2016.
- [MS17] Valerio Melani and Pavel Safronov. Derived coisotropic structures II: stacks and quantization. *arXiv preprint arXiv:1704.03201*, 2017.

- [PR01] Alexander Polishchuk and Mitchell Rothstein. Fourier transform for D -algebras, I. *Duke Mathematical Journal*, 109(1):123–146, 2001.
- [PTVV13] Tony Pantev, Bertrand Toën, Michel Vaquié, and Gabriele Vezzosi. Shifted symplectic structures. *Publications mathématiques de l’IHÉS*, 117(1):271–328, 2013.
- [Sha16] Alexander Shapiro. Poisson geometry of monic matrix polynomials. *Int. Math. Res. Not.*, (17):5427–5453, 2016.
- [Smi15] Benjamin H Smith. Singular G -monopoles on $S^1 \times \Sigma$. *arXiv preprint arXiv:1501.03337*, 2015.
- [Spa16] Theodore Spaide. Shifted symplectic and Poisson structures on spaces of framed maps. *arXiv preprint arXiv:1607.03807*, 2016.
- [STSS98] Michel Semenov-Tian-Shansky and Alexey Sevostyanov. Drinfeld–Sokolov reduction for difference operators and deformations of W -algebras II. the general semisimple case. *Communications in mathematical physics*, 192(3):631–647, 1998.
- [Zhu17] Xinwen Zhu. An introduction to affine Grassmannians and the geometric Satake equivalence. In *Geometry of moduli spaces and representation theory*, volume 24 of *IAS/Park City Math. Ser.*, pages 59–154. Amer. Math. Soc., Providence, RI, 2017.

INSTITUT DES HAUTES ÉTUDES SCIENTIFIQUES
35 ROUTE DE CHARTRES, BURES-SUR-YVETTE, 91440, FRANCE
`celliot@ihes.fr`
`pestun@ihes.fr`