

Untitled Project

Christopher Kirwan 17325310
Project Supervisor: Sinead Ryan

December 8, 2020

Contents

1	Introduction	2
1.1	The Need For Spin Models	2
1.2	Defintions for the Spin $O(n)$ (Symmetric) Model	3

Chapter 1

Introduction

1.1 The Need For Spin Models

The study of statistical mechanics arose due to the need to connect the physical descriptions of large particle systems in the macroscopic and microscopic realms. This is done by bridging thermodynamic quantities (macroscopic), such as temperature, with microscopic observables that fluctuate about some average. This is achieved using statistical methods and probability theory. Applying the methods developed by statistical mechanics to various mathematical models has given physicists enormous insight into various physical phenomena and gives justification to study these models in detail.

We begin by constructing a physical model, using a substance whose individual atoms are arranged in a regular crystalline lattice. Furthermore, suppose that each atom in the lattice has a magnetic moment which we call its spin. In this picture, we also assume that these moments tend to align with their neighbours¹ and an external magnetic field H . We introduce a measurement parameter called the magnetization, which is simply the global average of these spins.




Figure 1.1

Varying the external magnetic field, we can observe two distinct behaviours around $H = 0$, called paramagnetic and ferromagnetic behaviour, first observed by Pierre Curie in 1895. In the first case, as $H \rightarrow 0$, the global order (ie: spin alignment) is lost and the magnetization tends to zero. In the ferromagnetic case, the local spin interactions are strong enough that the substance maintains a global non-zero net magnetization. The value of this magnetization depends on the direction in which the magnetic field approaches 0. Thus, a ferromagnet displays a *spontaneous magnetization* $\pm M$ at $H = 0$. This is a discontinuity in the magnetization, which corresponds to a *first order phase transition*.

¹this assumption will be made concrete later

The observations by Curie also established that a temperature dependent transition occurs in ferromagnetic materials, in which the material changes into a paramagnetic material. This transition occurs at a well-defined temperature called the Curie Temperature. A goal of 20th century thermodynamics was to be able to describe this phase transition using the framework of statistical mechanics.

In 1920, Wilhelm Lenz introduced what is now called the Ising Model (a term coined by Rudolph Peierls) to help understand that phase transition. The one-dimensional case was solved by

1.2 Defintions for the Spin $O(n)$ (Symmetric) Model

We begin the discusiion of the model by defining the system on which the model "lives".

Definition 1.2.1. Let $d \geq 1$ and let $G = (V(G), E(G))$ be a finite, non-oriented graph. The set of vertices Λ is defined to be a subset of \mathbb{Z}^d (that is, $V(G) = \Lambda$, with the coordinates $i = (i_1, \dots, i_d) \in \Lambda$ being integers). We also define the egdes of the graph $E(G)$ to be between the *nearest neighbours* of Λ . That is, pairs (i, j) with $\|i - j\|_1 = 1$. We denote nearest neighbours as $\langle i, j \rangle$.

Throughout the remainder of this text, we shall call the graph $G = (\Lambda, E)$ a *lattice* and simply refer to it as Λ . We also remark that Λ can also be \mathbb{Z}^d itself. This is referred to as the *thermodynamic limit*. The need for this limit will be made apparent later.

Fixing $n \in \mathbb{N}$, we have the *single spin configuration* as

$$\Omega_0 = \{\sigma \in \mathbb{R}^n \text{ such that } \|\sigma\|_2 = 1\} \equiv \mathbb{S}^{n-1}$$

Following from this, we have a *microstate*, often referred to as a *spin configuration*, on Λ as

$$\Omega_\Lambda = \Omega_0^\Lambda$$

and in the thermodynamic limit $\Omega = \Omega_0^{\mathbb{Z}^d}$

For each $i \in \Lambda$, we call the random variable $\sigma_i : \Omega_\Lambda \rightarrow \mathbb{S}^{n-1}$ assigned to each point the lattice as the *spin*. We add two special restrictions onto these spins:

- That the spins themselves interact with their nearest neighbours
- This same interaction is invariant under a simultaneous rotation of the entire configuration²

With these restrictions, we can then assume that only a function of the inner/scalar product of the interacting spins contribute to the total energy of the system. We therefore arrive at what is *probably* the most important definition of the entire report

Definition 1.2.2. Let $U : [-1, 1] \rightarrow \mathbb{R}$ (usually referred to as the *potential function*) and \mathcal{H}_Λ be the Hamiltonian of an $O(n)$ -symmetric model, over a lattice Λ . We thus have

$$\mathcal{H}_\Lambda = \sum_{\langle i, j \rangle} U(\langle \sigma_i, \sigma_j \rangle) \quad (1.1)$$

In particular, if we have $f(x) = -x$, we then have the definiton of the Hamiltonian for the $O(n)$ (or n -vector) model

$$H_\Lambda = - \sum_{\langle i, j \rangle} \langle \sigma_i, \sigma_j \rangle \quad (1.2)$$

²The discussion of an external magnetic field will occur with the Ising Model

We can now plug in different values of n to get different models

- $n = 1$ gives the *Ising Model*, with spins taking values in $\{\pm 1\}$. The discussion on this model is given below
- $n = 2$, the spins lie along the circle with radius 1 (hence $O(2)$ symmetric). This is called the *XY Model* and is also central to the discussion.
- $n = 3, n \rightarrow \infty$ are called the *Heisenberg Model* and *Berlin-Kac Spherical Model*. $n = 4$ can be used as a "toy" model of the Higgs Sector.

Definition 1.2.3. At inverse temperature $\beta \in [0, \infty)$, configurations are randomly chosen from the probability measure μ defined as

$$d\mu = \frac{1}{Z} \exp(\beta H(\sigma_i, \sigma_j)) d\sigma \quad (1.3)$$

where Z is the normalization constant called the *partition function*, given by

$$Z = \int_{\Omega_\Lambda} \exp(\beta H(\sigma_i, \sigma_j)) d\sigma \quad (1.4)$$