

Survival outcomes with oral azacitidine maintenance in patients with acute myeloid leukemia in remission by receipt of initial chemotherapy: subgroup analyses from the phase 3 QUAZAR AML-001 trial

by Andrew H. Wei, Gail J. Roboz, Hervé Dombret, Hartmut Döhner, Andre C. Schuh, Pau Montesinos, Dominik Selleslag, Sergey N. Bondarenko, Thomas Prebet, Yinzhi Lai, Barry Skikne, C. L. Beach, and Farhad Ravandi

Received: October 25, 2022.

Accepted: March 16, 2023.

Citation: Andrew H. Wei, Gail J. Roboz, Hervé Dombret, Hartmut Döhner, Andre C. Schuh, Pau Montesinos, Dominik Selleslag, Sergey N. Bondarenko, Thomas Prebet, Yinzhi Lai, Barry Skikne, C. L. Beach, and Farhad Ravandi. Survival outcomes with oral azacitidine maintenance in patients with acute myeloid leukemia in remission by receipt of initial chemotherapy: subgroup analyses from the phase 3 QUAZAR AML-001 trial.

Haematologica. 2023 Mar 23. doi: 10.3324/haematol.2022.282296 [Epub ahead of print]

Publisher's Disclaimer.

E-publishing ahead of print is increasingly important for the rapid dissemination of science. Haematologica is, therefore, E-publishing PDF files of an early version of manuscripts that have completed a regular peer review and have been accepted for publication. E-publishing of this PDF file has been approved by the authors. After having E-published Ahead of Print, manuscripts will then undergo technical and English editing, typesetting, proof correction and be presented for the authors' final approval; the final version of the manuscript will then appear in a regular issue of the journal. All legal disclaimers that apply to the journal also pertain to this production process.

Survival outcomes with oral azacitidine maintenance in patients with acute myeloid leukemia in remission by receipt of initial chemotherapy: subgroup analyses from the phase 3 QUAZAR AML-001 trial

Andrew H. Wei,^{1,2*} Gail J. Roboz,^{3,4} Hervé Dombret,^{5,6} Hartmut Döhner,⁷ Andre C. Schuh,⁸ Pau Montesinos,⁹ Dominik Selleslag,¹⁰ Sergey N. Bondarenko,¹¹ Thomas Prebet,¹² Yinzhi Lai,¹² Barry Skikne,^{12,13†} C.L. Beach^{12†} and Farhad Ravandi¹⁴

¹Department of Clinical Haematology, The Alfred Hospital, Melbourne, Australia;

²Australian Centre for Blood Diseases, Monash University, Melbourne, Australia;

³Weill Cornell Medicine, New York, NY, USA;

⁴New York Presbyterian Hospital, New York, NY, USA;

⁵Hematology, Hôpital Saint-Louis, Assistance Publique – Hôpitaux de Paris (AP-HP), Paris, France;

⁶Institut de Recherche Saint-Louis, Université Paris Cité, Paris, France;

⁷Department of Internal Medicine III, Ulm University Hospital, Ulm, Germany;

⁸Princess Margaret Cancer Centre, Toronto, ON, Canada;

⁹Hospital Universitario y Politécnico La Fe, Valencia, Spain;

¹⁰AZ Sint-Jan Brugge-Oostende AV, Bruges, Belgium;

¹¹RM Gorbacheva Research Institute, Pavlov University, St. Petersburg, Russia;

¹²Bristol Myers Squibb, Summit, NJ, USA;

¹³University of Kansas Cancer Center, Kansas City, KS, USA;

¹⁴Department of Leukemia, The University of Texas MD Anderson Cancer Center, Houston, TX, USA.

*Currently at the Department of Clinical Haematology, Peter MacCallum Cancer Centre and Royal Melbourne Hospital, Melbourne, Australia; †Employed by Bristol Myers Squibb at the time the study was conducted.

Running head: Survival by prior chemotherapy in QUAZAR AML-001

Corresponding author: Andrew H. Wei, andrew.wei@petermac.org

Data sharing statement: BMS policy on data sharing may be found at

<https://www.bms.com/researchers-and-partners/independent-research/data-sharing-request-process.html>

Word count: Abstract, N/A; Main text, 1579

Number of main-text tables: 1

Number of main-text figures: 2

Number of supplemental files: 2 (1 containing 2 supplemental tables, 1 CONSORT diagram)

Trial registration: ClinicalTrials.gov identifier: NCT01757535

Acknowledgments: The authors would like to thank the patients, families, investigators, staff, and clinical study teams who participated in the QUAZAR AML-001 trial. All authors contributed to and approved the manuscript for submission. Writing and editorial support was provided by Korin Albert, PhD, of Excerpta Medica, funded by Bristol Myers Squibb.

Authors' contributions: The sponsors collected and analyzed data in conjunction with all authors. AHW wrote the first draft of the manuscript. All authors revised the manuscript and reviewed and approved the final version for submission.

Main text:

Oral azacitidine (Oral-AZA) is a hypomethylating agent approved for the treatment of adult patients with acute myeloid leukemia (AML) in first remission after intensive chemotherapy (IC).^{1,2} In the phase 3, randomized, double-blind, placebo-controlled QUAZAR AML-001 trial, Oral-AZA significantly prolonged relapse-free survival (RFS) and overall survival (OS) compared with placebo in patients with AML in first complete remission (CR) or CR with incomplete blood count recovery (CRi) after IC (induction ± consolidation) who were not candidates for hematopoietic stem cell transplantation (HSCT).³ The primary goal of QUAZAR AML-001 was to evaluate the effect of maintenance therapy with Oral-AZA for patients in remission after induction. While there were no protocol-specified criteria regarding prior chemotherapy used before study entry, including the use or number of consolidation cycles received, it is of clinical interest to assess whether the amount of pre-study chemotherapy may have influenced survival outcomes in this trial. Here, we present RFS and OS outcomes in patient subgroups defined by the use of consolidation and number of chemotherapy courses received prior to study entry.

IC is the cornerstone of initial AML therapy for patients fit enough to receive it, and most patients achieve CR with induction. Once in remission, patients may receive subsequent consolidation chemotherapy, but the optimal number of consolidation cycles is not well-defined, especially for older patients. After IC, the primary therapeutic goals for patients with AML in remission who are not eligible for HSCT are to delay relapse and prolong survival. Until Oral-AZA, no agent studied in the remission maintenance setting had significantly prolonged both RFS and OS.¹⁻⁸

Study design and key eligibility criteria of QUAZAR AML-001 have been reported in detail elsewhere.³ Briefly, eligible patients were aged ≥55 years with newly diagnosed AML in first remission after IC, had intermediate- or poor-risk cytogenetics (NCCN 2011 criteria⁹) and an Eastern Cooperative Oncology Group performance status (ECOG PS) score ≤3, and were HSCT-ineligible. Induction and consolidation regimens were administered at the discretion of the treating physician before study screening; trial eligibility was not contingent on the use of consolidation chemotherapy or amount of consolidation cycles received, but patients must have been screened for eligibility within 4 months of achieving initial CR/CRi during induction. Eligible patients were randomized 1:1 to Oral-AZA 300-mg or placebo once daily for 14 days per 28-day treatment cycle. Measurable residual disease (MRD) was assessed centrally via multiparameter flow cytometry, with a positivity threshold of 0.1% in the bone marrow for aberrant cells (different from normal or leukemia aberrant phenotype).

The primary trial endpoint was OS, defined as the time from randomization until death, and the key secondary endpoint was RFS, the time from randomization until relapse or death. Comparisons of OS and RFS between Oral-AZA and placebo within patient subgroups defined by use of consolidation therapy after induction (yes or no) were prospective exploratory endpoints in the trial protocol. Additional post hoc analyses were performed to assess survival outcomes in subgroups defined by the number of consolidation courses received (0, 1, or ≥2) and total number of induction and consolidation cycles. Induction courses were defined as AML-directed chemotherapy regimens administered prior to the date of first CR/CRi recorded on the electronic case report form and consolidation regimens were those given after that date.

Survival endpoints were estimated using Kaplan-Meier methods and compared between treatment arms using hazard ratios (HRs) and 95% confidence intervals (CIs) from stratified Cox proportional hazards models and *P* values from stratified log-rank tests. The post hoc survival analyses by number of consolidation cycles and total cycles of induction and consolidation were not sufficiently powered to determine statistically significant differences within or between treatment arms, precluding meaningful

interpretation of *P* values; HR point estimates and 95% CIs in these subgroups are provided for informational purposes only. The data cutoff was performed in July 2019.

The trial enrolled 472 patients (Oral-AZA 238, placebo 234) (Figure 1). Prior to enrollment, the most common agents used for induction and consolidation were cytarabine (99% and 80%, respectively), idarubicin (55% and 20%), and daunorubicin (33% and 8%); use of these agents was similar between the Oral-AZA and placebo arms. Most patients (80% [378/472]) received consolidation after induction, and use of consolidation was similar between treatment arms (Oral-AZA 78% [186/238], placebo 82% [192/234]) (Figure 1). Nearly half of patients in the Oral-AZA (*n*=110 [46%]) and placebo (*n*=102 [44%]) arms received 1 prior consolidation, and 32% (*n*=76) and 38% (*n*=90) of patients, respectively, received ≥2 prior consolidation cycles. The remaining 20% of patients (*n*=94) did not receive consolidation, including 52 patients (22%) in the Oral-AZA arm and 42 (18%) in the placebo arm. Baseline characteristics were generally similar among consolidation-defined subgroups within and between treatment arms (Supplementary Table S1). In both arms, patients who did not receive consolidation tended to be older than those who did. Rate of measurable residual disease (MRD) negativity at screening was similar between consolidation-defined cohorts within the Oral-AZA arm, whereas in the placebo arm, a larger proportion of patients who received consolidation were MRD-negative compared with those who did not (50% vs 36%, respectively).

Oral-AZA significantly prolonged both RFS and OS from the time of randomization compared with placebo, regardless of whether patients received consolidation prior to study entry. For patients who did not receive consolidation, median RFS was prolonged with Oral-AZA by 4.5 months vs placebo (median 8.4 vs 3.9 months, respectively; HR 0.58 [95%CI 0.36–0.94]; *P*=0.0258) and the estimated 1-year RFS rate was 18.7% higher with Oral-AZA (40.8% vs 22.0%) (Figure 2A and Table 1). Oral-AZA also prolonged median OS in this subgroup by approximately 12 months compared with placebo (median 23.3 vs 10.9 months, respectively; HR 0.54 [95%CI 0.33–0.87]; *P*=0.0103) and improved 1-year survival rate by 30.7% (71.2% vs 40.5%) (Figure 2B and Table 1). For patients who did receive consolidation following initial induction, median RFS was prolonged more than two-fold with Oral-AZA vs placebo—10.2 vs 5.0 months, respectively (HR 0.67 [95%CI 0.53–0.85]; *P*=0.001)—and 1-year RFS rates were 45.9% and 28.6%, respectively (Figure 2A and Table 1). Median OS was 24.7 months with Oral-AZA and 15.4 months with placebo (HR 0.74 [95%CI 0.58–0.94]; *P*=0.0147) and estimated 1-year survival rates were 73.2% and 59.2%, respectively (Figure 2B and Table 1). Estimated median RFS was approximately twice as long with Oral-AZA compared with placebo in both the 1 Consolidation and ≥2 Consolidation cohorts, and Oral-AZA increased 1-year survival rates in these cohorts by 17.3% and 19.6%, respectively (Table 1). Oral-AZA nominally improved OS regardless of the number of prior consolidation cycles received (0, 1, or ≥2), with median OS estimates ranging from 21.0 to 28.6 months in the Oral-AZA arm and 10.9 to 17.6 months in the placebo arm (Table 1). Intriguingly, median RFS appeared favorable for patients receiving Oral-AZA without any prior consolidation therapy (8.4 months), compared with patients receiving consolidation therapy but no maintenance in the placebo arm (5.0 months). Analogously, median OS was also longer for patients receiving Oral-AZA and no prior consolidation therapy (23.3 months), compared with patients receiving consolidation therapy but no maintenance treatment in the placebo arm (15.4 months).

Overall, 79% of patients (*n*=375) received a single induction course before achieving remission and 21% (*n*=97) received ≥2 inductions (Figure 1). When accounting for total chemotherapy received before study entry (ie, number of induction and consolidation courses), median RFS was numerically prolonged by 1.5 to 8.5 months with Oral-AZA vs placebo across all induction/consolidation cohorts (Supplementary Table S2). Patients who received a single induction followed by ≥2 cycles of consolidation appeared to have

the most favorable survival outcomes within each treatment arm, whereas the small subgroup of patients who received ≥ 2 courses of induction and no consolidation generally had poor outcomes, but sample sizes prevent meaningful interpretation.

The overall safety profile of Oral-AZA was similar among consolidation groups and was aligned with the overall QUAZAR population. No associations were found between the number of consolidation cycles received and Oral-AZA dose modifications (data not shown).

As mentioned, the primary objective of the QUAZAR AML-001 trial was to determine the efficacy of Oral-AZA as maintenance therapy subsequent to chemotherapy for patients already in remission. A broad assessment of the impact of consolidation therapy in the front-line management of AML is beyond the scope of this trial, and Oral-AZA is not meant to replace consolidation chemotherapy for patients who can receive it. Overall, Oral-AZA maintenance significantly prolonged both RFS and OS compared with placebo regardless of whether patients received consolidation after initial induction. With the caveat regarding small sample sizes and lack of statistical power, post hoc analyses in subgroups defined by number of consolidation cycles received suggest that Oral-AZA may prolong RFS and OS compared with a “watch-and-wait” approach (emulated with placebo) for patients with AML in first remission after IC, independent of the number of induction and consolidation courses received before beginning maintenance treatment. A previous analysis examining the relationship between survival outcomes and MRD in QUAZAR AML-001 found that although patients with MRD responses (ie, conversion from MRD-positive at baseline to MRD-negative) were more likely to have received consolidation chemotherapy before study entry than those who remained MRD-positive on-study, the number of chemotherapy cycles received before study entry was not significantly predictive of MRD response or duration on-study with MRD-negative status.¹⁰ Overall, these findings indicate that intensive induction chemotherapy followed by Oral-AZA maintenance therapy is effective regardless of the amount of prior consolidation delivered, and represents an important component of therapy in patients with intermediate- or poor-risk AML in remission not candidates for HSCT.

References:

1. Celgene Corporation. ONUREG® (azacitidine tablets) prescribing information. https://www.accessdata.fda.gov/drugsatfda_docs/label/2020/214120s000lbl.pdf Revised September 2020. Accessed September 22, 2022.
2. European Medicines Agency, Committee for Medicinal Products for Human Use. ONUREG® (azacitidine). https://www.ema.europa.eu/en/documents/smop-initial/chmp-summary-positive-opinion-onureg_en.pdf Accessed September 22, 2021.
3. Wei AH, Döhner H, Pocock C, et al. Oral azacitidine maintenance therapy for acute myeloid leukemia in first remission. *N Engl J Med*. 2020;383(26):2526-2537.
4. Palva IP, Almqvist A, Elonen E, et al. Value of maintenance therapy with chemotherapy or interferon during remission of acute myeloid leukaemia. *Eur J Haematol*. 1991;47(3):229-233.
5. Huls G, Chitu DA, Havelange V, et al. Azacitidine maintenance after intensive chemotherapy improves DFS in older AML patients. *Blood*. 2019;133(13):1457-1464.
6. Blum W, Sanford BL, Klisovic R, et al. Maintenance therapy with decitabine in younger adults with acute myeloid leukemia in first remission: a phase 2 Cancer and Leukemia Group B Study (CALGB 10503). *Leukemia*. 2017;31(1):34-39.
7. Lowenberg B, Suciu S, Archimbaud E, et al. Mitoxantrone versus daunorubicin in induction-consolidation chemotherapy--the value of low-dose cytarabine for maintenance of remission, and an assessment of prognostic factors in acute myeloid leukemia in the elderly: final report. European Organization for the Research and Treatment of Cancer and the Dutch-Belgian Hemato-Oncology Cooperative Hovon Group. *J Clin Oncol*. 1998;16(3):872-881.
8. Lowenberg B, Beck J, Graux C, et al. Gemtuzumab ozogamicin as postremission treatment in AML at 60 years of age or more: results of a multicenter phase 3 study. *Blood*. 2010;115(13):2586-2591.
9. National Comprehensive Cancer Network Clinical Practice Guidelines in Oncology for Acute Myeloid Leukemia. National Comprehensive Cancer Network website. http://www.nccn.org/professionals/physician_gls/PDF/aml.pdf Accessed 01 Mar 2011.
10. Roboz GJ, Ravandi F, Wei AH, et al. Oral azacitidine prolongs survival of patients with AML in remission independently of measurable residual disease status. *Blood*. 2022;139(14):2145-2155.

Table 1. Estimated relapse-free and overall survival with Oral-AZA vs placebo by number of consolidation cycles received before study entry

	Oral-AZA N = 238	Placebo N = 234	Difference, Oral-AZA vs Placebo [95% CI], months
No consolidation, n (%)	52 (22)	42 (18)	
RFS, months, median [95% CI]	8.4 [7.5–16.2]	3.9 [1.9–4.9]	+4.5 months [0.8–8.2]
Oral-AZA vs Placebo: HR [95% CI]	0.55 [0.34–0.88]		
1 year RFS rate	40.8%	22.0%	+18.7% [–0.6 to +38.1]
OS, months, median [95% CI]	23.3 [13.5–37.5]	10.9 [6.3–15.7]	+12.4 months [4.7–26.7]
Oral-AZA vs Placebo: HR [95% CI]	0.55 [0.34–0.89]		
1-year OS rate	71.2%	40.5%	+30.7% [11.4–50.0]
Any consolidation* n (%)	186 (78)	192 (82)	
RFS, months, median [95% CI]	10.2 [7.7–13.1]	5.0 [4.6–7.3]	+5.2 months [2.7–7.6]
Oral-AZA vs Placebo: HR [95% CI]	0.69 [0.54–0.87]		
1 year RFS rate	45.9%	28.6%	+17.3% [7.2–27.4]
OS, months, median [95% CI]	24.7 [17.9–31.0]	15.4 [12.9–21.0]	+9.3 months [3.4–15.2]
Oral-AZA vs Placebo: HR [95% CI]	0.76 [0.60–0.97]		
1-year OS rate	73.2%	59.2%	+14.0% [4.5–23.6]
1 consolidation, n (%)	110 (46)	102 (44)	
RFS, months, median [95% CI]	10.0 [7.4–11.7]	4.7 [4.0–7.4]	+5.3 months [2.2–8.3]
Oral-AZA vs Placebo: HR [95% CI]	0.72 [0.53–0.99]		
1 year RFS rate	40.6%	23.3%	+17.3% [4.4–30.2]
OS, months, median [95% CI]	21.0 [16.7–30.5]	14.3 [11.7–18.0]	+6.7 months [0.1–13.3]
Oral-AZA vs Placebo: HR [95% CI]	0.75 [0.55–1.02]		
1-year OS rate	68.8%	59.2%	+9.6% [–3.4 to +22.6]
≥ 2 consolidations, n (%)	76 (32)	90 (38)	
RFS, months, median [95% CI]	13.0 [7.7–21.2]	6.1 [4.6–7.5]	+6.9 months [0.7–13.1]
Oral-AZA vs Placebo: HR [95% CI]	0.59 [0.41–0.87]		
1 year RFS rate	54.1%	34.5%	+19.6% [3.7–35.4]
OS, months, median [95% CI]	28.6 [17.8–41.3]	17.6 [11.6–28.7]	+11.0 months [–0.1 to +22.1]
Oral-AZA vs Placebo: HR [95% CI]	0.75 [0.50–1.11]		
1-year OS rate	80.0%	59.2%	+20.9% [7.0–34.8]
*Includes patients in the 1 Consolidation and ≥ 2 Consolidations cohorts.			
CI, confidence interval; HR, hazard ratio; Oral-AZA, oral azacitidine; OS, overall survival; RFS, relapse-free survival.			

Figure legends:

Figure 1. Patient enrollment and prior chemotherapy details.

*The ≥ 2 Consolidations cohort included 19 patients (Oral-AZA 6, placebo 13) who received 3 consolidation cycles. CR, complete remission; CRi, CR with incomplete blood count recovery; Oral-AZA, oral azacitidine.

Figure 2. Survival outcomes by prior consolidation chemotherapy use.

Kaplan-Meier estimated relapse-free survival (A) and overall survival (B) with Oral-AZA vs placebo by prior use of consolidation chemotherapy before study entry.

RFS and OS estimates were derived using Kaplan-Meier methods and compared for Oral-AZA vs placebo using a log-rank test. HRs and 95% CIs were generated using a stratified Cox proportional hazards model. CI, confidence interval; HR, hazard ratio; Oral-AZA, oral azacitidine; OS, overall survival; pts, patients; RFS, relapse-free survival.

Randomized
N = 472

1 Induction
n = 375 (79%)

> 1 Induction
n = 97 (21%)

**CR/
CRi**

No Consolidation
n = 94 (20%)

Any Consolidation
n = 378 (80%)

Randomization

Oral-AZA: n = 52 (22%)

Placebo: n = 42 (18%)

Oral-AZA: n = 186 (78%)

Placebo: n = 192 (82%)

1 Consolidation
n = 212 (45%)

≥2 Consolidations*
n = 166 (35%)

Oral-AZA: n = 110 (46%)

Placebo: n = 102 (44%)

Oral-AZA: n = 76 (32%)


Placebo: n = 90 (38%)

— Oral-AZA

— Placebo

A


No Consolidation



No. of Pts

Oral-AZA	52	17	12	4	2	0
Placebo	42	7	4	1	0	

Any Consolidation




No. of Pts

Oral-AZA	186	75	35	4	1	1
Placebo	192	48	25	5	3	0

B


No Consolidation



No. of Pts

Oral-AZA	52	37	25	16	8	4	0
Placebo	42	17	10	5	3	2	1

Any Consolidation



No. of Pts

Oral-AZA	186	131	90	43	18	11	1
Placebo	192	110	72	29	16	9	0

Supplementary data for the manuscript titled: Survival outcomes with oral azacitidine maintenance in patients with acute myeloid leukemia in remission by receipt of initial chemotherapy: subgroup analyses from the phase 3 QUAZAR AML-001 trial

Supplementary Table S1. Demographic and disease characteristics by randomized treatment arm and number of consolidation cycles received

	Oral-AZA (N = 238)				Placebo (N = 234)			
	No Consolidation n = 52	Any Consolidation n = 186*	1 Consolidation n = 110	≥ 2 Consolidations n = 76	No Consolidation n = 42	Any Consolidation n = 192*	1 Consolidation n = 102	≥ 2 Consolidations n = 90
Age, median (range), years	71 (59–84)	67 (55–86)	68 (55–86)	66 (55–75)	70 (58–81)	68 (55–82)	68 (55–78)	68 (55–82)
WHO AML classification, n (%)								
Recurrent genetic abnormalities [†]	10 (19)	29 (16)	15 (14)	14 (18)	7 (17)	39 (20)	18 (18)	21 (23)
Myelodysplasia-related changes	16 (31)	33 (18)	24 (22)	9 (12)	10 (24)	32 (17)	17 (17)	15 (17)
Not otherwise specified	26 (50)	122 (66)	71 (65)	51 (67)	25 (60)	120 (63)	66 (65)	54 (60)
De novo AML, n (%)	46 (88)	167 (90)	95 (86)	72 (95)	40 (95)	176 (92)	93 (91)	83 (92)
ECOG PS score, n (%)								
0	21 (40)	95 (51)	58 (53)	37 (49)	20 (48)	91 (47)	56 (55)	35 (39)
1	24 (46)	77 (41)	44 (40)	33 (43)	19 (45)	87 (45)	36 (35)	51 (57)
2–3	7 (13)	14 (8)	8 (7)	6 (8)	3 (7)	14 (7)	10 (10)	4 (4)
Cytogenetic risk at diagnosis, n (%)								
Intermediate	44 (85)	159 (85)	93 (85)	66 (87)	36 (86)	167 (87)	85 (83)	82 (91)
Poor	8 (15)	27 (15)	17 (15)	10 (13)	6 (14)	25 (13)	17 (17)	8 (9)
CR/CRI status at randomization, n (%)								
CR	38 (73)	145 (78)	84 (76)	61 (80)	36 (86)	141 (73)	70 (69)	71 (79)
CRI	14 (27)	37 (20)	25 (23)	12 (16)	4 (10)	40 (21)	23 (23)	17 (19)
Not in CR/CRI or Missing [‡]	1 (2)	4 (2)	1 (1)	3 (4)	2 (5)	11 (6)	9 (9)	2 (2)
Days from CR/CRI to randomization,[§] median (range)	35.0 (7–128)	88.0 (8–154)	80.5 (8–154)	92.5 (64–130)	35.5 (7–125)	88.5 (37–263)	82.0 (37–263)	92.0 (68–134)
MRD status at screening, n (%)								
Negative	30 (58)	103 (55)	58 (53)	45 (59)	15 (36)	96 (50)	49 (48)	47 (52)
Positive	21 (40)	82 (44)	51 (46)	31 (41)	26 (62)	90 (47)	52 (51)	38 (42)
Missing	1 (2)	1 (1)	1 (1)	0	1 (2)	6 (3)	1 (1)	5 (6)
Reason(s) ineligible for HSCT, n (%)								
Age	41 (79)	113 (61)	77 (70)	36 (47)	33 (79)	119 (62)	66 (65)	53 (59)
Comorbidities	8 (15)	44 (24)	33 (30)	11 (14)	10 (24)	40 (21)	22 (22)	18 (20)
No available donor	9 (17)	28 (15)	8 (7)	20 (26)	7 (17)	28 (15)	14 (14)	14 (16)
Patient decision	1 (2)	18 (10)	9 (8)	9 (12)	5 (12)	27 (14)	16 (16)	11 (12)
Performance status	7 (13)	7 (4)	5 (5)	2 (3)	3 (7)	6 (3)	3 (3)	3 (3)
Unfavorable cytogenetics	0	6 (3)	1 (1)	5 (7)	1 (2)	9 (5)	4 (4)	5 (6)
Other	5 (10)	23 (12)	11 (10)	12 (16)	2 (5)	19 (10)	7 (7)	12 (13)
[*] Includes patients in the 1 Consolidation and ≥ 2 Consolidations cohorts. [†] Central assessment by flow cytometry, using a ≥ 0.1% MRD-positive threshold ("different-from-normal" method). [‡] All patients must have been in CR/CRI at study screening; CR/CRI status was missing at randomization for two patients in the placebo arm. [§] Four patients were enrolled beyond the 4-month (± 7 days) inclusion window (protocol violations). Individual patients may be accounted for across multiple categories. AML, acute myeloid leukemia; CR, complete remission; CRI, CR with incomplete blood count recovery; ECOG PS, Eastern Cooperative Oncology Group performance status; HSCT, hematopoietic stem cell transplant; MRD, measurable residual disease; Oral-AZA, oral azacitidine; WHO, World Health Organization.								

Supplementary Table S2. Median relapse-free and overall survival with Oral-AZA vs placebo by total number of induction and consolidation cycles received before study entry

	Oral-AZA N = 238	Placebo N = 234	Oral-AZA vs Placebo	
			Difference [95% CI], months	HR [95% CI]
1 induction, no consolidation, n (%)	38 (16)	35 (15)		
RFS, median [95% CI], months	10.4 [7.7–25.1]	3.9 [1.9–4.9]	+6.5 [0.6–12.5]	0.47 [0.27–0.82]
OS, median [95% CI], months	29.3 [13.4–45.3]	10.8 [6.2–15.7]	+18.5 [3.5–33.5]	0.48 [0.28–0.82]
1 induction + 1 consolidation, n (%)	84 (35)	81 (35)		
RFS, median [95% CI], months	9.8 [7.0–11.1]	5.0 [4.0–7.6]	+4.9 [0.5–7.6]	0.82 [0.58–1.17]
OS, median [95% CI], months	19.4 [14.3–24.8]	15.0 [12.2–24.3]	+4.3 [–2.6 to +11.3]	0.91 [0.64–1.29]
1 induction + ≥ 2 consolidations, n (%)	59 (25)	78 (33)		
RFS, median [95% CI], months	13.0 [7.7–21.1]	6.1 [4.6–7.7]	+6.9 [1.0–12.7]	0.56 [0.37–0.85]
OS, median [95% CI], months	28.6 [17.7–36.6]	16.6 [11.6–27.0]	+12.0 [0.9–23.1]	0.76 [0.49–1.17]
≥ 2 inductions, no consolidation, n (%)	14 (6)	7 (3)		
RFS, median [95% CI], months	4.2 [1.9–8.4]	2.7 [0.4–9.2]	+1.5 [–3.5 to +5.6]	0.66 [0.24–1.81]
OS, median [95% CI], months	16.2 [8.9–37.2]	11.6 [3.1–NE]	+4.7 [NE–NE]	0.90 [0.31–2.61]
≥ 2 inductions + ≥ 1 consolidation, n (%)	43 (18)	33 (14)		
RFS, median [95% CI], months	12.9 [6.1–46.1]	4.4 [2.0–7.5]	+8.5 [0.4–16.1]	0.58 [0.33–1.01]
OS, median [95% CI], months	36.0 [17.9–47.2]	14.2 [8.5–22.3]	+21.8 [6.0–37.6]	0.49 [0.28–0.86]
CI, confidence interval; HR, hazard ratio; NE, not estimable; Oral-AZA, oral azacitidine; OS, overall survival; RFS, relapse-free survival.				