



SHORT REPORT

A randomised comparison of FLAG-Ida versus daunorubicin combined with clofarabine in relapsed or refractory acute myeloid leukaemia: Results from the UK NCRI AML17 trial

Nigel H. Russell¹  | Robert K. Hills²  | Lars Kjeldsen³ | Richard E. Clark⁴ | Sahra Ali⁵ | Paul Cahalin⁶ | Ian F. Thomas⁷ | Alan K. Burnett⁸

¹Department of Haematology, Guy's Hospital, London, UK

²Nuffield Department of Population Health, University of Oxford, Oxford, UK

³Department of Haematology, Rigshospitalet, Copenhagen, Denmark

⁴Department of Molecular and Clinical Cancer Medicine, University of Liverpool, UK

⁵Department of Haematology, Castle Hill Hospital, Hull, UK

⁶Department of Haematology, Blackpool Victoria Hospital, Blackpool, UK

⁷Centre for Trials Research, Cardiff University, Cardiff, UK

⁸Paul O'Gorman Leukaemia Centre, University of Glasgow, Glasgow, UK

Correspondence

Nigel H. Russell, Department of Haematology, Guy's Hospital, London, UK.
Email: nigel.russell@nottingham.ac.uk

Funding information

Cancer Research UK

Summary

The prognosis for younger patients with relapsed acute myeloid leukaemia (AML) is generally dismal. Allogeneic stem cell transplantation is the preferred therapy for these patients. As part of the UK NCRI AML17 trial, daunorubicin/clofarabine (DClo) was compared with fludarabine, cytarabine, granulocyte colony-stimulating factor with idarubicin (FLAG-Ida) in 311 patients designated high-risk following course one of induction therapy, which has previously been reported. We now report the results of the same randomisation in patients who were refractory to two induction courses or subsequently relapsed. A total of 94 relapsed or refractory AML patients, usually less than 60 years of age and with mainly favourable or intermediate-risk cytogenetics, were randomised to receive up to three courses of DClo or FLAG-Ida, with the aim of proceeding to transplant. Complete remission was achieved in 74% of patients with no difference between the arms. Overall, 57% of patients received a transplant with no difference between the arms, likewise overall survival at five years showed no significant difference (21% for DClo vs. 22% for FLAG-Ida). No patient who did not receive a transplant survived beyond 21 months. A stratified analysis including the 311 post course 1 high-risk patients who underwent the same randomisation showed a consistent treatment benefit for FLAG-Ida.

KEYWORDS

chronic lymphocytic leukaemia, morphology, CLL lymphocytes

INTRODUCTION

Patients with relapsed and refractory acute myeloid leukaemia (AML) represent an important unmet therapeutic need, for which there is no universally accepted standard of care. European Leukemia Network (ELN) guidelines suggest a number of regimens including intermediate-dose cytarabine with or without an anthracycline, MEC [mitoxantrone, etoposide and cytosine arabinoside (Ara-C)] or

fludarabine, cytarabine and granulocyte colony-stimulating factor (GCSF) with idarubicin (FLAG-Ida).¹ Furthermore, several new chemotherapy drugs having failed to improve survival in randomised studies against standard regimens^{2,3} although the *FLT3* inhibitor gilteritinib was found to be superior to chemotherapy in relapsed *FLT3*-mutated AML.⁴ Recently FLAG-Ida combined with venetoclax has been reported as having promising results in both newly diagnosed and relapsed/refractory AML.⁵

The challenge with relapsed/refractory AML is twofold. Firstly to improve the remission rate and secondly to deliver more patients to transplant which is the only curative option for the majority of patients. For relapsed patients the FLAG-Ida (fludarabine/ara-C/GCSF and idarubicin) has been widely used¹ and in our MRC AML15 trial FLAG-Ida given for the first two treatment courses had a significantly superior anti-leukaemia effect when compared to '7 + 3'-like chemotherapy⁶. In the AML17 trial we chose as the comparative treatment to replace Ara-C in a daunorubicin/Ara-C combination, with the alternative nucleoside, clofarabine.

Clofarabine (2-chloro-2'-fluoro-deoxy-9-β-D-arabinofuranosyladenine) is a novel nucleoside analogue developed as the result of a series of chemical modifications to minimise cleavage while retaining activity.⁷ Although not approved as upfront therapy in AML despite activity in adverse-risk disease^{8,9} promising results have been reported when given in combination with Ara-C and GCSF (GCLAC regimen) in the relapsed setting.¹⁰ Following a feasibility study combining daunorubicin with clofarabine (DClo), we prospectively compared the combination with FLAG-Ida in both high-risk AML following course 1 of induction and in relapsed/refractory patients who had previously entered the AML17 trial. The high-risk experience has previously been reported.¹¹ Here we report on the outcome of the relapsed and refractory patients.

PATIENTS AND METHODS

As part of the UK National Cancer Research Institute (NCRI) AML17 trial (ISRCTN55675535) designed primarily for patients under 60 years of age with untreated *de novo* or secondary AML and high-risk myelodysplastic syndrome (defined as >10% marrow blasts at diagnosis), patients who were high-risk post course 1 ($n = 311$) could enter a randomisation to compare FLAG-Ida with DClo. In addition, this randomisation was open to patients in first morphological relapse and to those who were refractory to two courses of induction chemotherapy. Patients who entered the high-risk randomisation in CR1 were not eligible to re-enter the randomisation if they subsequently relapsed. Patients older than 60 years could enter the trial if considered fit for intensive therapy.

The trial was conducted in accordance with the Declaration of Helsinki, was sponsored by Cardiff University, and was approved by the Wales Research Ethic Committee 3. All patients provided written informed consent to random assignment. Patients needed to be aged over 16 years at trial entry. Relapse of disease was defined as more than 5% blasts in the marrow. Patients who relapsed after treatment in NCRI trials prior to AML17 could be included. Patients were randomly assigned in a 2:1 ratio to DClo or FLAG-Ida, each for up to three courses stratified by baseline information. The intention was that every patient would be eligible to proceed to allogeneic transplant. Consenting patients

were randomised^{1,2} to receive up to three courses of DClo (daunorubicin 50 mg/m² on days 1, 3 and 5 and clofarabine 20 mg/m² on days 1–5) or FLAG-Ida (fludarabine 30 mg/m² on days 2–6, Ara-C 2 g/m² on days 2–6, GCSF 263 µg on days 1–7, idarubicin 8 mg/m² on days 4–6).

Statistical methods

The primary outcome measures for the trial were the number of patients delivered to transplant and overall survival (OS). All end-points were defined according to the revised International Working Group criteria.¹²

All analyses are by intention-to-treat. Categorical end-points [e.g. complete remission (CR) rates] were compared using Mantel–Haenszel tests, giving Peto odds ratios (OR) and confidence intervals (CI). Continuous/scale variables were analysed by non-parametric (Wilcoxon rank sum) tests. Time-to-event outcomes were analysed using the log-rank test, with Kaplan–Meier survival curves. Odds/hazard ratios (OR/HR) less than 1 indicate benefit for the investigational therapy [daunorubicin/clofarabine (DClo)] versus standard therapy (FLAG-Ida). All survival percentages are at five years unless otherwise stated except for survival censored at transplant where because of lack of follow-up to five years among surviving non-transplanted patients percentages are given at four years. Median follow-up is 46.7 months (range 5.0–68.0 months). Stratified log-rank tests with tests for interaction were performed using the methodology of the Early Breast Cancer Trialists' Collaborative Group¹³

RESULTS

Between November 2009 and December 2012, 94 patients (88 relapsed and six refractory) entered the randomisation, of whom 79 had received initial induction therapy within AML17. The induction that the patients had received within the AML17 trial overall varied over the period of the trial and included ADE (Ara-C/daunorubicin/etoposide) with or without the addition of gemtuzumab ozogamicin (GO) either at a dose of 3 mg/m² (GO3) or 6 mg/m² (GO6); DA (daunorubicin/Ara-C) with GO3 or GO6, or DA with the daunorubicin dose being 60 mg/m² (DA60) or 90 mg/m² (DA90).^{14,15} For the purposes of this analysis relapsed and refractory patients are combined. A CONSORT diagram for the whole trial is shown in Figure S1. The patients' characteristics and details of prior upfront treatments are given in Table 1; these are well balanced between the arms with 51% of patients in the DClo arm receiving prior GO compared to 59% in the FLAG-Ida arm. Sixty-two patients were allocated to DClo and 32 to FLAG-Ida. The median age of randomised patients was 47 years (range 19–63). Of note, 13 (14%) recruited were over 60 years. Only 3/94 patients had adverse-risk cytogenetics reflecting their categorisation as high-risk

TABLE 1 Patient characteristics

	DClo	FLAG-Ida
Number randomised	62	32
Age group (years)		
15–29	8	7
30–39	8	2
40–49	15	8
50–59	22	11
60+	9	4
Gender		
Female	35	16
Male	27	16
Type of disease		
<i>De novo</i>	58	31
Secondary	1	0
High-risk MDS	3	1
Performance status		
0	46	26
1	12	5
2	3	1
3	1	0
4	0	0
Induction treatment		
ADE	12	5
ADE+GO3	7	4
ADE+GO6	7	5
DA+GO3	12	4
DA+GO6	11	6
DA60	5	3
DA90	2	1
Not AML17	6	4
Cytogenetics		
Favourable	9	5
Intermediate	40	19
Adverse	2	1
NK	11	7
FLT3 ITD		
WT	30	17
Mutant	23	10
Not known	9	5
NPM1		
WT	34	18
Mutant	17	7
Not known	11	7
Status of rel/ref		
Relapsed	58	30
Refractory	4	2

Abbreviations: ADE, cytosine arabinoside, daunorubicin and etoposide; AML, acute myeloid leukaemia; DA, daunorubicin and cytosine arabinoside; DClo, daunorubicin/clofarabine; FLAG-Ida, fludarabine, cytarabine, granulocyte colony-stimulating factor with idarubicin; GO, gemtuzumab ozogamicin; ITD, internal tandem duplication; MDS, myelodysplastic syndrome; WT, wild type.

post course 1 and hence ineligibility to re-enter the randomisation at relapse. Thirty-three of the 94 patients had a *FLT3* mutation present at diagnosis but *FLT3* status was not determined at relapse. Seventy of the 94 patients (74%) achieved a CR or CR with incomplete bone marrow recovery (CR/CRi) after randomisation. The median time to remission from randomisation was 46 days among remitters, with only 63% of remitters achieving their remission within 60 days of randomisation. Twenty-four patients never achieved a CR or CRi after two courses of induction. The response by treatment allocation for randomised patients is shown in Table 2 and was 74% and 75% for DClo and FLAG-Ida respectively with no significant difference in 30- or 60-day mortality between the arms (6% and 11% for DClo vs 3% and 6% for FLAG-Ida, $p = 0.7$ and 0.8 respectively).

Transplant

Fifty-three patients (57%) received an allogeneic transplant; in addition, nine received a transplant of unknown type; two allografts occurred prior to randomisation (plus one additional transplant of unknown type). Of the remaining 51 allogeneic transplants there were 42 myeloablative transplants (DClo 28 and FLAG-Ida 14;), and nine reduced-intensity transplants (DClo six, FLAG-Ida three). Overall, excluding patients transplanted before randomisation, the rate of allogeneic transplantation did not differ between arms [57% vs 55%; OR 0.81 (0.35–1.90); $p = 0.6$].

The OS at five years from the point of randomisation did not differ for DClo versus FLAG-Ida [21% vs 22%; HR 1.24 (0.76–2.33); $p = 0.4$] (Figure 1A). A test for interaction shows no evidence of heterogeneity of effect by GO upfront ($p = 0.17$) No patient who was not transplanted survived beyond 21 months; the HR for survival censored at stem cell transplantation (SCT) was 1.11 (0.52–2.34), $p = 0.8$. When looking at transplantation in the patients who entered remission, there was no significant difference in relapse-free survival (RFS) post transplant (38% vs 29% at four years, HR 1.11 (0.49–2.52); $p = 0.8$) (Table 2).

Stratified analysis of high-risk and relapsed/refractory AML

In total 405 patients in AML17 entered the DClo versus FLAG-Ida randomisation. This included the 311 from the high-risk randomisation post course 1 whose outcome has been previously published¹¹ and the 94 relapsed/refractory patients reported here. The two groups of patients were combined using standard stratified analytical techniques. The results of the two groups were consistent with each other ($p = 1.0$ between treatment and group). Overall, the whole trial showed a consistent benefit for FLAG-Ida (HR 1.35, CI 1.06–1.73; $p = 0.02$) with no evidence that the relapsed/refractory patients are any different.

TABLE 2 Outcomes of relapsed/refractory randomisation

	DClo	FLAG-Ida	HR/OR, 95% CI	p-value
ORR (CR + CRi)	74%	75%	1.04 (0.39–2.76)	0.9
30-days mortality	6%	3%	1.92 (0.30–12.2)	0.5
60-days mortality	11%	6%	1.76 (0.45–6.93)	0.4
5-year OS	21%	22%	1.24 (0.76–2.33)	0.4
4-year OS censored at SCT	0%	0%	1.11 (0.52–2.34)	0.8

Abbreviations: CI, confidence interval; CR/CRi, complete remission /with incomplete bone marrow recovery; DClo, daunorubicin/clofarabine; FLAG-Ida, fludarabine, cytarabine, granulocyte colony-stimulating factor with idarubicin; HR, hazard ratio; OR, odds ratio; ORR, overall response rate; OS, overall survival; SCT, stem cell transplantation.




FIGURE 1 (A) Overall survival by randomisation; (B) stratified survival analysis for all high-risk patients [Colour figure can be viewed at wileyonlinelibrary.com]

DISCUSSION

In this trial the aim was to compare a novel salvage regimen against FLAG-Ida and to improve upon the number of patients proceeding to transplant and survival post transplant. In the relapse/refractory patients reported here there was a high overall response rate (CR + CRi) of 74% with no difference between DClo and FLAG-Ida. This

unexpectedly high response rate reflects the facts that although the median first remission duration was 10 months and approximately two thirds of patients relapsed after a remission duration of less than one year, nonetheless, 19% of randomised patients had favourable cytogenetics, of whom 93% achieved CR2. This compares with only 1% of patients entering the randomisation having adverse-risk cytogenetics; and only four patients had received a prior

stem cell transplantation. This disposition of patients reflects the trial structure that excluded patients from the relapse randomisation if they had previously been defined as high risk post course 1. Although the total number of patients was relatively small, there was no difference in the proportion of patients in each arm receiving an allogeneic transplant (57% vs 55%). There was no difference in the risk of relapse after transplant and there was no difference in OS between the arms. No patient who was not transplanted survived beyond 21 months, which emphasises the importance of transplant in relapsed disease. The disposition of patients entering the randomisation makes comparison with the FLAG-Ida-venetoclax combination problematic as that study included more patients with adverse cytogenetics and prior transplant as well as patients receiving second salvage treatment, although of note, patients with favourable-risk cytogenetics in that study performed less well with FLAG-Ida-venetoclax.⁴

In conclusion, although the FLAG-Ida schedule was not superior in the relapsed setting we previously found it to be so in high-risk patients upfront and a stratified analysis of all patients entering the high-risk randomisation showed overall benefit making it the standard of care for relapsed disease, certainly for patients with favourable or intermediate-risk cytogenetics without a *FLT3* mutation. Whether the combination of FLAG-Ida with venetoclax improves outcome further is a question for future studies.

ACKNOWLEDGEMENTS

The research costs of the trial were provided by Cancer Research UK. Clofarabine was provided by Genzyme Inc. (Boston, MA, USA). The following institutions and investigators recruited patients to this study:

Aalborg University Hospital: Dr Maria Kallenbach; Aarhus University Hospital: Dr Jan Maxwell Norgaard, Dr Mette Holm; Addenbrooke's Hospital: Dr Charles Crawley, Dr Jenny Craig; Auckland City Hospital: Dr Richard Doocey, Dr Timothy Hawkins; Ayr Hospital: Dr Paul Eynaud; Barnet General Hospital: Dr Andres Virchis; Barts and the London NHS Trust: Dr Jamie Cavenagh, Dr Matthew Smith; Basingstoke and North Hampshire Foundation NHS Trust: Dr Sylwia Simpson; Beatson West of Scotland Cancer Centre: Dr Andrew Clark, Dr Anne Parker, Dr Mark Drummond, Dr Pam McKay; Belfast City Hospital: Dr Mary Francis McMullin; Birmingham Heartlands Hospital: Dr Donald Milligan, Dr Guy Pratt, Dr Matthew Lumley, Dr Neil Smith, Dr Richard Lovell, Dr Shankara Paneesha; Blackpool Victoria Hospital NHS Foundation Trust: Dr Paul Cahalin; Borders General Hospital: Dr Ashok Okhandiar; Bradford Royal Infirmary: Dr Lisa Newton, Dr Nitin Sood, Dr Sam Ackroyd; Bristol Haematology and Oncology Centre: Dr David Marks, Dr Roger Evelyn; Chesterfield Royal Hospital: Dr Emma Welch, Dr Robert Cutting; Christchurch Hospital: Dr Liam Fernyhough, Dr Mark Smith, Dr Ruth Spearing, Dr Steve Gibbons; Christie Hospital NHS Trust: Dr Mike Dennis; Countess of Chester Hospital: Dr Salah Tuegar; Darent Valley Hospital: Dr Anil Kamat, Dr Tariq

Shafi; Derby Hospitals NHS Foundation Trust: Dr Adrian Smith, Dr Juanah Addada; Derriford Hospital: Dr Adrian Copplestone, Dr Hannah Hunter, Dr Mike Hamon, Dr Tim Nokes, Dr Wayne Thomas; Doncaster Royal Infirmary: Dr Joe Joseph; East Kent Hospitals University NHS Foundation Trust: Dr Christopher Pocock, Dr Jindriska Lindsay, Dr Kamiran Saied, Dr Vijay Ratnayake; East Sussex Hospitals NHS Trust: Dr Judy Beard, Dr Satyajit Sahu; Falkirk and District Royal Infirmary: Dr Christopher Brammer, Dr Marie Hughes; Glan Clwyd Hospital: Dr Christine Hoyle; Gloucestershire Royal Hospital: Dr Adam Rye, Dr Eve Blundell, Dr Rebecca Frewin, Dr Sally Chown; Great Western Hospital: Dr Alex Sternberg; Guys and St Thomas' Foundation Trust: Dr Matthew Smith, Dr Robert Carr; Hammersmith Hospital: Dr Jiri Pavlu; Herlev University Hospital: Dr Morten Krogh Jensen; Hillingdon Hospital: Dr Riaz Janmohamed, Dr Richard Kaczmarek; Hull Royal Infirmary: Dr Sahra Ali; Ipswich Hospital NHS Trust: Dr Isobel Chalmers; James Cook University Hospital: Dr Chris Millar, Dr Ray Dang; James Paget University Hospital: Dr Cesar Gomez, Dr Shala Sadullah; John Radcliffe Hospital: Dr Pares Vyas; Kettering General Hospital: Dr Isaac Wilson-Morke, Dr Karyn Longmuir, Dr Mark Kwan; Leicester Royal Infirmary: Dr Ann Hunter, Dr Murray Martin; Lincoln County Hospital: Dr Kandeepan Saravanamuttu; Maidstone Hospital: Dr Don Gillett, Dr Richard Gale; Manchester Royal Infirmary: Dr Eleni Tholouli, Dr Guy Lucas; Medway Maritime Hospital: Dr Ayed Eden, Dr Vivienne Andrews; Milton Keynes Hospital NHS Foundation Trust: Dr Moez Dungarwalla; Monklands Hospital: Dr John Murphy, Dr Pamela Paterson; New Cross Hospital: Dr Abraham Jacob, Dr Sunil Hada; NHS Grampian: Dr Dominic Culligan, Dr Jane Tighe; Ninewells Hospital: Dr David Meiklejohn, Dr Ron Kerr; Norfolk and Norwich University Hospital NHS Foundation Trust: Dr Matthew Lawes; Northampton General Hospital: Dr Angela Bowen, Dr Suchitra Krishnamurthy@nhs.uk; Northwick Park Hospital: Dr Charalampia Kyriakov, Dr Nicki Panoskaltis, Dr Robert Ayto; Nottingham University Hospitals NHS Trust: Dr Emma Dasgupta, Dr Jenny Byrne, Dr Nigel Russell; Odense University Hospital: Dr Lone Friis; Palmerston North Hospital: Dr Bart Baker; Peterborough District Hospital: Dr Sateesh Nagumantry; Pinderfields General Hospital: Dr Paul Moreton; Poole General Hospital: Dr Fergus Jack, Dr Rebecca Maddams; Princess Royal University Hospital: Dr Bipin Vadhir, Dr Corinne DeLord; Queen Alexandra Hospital: Dr Helen Dignum, Dr Robert Corser; Queen Elizabeth Hospital Birmingham: Dr Jim Murray; Queen Elizabeth Hospital, Norfolk: Dr Jane Keidan; Queens Hospital: Dr Claire Hemmaway, Dr Jane Stevens; Raigmore Hospital: Dr Chris Lush, Dr Joanne Craig; Rigshospitalet: Dr Carsten Niemann, Dr Lars Kjeldsen, Dr Ove Juul Nielsen, Dr Peter Kampmann, Dr Soeren Lykke Petersen; Rotherham General Hospital: Dr Arun Alfred, Dr Richard Went; Royal Berkshire Hospital: Dr Rebecca Sampson; Royal Bournemouth Hospital: Dr Joseph Chacko, Dr Rachel Hall; Royal Cornwall Hospital:

Dr Bryson Pottinger; Royal Devon and Exeter Hospital: Dr Anthony Todd, Dr Claudius Rudin, Dr Loretta Ngu, Dr Malcolm Hamilton, Dr Paul Kerr; Royal Free Hospital: Dr Christopher McNamara, Dr Panos Kottaridis; Royal Marsden Hospital: Dr Mark Ethell, Dr Mike Potter; Royal Oldham: Dr Allameddine Allameddine; Royal Oldham: Dr David Osborne; Royal Oldham: Dr Vivek Sen; Royal Surrey County Hospital: Dr Elisabeth Grey-Davies, Dr Johannes DeVos, Dr Louise Hendry; Royal United Hospital Bath: Dr Josephine Crowe; Russells Hall Hospital: Dr Savio Fernandes, Dr Stephen Jenkins; Salford Royal Hospital: Dr John Houghton, Dr Mark Williams; Sandwell Hospital: Dr Farooq Wandoo, Dr Richard Murrin, Dr Yasmin Hasan; Singleton Hospital: Dr Hamdi Sati, Dr Saad Ismail; South Devon Healthcare NHS Foundation Trust: Dr Deborah Turner, Dr Patrick Roberts; Southampton General Hospital: Dr Matthew Jenner; Southern General Hospital: Dr Anne Morrison; St James's University Hospital: Dr David Bowen; St Richard's Hospital: Dr Phillip Bevan; Stoke Mandeville Hospital: Dr Helen Eagleton, Dr Jonathan Pattinson; Sunderland Royal Hospital: Dr Mike Galloway, Dr Scott Marshall; Taunton and Somerset Foundation Trust: Dr Belinda Austen; The Newcastle upon Tyne NHS Foundation Trust: Dr Anne Lennard, Dr Gail Jones; The Royal Liverpool University Hospital: Dr Rahuman Salim, Dr Richard Clark; University College London Hospitals: Dr Anthony Goldstone, Dr Asim Khwaja, Dr David Linch; University Hospital Aintree: Dr Barbara Hammer, Dr Barrie Woodcock, Dr Walid Sadik; University Hospital Coventry (Walsgrave): Dr Nicholas Jackson, Dr Shailesh Jobanputra, Dr Syed Bokhari; University Hospital Lewisham: Dr Naheed Mir; University Hospital of North Staffordshire NHS Trust: Dr Deepak Chandra, Dr Richard Chasty; University Hospital of North Tees and Hartlepool: Dr Philip Mounter, Dr Zor Maung; University Hospital of Wales: Dr Jonathan Kell, Dr Steve Knapper; Victoria Hospital NHS, Fife: Dr Peter Williamson, Dr Stephen Rogers; Waikato Hospital: Dr Gillian Corbett; Western General Hospital: Dr Peter Johnson, Dr P. H. Roddie; Wishaw General Hospital: Dr Annielle Hung; Worcestershire Royal Hospital: Dr Juliet Mills; Worthing Hospital: Dr Santosh Narat; York Hospital: Dr Laura Munro, Dr Lee Bond.

CONFLICT OF INTEREST

The authors declare no competing interests.

AUTHOR CONTRIBUTIONS

Conception and design: Alan K. Burnett, Nigel H. Russell and Robert K. Hills. Provision of study materials or patients: Alan K. Burnett, Nigel H. Russell, Richard E. Clark, Sahra Ali, Lars Kjeldsen, Paul Cahalin. Collection and assembly of data: Alan K. Burnett, Robert K. Hills, Nigel H. Russell, Richard E. Clark, Paul Cahalin, Sahra Ali, Lars Kjeldsen, Ian F. Thomas. Data analysis and interpretation: Alan K. Burnett, Robert K. Hills, Nigel H. Russell. Manuscript writing: Nigel H. Russell and Alan K. Burnett drafted the paper which was revised and approved by all authors.

CLINICAL TRIAL INFORMATION

The study is part of the National Cancer Research Institute (NCRI) Acute Myeloid Leukaemia and High-Risk Myelodysplastic Syndrome Trial 17 (Trial Reference ISRCTN 55675535).

ORCID

Nigel H. Russell  <https://orcid.org/0000-0003-1893-8155>

Robert K. Hills  <https://orcid.org/0000-0003-0166-0062>

REFERENCES

1. Döhner H, Estey E, Grimwade D, Amadori S, Appelbaum FR, Büchner T, et al. Diagnosis and management of AML in adults: 2017 ELN recommendations from an international expert panel. *Blood*. 2017;129(4):424–47.
2. Roboz GJ, Rosenblat T, Arellano M, Gobbi M, Altman JK, Montesinos P, et al. International randomized phase III study of elacytarabine versus investigator choice in patients with relapsed/refractory acute myeloid leukemia. *J Clin Oncol*. 2014;32(18):1919–26.
3. Ravandi F, Ritchie EK, Sayar H, Lancet JE, Craig MD, Vey N, et al. Vosaroxin plus cytarabine versus placebo plus cytarabine in patients with first relapsed or refractory acute myeloid leukaemia (VALOR): a randomised, controlled, double-blind, multinational, phase 3 study. *Lancet Oncol*. 2015;16(9):1025–36.
4. Perl AE, Martinelli G, Cortes JE, Neubauer A, Berman E, Paolini S, et al. Gilteritinib or chemotherapy for relapsed or refractory *FLT3*-mutated AML. *N Engl J Med*. 2019;381(18):1728–40.
5. DiNardo CD, Lachowicz CA, Takahashi K, et al. Venetoclax combined with FLAG-IDA induction and consolidation in newly diagnosed and relapsed or refractory acute myeloid leukemia. *J Clin Oncol*. 2021;39(25):2768–78.
6. Burnett AK, Russell NH, Hills RK, Hunter AE, Kjeldsen L, Yin J, et al. Optimization of chemotherapy for younger patients with acute myeloid leukemia: results of the medical research council AML15 trial. *J Clin Oncol*. 2013;31(27):3360–8.
7. Kantarjian H, Gandhi V, Cortes J, Verstovsek S, du M, Garcia-Manero G, et al. Phase 2 clinical and pharmacologic study of clofarabine in patients with refractory or relapsed acute leukemia. *Blood*. 2003;102(7):2379–86.
8. Kantarjian HM, Erba HP, Claxton D, Arellano M, Lyons RM, Kvasovics T, et al. Phase II study of clofarabine monotherapy in previously untreated older adults with acute myeloid leukemia and unfavorable prognostic factors. *J Clin Oncol*. 2010;28(4):549–55.
9. Burnett AK, Russell NH, Kell J, Dennis M, Milligan D, Paolini S, et al. European development of clofarabine as treatment for older patients with acute myeloid leukemia considered unsuitable for intensive chemotherapy. *J Clin Oncol*. 2010;28:2389–95.
10. Becker PS, Kantarjian HM, Appelbaum FR, Petersdorf SH, Storer B, Pierce S, et al. Clofarabine with high dose cytarabine and granulocyte colony-stimulating factor (G-CSF) priming for relapsed and refractory acute myeloid leukaemia. *Br J Haematol*. 2011;155(2):182–9.
11. Burnett AK, Hills RK, Nielsen OJ, Freeman S, Ali A, Cahalin P, et al. A comparison of FLAG-Ida and daunorubicin combined with clofarabine in high-risk acute myeloid leukaemia: data from the UKNCRI AML17 trial. *Leukemia*. 2018;32(12):2693–7.
12. Cheson BD, Bennett JM, Kopecky KJ, Büchner T, Willman CL, Estey EH, et al. Revised recommendations of the international working Group for Diagnosis, standardization of response criteria, and reporting standards for therapeutic trials in acute myeloid leukemia. *J Clin Oncol*. 2003;21:4642–9.
13. Early Breast Cancer Trialists' Collaborative Group. Treatment of early breast cancer, volume 1: worldwide evidence 1985–1990. Oxford, United Kingdom: Oxford University Press; 1990.
14. Burnett A, Cavenagh J, Russell N, Hills R, Kell J, Jones G, et al. Defining the dose of gemtuzumab ozogamicin in combination with induction chemotherapy in acute myeloid leukemia: a comparison of

3 mg/m² with 6 mg/m² in the NCRI AML17 trial. *Haematologica*. 2016;101(6):724–31.

15. Burnett AK, Russell NH, Hills RK, Kell J, Cavenagh J, Kjeldsen L, et al. A randomized comparison of daunorubicin 90 mg/m² vs 60 mg/m² in AML induction: results from the UKNCRI AML17 trial in 1206 patients. *Blood*. 2015;125(25):3878–85.

SUPPORTING INFORMATION

Additional supporting information may be found in the online version of the article at the publisher's website.

How to cite this article: Russell NH, Hills RK, Kjeldsen L, Clark RE, Ali S, Cahalin P, et al. A randomised comparison of FLAG-Ida versus daunorubicin combined with clofarabine in relapsed or refractory acute myeloid leukaemia: Results from the UK NCRI AML17 trial. *Br J Haematol*. 2022;198:528–534. doi:[10.1111/bjh.18195](https://doi.org/10.1111/bjh.18195)