

MYELOID NEOPLASIA

CEBPA mutations in 4708 patients with acute myeloid leukemia: differential impact of bZIP and TAD mutations on outcome

Franziska Taube,^{1,*} Julia Annabell Georgi,^{1,*} Michael Kramer,¹ Sebastian Stasik,¹ Jan Moritz Middeke,¹ Christoph Röllig,¹ Utz Krug,² Alwin Krämer,³ Sebastian Scholl,⁴ Andreas Hochhaus,⁴ Tim H. Brümmendorf,⁵ Ralph Naumann,⁶ Andreas Petzold,⁷ Roger Mulet-Lazaro,⁸ Peter J. M. Valk,⁸ Björn Steffen,⁹ Hermann Einsele,¹⁰ Markus Schaich,¹¹ Andreas Burchert,¹² Andreas Neubauer,¹² Kerstin Schäfer-Eckart,¹³ Christoph Schliemann,¹⁴ Stefan W. Krause,¹⁵ Mathias Hänel,¹⁶ Richard Noppeney,¹⁷ Ulrich Kaiser,¹⁸ Claudia D. Baldus,¹⁹ Martin Kaufmann,²⁰ Sylvia Herold,²¹ Friedrich Stölzel,¹ Katja Sockel,¹ Malte von Bonin,¹ Carsten Müller-Tidow,³ Uwe Platzbecker,²² Wolfgang E. Berdel,¹⁴ Hubert Serve,⁹ Gerhard Ehninger,²³ Martin Bornhäuser,^{1,24} Johannes Schetelig,^{1,25} Christian Thiede,^{1,26} for the Study Alliance Leukemia (SAL)

¹Medizinische Klinik und Poliklinik I, Universitätsklinikum Carl Gustav Carus, Dresden, Germany; ²Medizinische Klinik III, Klinikum Leverkusen, Leverkusen, Germany; ³Universität Heidelberg, Medizinische Klinik und Poliklinik, Abteilung Innere Medizin V, Heidelberg, Germany; ⁴Klinik für Innere Medizin II, Universitätsklinikum Jena, Jena, Germany; ⁵Medizinische Klinik IV, Uniklinik Rheinisch-Westfälische Technische Hochschule (RWTH) Aachen, Aachen, Germany; ⁶Medizinische Klinik III, St Marien-Krankenhaus Siegen, Siegen, Germany; ⁷DRESDEN-Concept Genome Center, Center for Molecular and Cellular Bioengineering, Technische Universität (TU) Dresden, Dresden, Germany; ⁸Department of Hematology, Erasmus Medical Center (MC) Cancer Institute, Rotterdam, The Netherlands; ⁹Medizinische Klinik 2, Hämatologie/Onkologie, Johann Wolfgang Goethe-Universität, Frankfurt am Main, Germany; ¹⁰Medizinische Klinik und Poliklinik II, Universitätsklinikum Würzburg, Würzburg, Germany; ¹¹Klinik für Hämatologie, Onkologie und Palliativmedizin, Rems-Murr-Klinikum Winnenden, Winnenden, Germany; ¹²Klinik für Innere Medizin, Schwerpunkt Hämatologie, Onkologie und Immunologie, Philipps Universität Marburg, Marburg, Germany; ¹³Klinikum Nürnberg, Paracelsus Medizinische Privatuniversität, Medizinische Klinik 5, Nürnberg, Germany; ¹⁴Medizinische Klinik A, Universitätsklinikum Münster, Münster, Germany; ¹⁵Medizinische Klinik 5, Universitätsklinikum Erlangen, Erlangen, Germany; ¹⁶Klinik für Innere Medizin III, Klinikum Chemnitz, Chemnitz, Germany; ¹⁷Klinik für Hämatologie, Universitätsklinikum Essen, Essen, Germany; ¹⁸Medizinische Klinik II, St Bernward Krankenhaus, Hildesheim, Germany; ¹⁹Klinik für Innere Medizin II, Universitätsklinikum Schleswig-Holstein, Campus Kiel, Kiel, Germany; ²⁰Abteilung für Hämatologie, Onkologie und Palliativmedizin, Robert-Bosch-Krankenhaus, Stuttgart, Germany; ²¹Institut für Pathologie, Universitätsklinikum Carl Gustav Carus, Dresden, Germany; ²²Klinik und Poliklinik für Hämatologie, Zelltherapie und Hämostaseologie, Universitätsklinikum Leipzig, Leipzig, Germany; ²³GEMoAB Monoclonals GmbH, Dresden, Germany; ²⁴National Center for Tumor Diseases NCT, Dresden, Germany; ²⁵DKMS Clinical Trials Unit, Dresden, Germany; and ²⁶Agendix GmbH, Dresden, Germany

KEY POINTS

- **CEBPA^{smbZIP}- and CEBPA^{bi}-mutant AML share clinical and mutational characteristics and are distinct from CEBPA^{smTAD}-mutant AML.**
- **Only in-frame mutations in CEBPA-bZIP are associated with favorable clinical response in monoallelic and biallelic constellations.**

Biallelic mutations of the CEBPA gene (CEBPA^{bi}) define a distinct entity associated with favorable prognosis; however, the role of monoallelic mutations (CEBPAsm) is poorly understood. We retrospectively analyzed 4708 adults with acute myeloid leukemia (AML) who had been recruited into the Study Alliance Leukemia trials, to investigate the prognostic impact of CEBPAsm. CEBPA mutations were identified in 240 patients (5.1%): 131 CEBPA^{bi} and 109 CEBPAsm (60 affecting the N-terminal transactivation domains [CEBPA^{smTAD}] and 49 the C-terminal DNA-binding or basic leucine zipper region [CEBPA^{smbZIP}]). Interestingly, patients carrying CEBPA^{bi} or CEBPA^{smbZIP} shared several clinical factors: they were significantly younger (median, 46 and 50 years, respectively) and had higher white blood cell (WBC) counts at diagnosis (median, $23.7 \times 10^9/L$ and $35.7 \times 10^9/L$) than patients with CEBPA^{smTAD} (median age, 63 years, median WBC $13.1 \times 10^9/L$; $P < .001$). Co-mutations were similar in both groups: GATA2 mutations (35.1% CEBPA^{bi}; 36.7% CEBPA^{smbZIP} vs 6.7% CEBPA^{smTAD}; $P < .001$) or NPM1 mutations (3.1% CEBPA^{bi}; 8.2% CEBPA^{smbZIP} vs 38.3% CEBPA^{smTAD}; $P < .001$). CEBPA^{bi} and CEBPA^{smbZIP}, but not CEBPA^{smTAD} were associated with significantly improved overall (OS; median 103 and 63

vs 13 months) and event-free survival (EFS; median, 20.7 and 17.1 months vs 5.7 months), in univariate and multivariable analyses. Additional analyses revealed that the clinical and molecular features as well as the favorable survival were confined to patients with in-frame mutations in bZIP (CEBPA^{bZIP-inf}). When patients were classified according to CEBPA^{bZIP-inf} and CEBPA^{other} (including CEBPA^{smTAD} and non-CEBPA^{bZIP-inf}), only patients bearing CEBPA^{bZIP-inf} showed superior complete remission rates and the longest median OS and EFS, arguing for a previously undefined prognostic role of this type of mutation.

Introduction

Patients with normal karyotype represent the largest subgroup of patients with acute myeloid leukemia (AML).¹ Research performed during the past 20 years has revealed several novel abnormalities that are particularly common in this subgroup of patients (eg, activating mutations of the FLT3 receptor tyrosine kinase²⁻⁵ and mutations of the *NPM1* gene encoding the nucleophosmin protein.⁶⁻⁸) In addition, mutations of the gene encoding CCAAT/enhancer-binding protein- α (*CEBPA*) have been described in patients with AML and normal karyotype.⁹⁻¹² The *CEBPA* gene encodes a transcription factor that serves as a master regulator of granulopoiesis.¹³ Targeted disruption of the gene in mice is associated with a block of granulocyte development and downregulation of target genes such as granulocyte colony-stimulating factor receptor.¹³ The intronless *CEBPA* gene on chromosome 19q13.1 encodes 2 major protein isoforms, the 42-kDa full-length protein (p42), which has the full transcriptional activity, and a shorter, 30-kDa isoform (p30), produced from a second, alternative start codon and shown to have complex functions, including an inhibitory effect on⁹ and increased degradation¹⁴ of the p42 full-length protein.

Since the initial report,⁹ several groups have investigated *CEBPA* mutations in AML.^{11,12,15-18} Besides the reproducible association with certain morphological and clinical features (eg, FAB M1 and M2 morphology, high CD34 expression on blasts, and predominance in normal karyotype), nearly all studies have shown that patients with *CEBPA* mutations have a more favorable outcome. However, more recent results have indicated that a good prognosis is confined to patients with biallelic or double mutations (*CEBPA*^{bi}),¹⁹⁻²¹ whereas patients with monoallelic *CEBPA* mutations (*CEBPA*sm) did not differ in their response to treatment from patients with wild-type *CEBPA* (*CEBPA*^{wt}) and had a less favorable outcome.¹⁹⁻²¹ This finding led to the inclusion of biallelic *CEBPA* mutations as an independent entity in the most recent World Health Organization classification,²² as well as a favorable prognostic group in the ELN2017 recommendations.²³ However, the impact of monoallelic *CEBPA* mutations has been investigated in more detail in only a few studies, especially in light of the different biological effects of N- and C-terminal mutations.^{21,24,25}

To investigate the prevalence and prognostic role of *CEBPA* mutations, in particular *CEBPA*sm mutations, in adult patients with AML, we studied 4708 patients with newly diagnosed AML. In contrast to many previous studies, we included patients of all age groups, as well as secondary AML (sAML) after prior myelodysplasia (MDS) or therapy-related AML (tAML). Because most *CEBPA* mutations reported so far consist of insertions or deletions,²⁶⁻²⁹ we used high-resolution fragment analysis for mutation screening and found *CEBPA* mutations in 5.1% of patients. The results of targeted next-generation sequencing (NGS) indicated profound differences in the co-mutational spectrum of individual *CEBPA* mutations. Our results point to a differential effect of *CEBPA*sm mutations that appear to be associated with similar clinical parameters, co-mutations, and outcome compared with *CEBPA*^{bi} mutations. If confirmed, these results, generated in one of the largest cohorts of patients with AML analyzed so far, could build the basis of a refined clinical classification of *CEBPA* mutations.

Patients, materials, and methods

Patients

We screened 4708 adult patients with newly diagnosed AML ($n = 3729$ with de novo AML; $n = 644$ with AML and a history of MDS; and $n = 335$ with tAML) for the presence of mutations in the *CEBPA* gene. Most individuals ($n = 3104$; 67.45%) were treated in prospective studies, including the AML96 ($n = 1457$), AML2003 ($n = 1081$), AML60⁺ ($n = 359$), and SORAML ($n = 207$) protocols of the Study Alliance Leukemia (SAL). The remaining patients ($n = 1604$) were recruited to the SAL registry and biorepository. Detailed treatment protocols have been published³⁰⁻³³ and are summarized in the supplemental Data (available on the Blood Web site), including the number of patients treated in each protocol. All studies involved risk-stratified consolidation therapy according to cytogenetic risk groups (*CEBPA* mutations were not used for risk stratification in any of those studies). Patients <60 years of age received standard cytosine arabinoside (Ara-C)/anthracycline (DA 3 + 7)-based, double-induction chemotherapy followed by consolidation with high-dose Ara-C in patients who had favorable risk. Patients in the intermediate- and high-risk groups had the option of upfront allogeneic transplantation and underwent autologous transplantation or chemotherapy-based consolidation in the absence of suitable donors. In older patients treated with curative intention, the chemotherapy regimen was adjusted according to predefined algorithms that integrate performance status and organ function.

This study was approved by the ethics board of the Technical University Dresden. Each patient gave written informed consent to participate in the respective study protocols.

Patient samples

All materials investigated were obtained at the time of diagnosis. Bone marrow was used whenever available; in all other cases, peripheral blood samples were examined. Genomic DNA was extracted from mononuclear cells by using standard procedures (DNA blood minikit; Qiagen, Hilden, Germany).

PCR for *CEBPA*

All polymerase chain reaction (PCR) analyses were performed on genomic DNA. Details of the PCR primers and cycling conditions are given in supplemental Tables 1 and 2 and supplemental Figure 1.

PCR-amplified mutant samples were purified and sequenced on an ABI3130xl instrument. Sequences were compared with the WT-*CEBPA* messenger RNA (mRNA) sequence (U34070).

NGS-based characterization of co-mutations in patients with *CEBPA* mutations

Profiling of mutations was achieved by targeted NGS-based resequencing with the TruSight Myeloid assay (Illumina, Chesterford, United Kingdom) covering 54 genes frequently mutated in AML, as described recently³⁴ (details in the supplemental Data). Data alignment of demultiplexed FastQ files, variant calling, and filtering were performed with the Sequence Pilot software package (JSI Medical Systems GmbH, Ettenheim, Germany), with default settings and a 5% variant allele frequency cutoff. *FLT3*-internal tandem duplication (ITD) and *NPM1* mutations were evaluated as reported previously.^{5,8}

RNA sequencing

RNA sequencing (RNA-Seq) was performed on total RNA isolated at diagnosis from 20 patients with a *CEBPA* mutation (5 *CEBPA*^{bi-inf}, 5 *CEBPA*^{sm-inf}, 5 *CEBPA*^{sm-other}, and 5 *CEBPA*^{bi-other}), by using strand-specific RNA-Seq library preparation (Ultra II Directional RNA Library Prep; New England Biolabs) and sequenced on an Illumina NovaSeq 6000 instrument. The complete workflow and the bioinformatic analyses are detailed in the supplemental Data.

Statistical analysis

Clinical variables across groups were compared by using the χ^2 test or 2-sided Fisher's exact test for categorical variables. The nonparametric Mann-Whitney *U* test was applied for continuous variables. *P* < .05 indicated a significant difference. Numerical variables are expressed as the median with interquartile range (IQR). Univariate analyses for the influence of the *CEBPA* mutational status on complete response (CR) rates were performed by using the χ^2 test. The log-rank test was used to evaluate OS and EFS. For multivariable analysis of prognostic factors, Cox proportional hazards regression models were used. Allogeneic hematopoietic stem cell transplantation (allo-HSCT) was modeled as a time-dependent covariate. *P*-values of association analyses of *CEBPA* mutations with clinical variables and other molecular abnormalities were adjusted for multiplicity by using the Bonferroni-Holm correction. All statistical analyses were performed with the SPSS software package, version 26 (SPSS, Chicago, IL) and R version 3.5.3 (<https://www.R-project.org/>).

Results

A total of 371 individual *CEBPA* mutations were identified in 240 of the 4708 patients (5.1%) by the screening procedure. In all patients with a mutation identified in any part of *CEBPA*, Sanger sequencing of the entire gene was performed to assess the presence of additional mutations, including single nucleotide variants, and to confirm the alterations identified using fragment analysis. Patients showing the previously described 6-bp polymorphism in TAD2²⁶ were included in the WT-*CEBPA* group.

As illustrated in supplemental Figure 2 and supplemental Table 3 and reported in previous studies, mutations largely clustered

in the N-terminal first transactivation domain (TAD1) and the DNA-binding and basic leucine zipper region (bZIP) in the C-terminal part of *CEBPA*. In our cohort, 131 of 240 patients (54.6%) presented with 2 *CEBPA* mutations, mostly consisting of combined mutations in bZIP and TAD, denoted as *CEBPA*^{bi}. *CEBPA*sm mutations were found in 109 of 240 patients (45.4%), of which 60 had N-terminal (*CEBPA*^{smTAD}) and 49 C-terminal (*CEBPA*^{smbZIP}) mutations. As reported before, most of the mutations in the TAD domains caused a frameshift, whereas mutations in the bZIP-region were predominantly in-frame insertions and duplications. Eight patients had monoallelic mutations with high variant allele frequency (4 *CEBPA*^{smTAD} and 4 *CEBPA*^{smbZIP}), indicating a homozygous state.

CEBPA mutations and clinical characteristics

The association of clinical parameters according to the localization of the *CEBPA* mutation (ie, patients with single N- or C-terminal mutations and patients with biallelic mutations) is summarized in Table 1. Compared with patients bearing *CEBPA*^{wt}, those with *CEBPA*^{bi} were significantly younger (median age, 46 years; IQR, 38-59) at diagnosis, similar to those with *CEBPA*^{smbZIP} (median age, 50 years; IQR, 39-57), whereas patients with *CEBPA*^{smTAD} were significantly older (median age, 63 years; IQR, 55-69.3) and were more comparable to the *CEBPA*^{wt} group (median, 57 years; IQR, 46-67; *P* < .001). When patients were categorized in 10-year age intervals (Figure 1), a continuous decrease in *CEBPA*^{bi} and *CEBPA*^{smbZIP} mutations was seen with increasing age, whereas *CEBPA*^{smTAD} alterations increased. In line with this finding, only a single patient with a *CEBPA*^{smTAD} mutation was observed in the group of patients <30 years of age.

As outlined in Table 1, patients with *CEBPA*^{smTAD} also had significantly lower WBC counts and significantly lower rates of CD34 positivity, compared with the *CEBPA*^{bi} and *CEBPA*^{smbZIP} groups, and more commonly had prior myelodysplasia syndrome (MDS) or tAML that evolved as a secondary disease.

Association of CEBPA mutations with other molecular abnormalities

Based on the targeted NGS approach, additional mutations were identified in 208 of 240 patients with a *CEBPA*




Figure 1. Age distribution of the 240 *CEBPA* mutations (*CEBPA*^{bi}, *CEBPA*^{smTAD}, *CEBPA*^{smbZIP}) identified in this study.

Table 1. Clinical and cytogenetic characteristics *CEBPA*^{wt} compared with *CEBPA*^{bi}, *CEBPA*^{smbZIP} and *CEBPA*^{smTAD} mutant patients

	<i>CEBPA</i> ^{wt} n = 4468	<i>CEBPA</i> ^{smTAD} n = 60	<i>CEBPA</i> ^{smbZIP} n = 49	<i>CEBPA</i> ^{bi} n = 131	P (adj.)
Age, y, median (IQR)	57 (46-67)	63 (55-69)	50 (39-57)	46 (38-59)	<.001
Sex, n (%)					
Female	2170 (49)	32 (53)	16 (33)	70 (53)	.26
Male	2298 (51)	28 (47)	33 (67)	61 (47)	
AML type, n (%)					
de novo	3510 (78)	50 (84)	47 (96)	122 (93)	.001
sAML	630 (14)	8 (13)	2(4)	4 (3)	
tAML	328 (7)	2 (3)	—	5 (4)	
Laboratory, median (IQR)					
BM blasts, %	60 (38-80)	60 (43.3-72.8)	65.3 (43.9-80)	64 (50-87)	.823
CD34-positivity, %	16 (2-47)	19.2 (3.8-65)	45 (19.5-68)	55.6 (32.9-77)	<.001
WBC, ×10 ⁹ /L	11.7 (3-45.7)	13.2 (3.3-51.2)	35.7 (11.5-93)	23.7 (9.3-64.7)	<.001
LDH, U/L	412.9 (272-727)	412 (291-569)	491 (344-941)	445 (292-679)	.99
FAB, n (%)					
M0	223 (5)	—	1 (2)	—	<.001
M1/M2	2040 (46)	40 (67)	39 (80)	101 (77)	
M4-M7	1535 (34)	15 (25)	8 (16)	19 (15)	
Unknown	670 (15)	5 (8)	1 (2)	10 (8)	
Cytogenetics, n (%)					
Normal karyotype	1923 (43)	37 (62)	34 (72)	102 (78)	<.001
Aberrant karyotype	2359 (53)	23 (38)	15 (28)	29 (22)	
Unknown	186 (4)	—	—	—	
Risk, n (%)					
Favorable risk (MRC)	353 (8)	1 (2)	1 (2)	—	<.001
Intermediate risk (MRC)	2928 (68)	54 (90)	43 (92)	119 (98)	
Adverse risk (MRC)	1001 (23)	5 (8)	3 (6)	2 (2)	
Treatment, n (%)					
Primary allo-HCT	747 (17)	3 (5)	10 (20)	36 (27)	<.001
Salvage allo-HCT	915 (20)	13 (22)	8 (16)	22 (17)	.189

Bold P-values indicate statistically significant results.

BM, bone marrow; FAB, French-American-British; LDH, lactate dehydrogenase; MRC, medical research council.

mutation (86.7%). The most frequently mutated genes were *TET2* (70 of 240; 29.2%), *GATA2* (68 of 240; 28.3%), *DNMT3A* (45 of 240; 18.8%), *FLT3*-ITD (39 of 240; 16.3%), *NPM1* and *NRAS* (31 of 240; 12.9% each), and *WT1* (30 of 240; 12.5%).

Figure 2 illustrates the distribution of co-mutations in the 3 *CEBPA* subgroups. Significant differences were observed for several genes, the most striking being *GATA2*, which was mutated in 35.1% of patients with *CEBPA*^{bi}, 36.7% of those with *CEBPA*^{smbZIP}, but in only 6.7% of those with *CEBPA*^{smTAD} ($P = .001$), and *NPM1*, which was mutated in 3.1% of patients with

CEBPA^{bi} and 8.2% of those with *CEBPA*^{smbZIP}, but in 38.3% of patients with *CEBPA*^{smTAD} ($P < .001$). Significant differences were also found for mutations in *DNMT3A*, *FLT3*-TKD, *IDH1*, and *IDH2*, as well as in *SRSF2* and *WT1* (Figure 2B). In general, the spectrum of mutations of *CEBPA*^{smbZIP} was more comparable to that of patients with *CEBPA*^{bi} and differed markedly from the *CEBPA*^{smTAD} group, the latter being more similar to patients with *CEBPA*^{wt}, who frequently carried mutations in genes associated with AML after prior MDS, such as spliceosome mutations (ie, *SRSF2*, *SF3B1*, or *U2AF1*) and alterations associated with DNA methylation (ie, *DNMT3A*, *ASXL1*, *TET2*, *IDH1*, and *IDH2*). Interestingly, only 7 patients in this analysis




Figure 2.




Figure 3. Survival analysis according to CEBPA mutation status (CEBPA^{wt}, CEBPA^{bi}, and CEBPAsm) within the cohort of 4461 patients receiving intensive treatment. Kaplan-Meier plots showing OS (A) and EFS (B). Numbers below the figures denote the patients at risk. The tables provide the results of pairwise univariate analyses.

had mutations in *CSF3R*, with no significant differences found between the groups (3.1% CEBPA^{bi}, 2% CEBPAsm, and 3.3% CEBPA^{wt}).

CEBPA mutations and outcome

The prognostic relevance of CEBPA mutations was analyzed in 4461 intensively treated patients, with a median follow-up time of patients remaining alive of 61 months (IQR, 36-96 months). When analyzed according to the current recommendations (ie, separating only CEBPA^{bi} and CEBPAsm), only patients with CEBPA^{bi} had a significantly higher rate of complete remission (CR; CEBPA^{bi}, 95% vs CEBPAsm, 75% and CEBPA^{wt}, 73%; $P < .001$), as well as a higher median OS (CEBPA^{bi}, 103.2 months; 95% confidence interval [CI], 70.7-inf vs CEBPAsm, 21.9 months; 95% CI, 12.7-54, and CEBPA^{wt}, 19.3 months; 95% CI, 17.9-21; $P < .001$) and EFS (CEBPA^{bi}, 20.7 months; 95% CI, 13.1-101.1 vs CEBPAsm, 9.4 months; 95% CI, 5.7-15.3, and CEBPA^{wt}, 7.0 months; 95% CI, 6.5-7.6; $P < .001$; Figure 3), which was confirmed in multivariable analyses (Table 2).

Given the marked differences in the clinical and molecular alterations associated with the different localization of single-mutant CEBPA, we also looked for the impact of the localization of CEBPAsm on outcome. Patients carrying CEBPAsm showed a

significantly higher CR rate (CEBPA^{bi} 95% vs CEBPAsm 86% vs CEBPA^{wt} 73%) and a significantly longer OS (CEBPA^{bi} 103.2 months; 95% CI, 70.7-inf, vs CEBPAsm, 63.3 months; 95% CI, 20.5-inf, vs CEBPA^{wt}, 17.9 months; 95% CI, 17.9-21) and EFS (CEBPA^{bi}, 20.7 months; 95% CI, 13.1-101.1 vs CEBPAsm, 9.4 months; 95% CI, 5.7-15.3, vs CEBPA^{wt}, 7.0 months; 95% CI, 6.5-7.6; Figure 4A-B). Multivariable analysis confirmed that CEBPAsm mutations represented an independent favorable risk for outcome (Table 3).

Effect of co-mutations on outcome

We compared the effect of co-mutations on the survival of the patients according to the mutational status of the 3 most common mutations previously associated with outcome (ie, *GATA2*, *TET2*, and *WT1* in the 3 CEBPA mutation subgroups (CEBPA^{bi}, CEBPAsm, and CEBPA^{wt}). Although the presence of *GATA2* mutations was associated with improved OS and EFS in the CEBPA^{bi} and CEBPAsm groups (Figure 5A-B); however, these differences were not significant in pairwise comparisons (all P -values were adjusted for multiple testing with the Bonferroni-Holm procedure). However, for *TET2* (Figure 5B-C) a significant difference was found between CEBPA^{bi}/TET2^{wt}

Figure 2. Alignment of additional gene mutations for 240 patients with CEBPA^{mut}. (A) Co-mutations organized by categories of related genes, as labeled on the left. Patients are shown in order by CEBPA subgroup. The heat map includes all mutated genes in patients with CEBPA^{bi}, CEBPAsm, or CEBPA^{wt}. Each column represents one of the 240 analyzed samples. Mutations in the investigated genes are shown by black bars, light gray bars indicate WT status. (B) Frequency distribution of additional gene mutations identified in patients with CEBPA^{bi}, CEBPAsm, or CEBPA^{wt} mutations (frequency of at least 10% in 1 subgroup). RAS signaling including *KRAS*, *NRAS*, *PTPN11*, and *CBLB*; spliceosome, including *SF3B1*, *SRSF2*, and *ZRSR2*; and methylation, including *DNMT3A*, *IDH1*, *IDH2*, and *TET2*.

Table 2. Results of the multivariable analysis for CEBPA^{bi} vs CEBPAsm

	OS (HR)	P	95% CI	EFS (HR)	P	95% CI	CR1 (OR)	P	95% CI
Intermediate karyotype	1			1			1		
Favorable karyotype	0.427	<.001	(0.35-0.52)	0.400	<.001	(0.34-0.47)	3.063	<.001	(2.07-4.54)
Adverse karyotype*	1.858	<.001	(1.68-2.05)	1.633	<.001	(1.49-1.79)	0.477	<.001	(0.40-0.57)
Age	1.032	<.001	(1.03-1.04)	1.023	<.001	(1.02-1.03)	0.948	<.001	(0.94-0.95)
Log ₁₀ WBC	1.155	<.001	(1.08-1.24)	1.228	<.001	(1.15-1.31)	0.754	<.001	(0.66-0.87)
De novo AML	1			1			1		
sAML	1.086	.131	(0.98-1.21)	1.069	.196	(0.97-1.18)	0.733	.002	(0.60-0.90)
tAML	1.386	<.001	(1.20-1.60)	1.098	.186	(0.96-1.26)	0.691	.009	(0.52-0.91)
No FLT3-ITD ^{mut}	1			1			1		
FLT3-ITD ^{mut}	1.206	<.001	(1.09-1.33)	1.242	<.001	(1.14-1.36)	1.017	.870	(0.83-1.25)
No NPM1 ^{mut}	1			1			1		
NPM1 ^{mut}	0.629	<.001	(0.57-0.69)	0.535	<.001	(0.49-0.59)	2.127	<.001	(1.75-2.59)
CEBPA ^{wt}	1			1			1		
CEBPA ^{bi}	0.510	<.001	(0.39-0.67)	0.534	<.001	(0.42-0.68)	6.328	<.001	(2.73-14.67)
CEBPA sm	0.825	.122	(0.65-1.05)	0.746	.011	(0.56-0.94)	1.117	.644	(0.70-1.78)
No allo-HSCT in CR1	1			1			1		
Allo-HSCT in CR1	0.773	<.001	(0.69-0.87)	0.630	<.001	(0.56-0.79)	—	—	—

The multivariable analysis performed includes the different study regimens as strata.

*Adverse karyotype according to ELN 2017.




Figure 4. Survival analysis according to CEBPA mutation status (CEBPA^{bi}, CEBPA^{smTAD}, CEBPA^{smbZIP}, and CEBPA^{wt}) within the cohort of 4461 patients receiving intensive treatment. Kaplan-Meier plots showing OS (A) and EFS (B). Numbers of patients at risk and results of log-rank tests from pairwise comparisons are provided below the x-axis.

and TET2^{mut} (median OS: CEBPA^{bi}/TET2^{wt}, not reached; 95% CI, 101.1-inf, vs CEBPA^{bi}/TET2^{mut}, 22.5 months; 95% CI, 13.9-inf; $P = .012$; median EFS: CEBPA^{bi}/TET2^{wt}, 34.7 months; 95% CI, 13.5-inf vs CEBPA^{bi}/TET2^{mut}, 9.5 months; 95% CI, 5.23-50.2; $P = .02$) and a trend for a different EFS in patients with CEBPA^{smbZIP}/TET2^{wt} or CEBPA^{smbZIP}/TET2^{mut} (median EFS CEBPA^{bi}/TET2^{wt}, not reached; 95% CI, 9.59-inf vs CEBPA^{bi}/TET2^{mut}, 9.2 months; 95% CI, 2.40-66.2; $P = .06$). WT1 mutations were associated with lower probability of survival (supplemental Figure 4); however, the comparison between the groups did not reach statistical significance.

NPM1 mutations, the most common co-mutation in patients carrying CEBPA^{smTAD}, were associated with slightly better OS and EFS; however, these differences were also not significant (supplemental Figure 5).

Presence of "typical" bZIP-mutations and outcome

Mutations in the CEBPA-bZIP region are typically in-frame (ie, multiples of 3 bp) and affect the DNA-binding-, fork-, or bZIP-region (for simplicity, summarized as bZIP) between amino acid positions 278 and 345 of the CEBPA protein (supplemental Figure 3), whereas frameshift mutations, the hallmark of mutations affecting the TAD1 and TAD2 domains, are less common in C-terminal mutations. To investigate whether the presence of typical in-frame bZIP mutations (irrespective of the biallelic or monoallelic status) actually represents the decisive molecular factor for the favorable outcome observed, we regrouped the 240

patients with CEBPA^{mut} according to the presence or absence of these mutations. Typical bZIP mutations according to this classification were found in 118 of 131 who had CEBPA^{bi} and 39 of 49 of those with CEBPA^{smbZIP} (denoted CEBPA^{bi-inf} and CEBPA^{sm-inf}). Thirteen of 131 CEBPA^{bi} had nontypical biallelic mutations (eg, consisting of 2 different TAD domain mutations or a combination of TAD mutations and frameshift or nonsense bZIP mutations, denoted CEBPA^{bi-other}). Patients with mutation in TAD as well as patients with a nontypical single-allele bZIP mutation were grouped as "other single-allele mutations" (CEBPA^{sm-other}). We again observed profound differences in the age distribution, with typical bZIP mutations (double and single allele) predominating in younger adults, whereas other mutations, especially the few nontypical biallelic mutations, were predominantly detected in patients >50 years of age (supplemental Table 4; supplemental Figure 6).

To further characterize potential biological similarities between CEBPA^{bi-inf} and CEBPA^{sm-inf}, we performed RNA-seq of 20 samples from patients with CEBPA^{bi-inf}, CEBPA^{sm-inf}, CEBPA^{bi-other}, and CEBPA^{sm-other} (5 samples per group). As illustrated in the volcano-plots in supplemental Figure 7, we found between 34 and 129 differentially expressed genes in pairwise comparisons between all groups, the only exception being CEBPA^{bi-inf} and CEBPA^{sm-inf}, where no significant differences were found at a false discovery rate of 5% (supplemental Figure 7C).

The survival analysis performed also revealed profound differences. Superimposable, favorable outcomes (OS and EFS) were

Table 3. Results of the multivariable analysis for CEBPA^{bi} vs CEBPA^{smbZIP} vs CEBPA^{smTAD}

	OS (HR)	P	95% CI	EFS (HR)	P	95% CI	CR1 (OR)	P	95% CI
Intermediate karyotype	1			1			1		
Favorable karyotype	0.426	<.001	(0.35-0.51)	0.400	<.001	(0.34-0.47)	3.076	<.001	(2.08-4.56)
Adverse karyotype*	1.853	<.001	(1.68-2.04)	1.625	<.001	(1.49-1.78)	0.476	<.001	(0.40-0.58)
Age	1.032	<.001	(1.03-1.04)	1.023	<.001	(1.02-1.03)	0.948	<.001	(0.94-0.95)
Log ₁₀ WBC	1.158	<.001	(1.08-1.24)	1.231	<.001	(1.16-1.31)	0.751	<.001	(0.66-0.86)
De novo AML	1			1			1		
sAML	1.088	.127	(0.98-1.21)	1.069	.189	(0.97-1.18)	0.735	.003	(0.60-0.90)
tAML	1.380	<.001	(1.19-1.60)	1.098	.186	(0.96-1.26)	0.697	.009	(0.53-0.91)
No FLT3-ITD ^{mut}	1			1			1		
FLT3-ITD ^{mut}	1.204	<.001	(1.09-1.33)	1.238	<.001	(1.13-1.35)	1.018	.866	(0.83-1.25)
No NPM1 ^{mut}	1			1			1		
NPM1 ^{mut}	0.625	<.001	(0.57-0.69)	0.531	<.001	(0.49-0.58)	2.151	<.001	(1.77-2.62)
CEBPA ^{wt}	1			1			1		
CEBPA ^{bi}	0.507	<.001	(0.39-0.67)	0.530	<.001	(0.42-0.67)	6.367	<.001	(2.75-14.77)
CEBPA ^{smbZIP}	0.620	.019	(0.42-0.92)	0.537	.001	(0.37-0.77)	1.876	.139	(0.81-4.32)
CEBPA ^{smTAD}	1.024	.878	(0.75-1.39)	0.981	.891	(0.74-1.31)	0.825	.515	(0.46-1.48)
No allo-HSCT in CR1	1			1			1		
Allo-HSCT in CR1	0.776	<.001	(0.69-0.87)	0.632	<.001	(0.56-0.71)	—	—	—

The multivariable analysis performed includes the different study regimens as strata.

*Adverse karyotype according to ELN 2017.




Figure 5. Survival analysis according to CEBPA and GATA2 mutational status within the cohort of 4461 patients receiving intensive treatment. Kaplan-Meier plots showing OS (A) and EFS (B) of CEBPA^{wt}, compared with CEBPA^{bi}/GATA2^{wt} (n = 82), CEBPA^{bi}/GATA2^{mut} (n = 46), CEBPA^{smbZIP}/GATA2^{wt} (n = 31), CEBPA^{smbZIP}/GATA2^{mut} (n = 18) mutant cases. Survival analysis according to CEBPA mutation status and TET2 status within a cohort of 4461 patients. Kaplan-Meier plots showing OS (C) and EFS (D) of CEBPA^{wt}, compared with CEBPA^{bi}/TET2^{wt} (n = 98), CEBPA^{bi}/TET2^{mut} (n = 30), CEBPA^{smbZIP}/TET2^{wt} (n = 33), CEBPA^{smbZIP}/TET2^{mut} (n = 16) mutant cases. Numbers of patients at risk and results of log-rank tests from pairwise comparisons are provided below the x-axis.




Figure 6. Grouping of CEBPA mutations according to localization and type.

confined to patients in the CEBPA^{bi-inf} and the CEBPA^{sm-inf} groups, whereas patients with other mutational constellations, in particular patients with nontypical biallelic CEBPA mutations, showed an inferior outcome (supplemental Table 5; supplemental Figure 8). Co-mutations retained their prognostic impact in patients carrying CEBPA^{bi-inf} and CEBPA^{sm-inf} (as illustrated for the most common GATA2 mutations in supplemental Figure 9).

Prognostic classification for CEBPA mutations

Because these results strongly support the notion that mutation constellations containing typical in-frame bZIP mutations (ie, CEBPA^{bi-inf} and CEBPA^{sm-inf}) are comparable with respect to most clinical and molecular factors studied, we combined these 2 mutation types (denoted CEBPA^{bZIP-inf}) and compared this new group to those patients without these mutations (CEBPA^{other}; allocation summarized in Figure 6).

As illustrated in Table 4, the clinical associations already observed for the individual groups were even more pronounced in this analysis, especially the highly divergent median age (46 vs 62 years; $P < .001$), the difference with respect to the association with sAML (4% vs 17%; $P < .001$), the median CD34 positivity (51% vs 34%; $P < .001$), and the median WBC counts ($25 \times 10^9/L$ vs $14.7 \times 10^9/L$; $P < .001$). As expected, the mutational spectrum of patients belonging to these 2 subgroups showed highly significant differences. Only 1 of 157 patients with

CEBPA^{bZIP-inf} (0.6%) showed an NPM1 mutation, compared with 30 of 83 patients with CEBPA^{other} mutations (36.1%; $P < .001$; Figure 7C; supplemental Figure 9). Other mutations that were significantly more common in the CEBPA^{other} subgroup were alterations affecting DNA methylation (43 of 157, 27.4% vs 60 of 83, 72.3%; $P < .001$) and the spliceosome (4 of 157, 2.6% vs 21 of 83, 25.3%; $P < .001$).

Outcome analyses performed for CEBPA^{bZIP-inf} patients confirmed a strong prognostic impact of this subgroup, whereas CEBPA mutations in the other patients did not show significant differences compared with CEBPA^{wt} (Figure 7A-B). Interestingly, patients with CEBPA^{bZIP-inf} appear not to benefit from HSCT performed in the first complete remission (CR1; supplemental Figure 10). Multivariable analysis performed for this classification confirmed that CEBPA^{bZIP-inf} positivity was the strongest predictor for achievement of CR (OR: 6.06; 95% CI, 2.78-13.23, $P < .001$) and a strong prognostic factor for OS (hazards ratio [HR], 0.57; 95% CI, 0.46-0.71; $P < .001$) and EFS (HR, 0.53; 95% CI, 0.43-0.64, $P < .001$; Table 5). We also looked for the effect of mutant GATA2, TET2, and WT1 on outcome in the novel subgroups. As observed in the previous analyses, mutant GATA2 and the absence of TET2 and WT1 mutations were associated with improved OS and EFS in patients carrying CEBPA^{bZIP-inf} but not in those with CEBPA^{other}, although these differences were not significant (supplemental Figure 11).

Table 4. Clinical variables in patients with *CEBPA*^{bZIP-inf} and *CEBPA*^{other} mutations

	<i>CEBPA</i> ^{wt} n = 4468	<i>CEBPA</i> ^{bZIP-inf} n = 157	<i>CEBPA</i> ^{other} n = 83	P (adj.)
Age in years, median (IQR)	57 (46-67)	46 (36-57)	62 (55-69)	<.001
Sex, n (%)				
Female	2170 (49)	76 (48)	42 (51)	.885
Male	2298 (51)	81 (52)	41 (49)	
AML type, n (%)				
De novo	3510 (78)	149 (96)	69 (83)	<.001
sAML	630 (14)	5 (3)	9 (11)	
tAML	328 (7)	2 (1)	5 (6)	
FAB subtype, n (%)				
M0	223 (5)	1 (1)	/	<.001
M1/M2	2040 (46)	127 (81)	53 (65)	
M4-M7	1535 (34)	22 (14)	20 (24)	
Unknown	670 (15)	7 (4)	9 (11)	
Laboratory, median (IQR)				
BM blasts in %	60 (38-80)	64 (50-78)	62 (43.5-78)	.853
CD34-positivity in %	16 (2-47)	51 (29-75.2)	34 (6.6-72.2)	<.001
WBC in 10 ⁹ /l	11.7 (3-45.7)	25 (9.2-70)	14.7 (4.6-53.4)	.001
LDH in U/l	412.9 (272-727)	445 (295-758)	449 (296-601)	.242
Treatment, n (%)				
Primary allo-HCT	747 (17)	52 (33)	5 (6)	<.001
Salvage allo-HCT	915 (20)	21 (13)	16 (19)	.209

Bold P-values indicate statically significant results.

BM, bone marrow; LDH, lactate dehydrogenase; FAB, French-American-British; MRC, medical research council.

Discussion

We analyzed 4708 adult patients with newly diagnosed AML for *CEBPA* alterations and identified mutations in *CEBPA* in 5.1%. Our cohort differed from patient cohorts in previous studies investigating the role of *CEBPA* mutations,^{9-12,15,21,24,35-42} because we did not select for age, cytogenetic subgroups, or disease status. Within the entire cohort, 49% showed an aberrant karyotype, and the median patient age at diagnosis was 57 years (IQR, 46-67 years), more accurately reflecting the entirety of patients with AML in general. This fact may explain why the prevalence of *CEBPA* mutations found in our study is at the lower end of the previously reported range of 4% to 20%.^{9-12,15,21,24,35-42}

Our main focus was the role of monoallelic *CEBPA* mutations; therefore, we performed an extensive evaluation of this subgroup and a detailed analysis of the individual type and localization of the mutation.

A first aspect observed with respect to the different mutational subgroups (*CEBPA*^{bi}, *CEBPA*^{smbZIP}, and *CEBPA*^{smTAD}) was the highly significant difference in the age distribution. Whereas *CEBPA*^{bi} and *CEBPA*^{smbZIP} were predominantly found in younger adults and decreased with age, *CEBPA*^{smTAD} mutations were rare in patients up to the age of 40 years and were

particularly common in older individuals. The lower age of patients with *CEBPA*^{bi} mutations has already been reported in previous studies.^{21,24,41} In contrast, the major age difference between *CEBPA*^{smTAD} and *CEBPA*^{smbZIP} mutations has not been reported before. Interestingly, a very recently published work in pediatric AML found only monoallelic *CEBPA*^{bZIP} mutations in their analysis, indicating that these mutations are indeed significantly more prevalent in younger individuals.⁴³ *CEBPA*^{bi} and *CEBPA*^{smbZIP} also showed overlapping laboratory profiles, with higher rates of CD34 positivity and higher WBC counts.⁴³ We also observed a favorable prognostic impact of *CEBPA*^{smbZIP} comparable to biallelic *CEBPA* mutations. This finding again is in concordance with the data reported by Tarlock et al in pediatric AML, showing that the outcome of patients with *CEBPA*^{bi} or *CEBPA*^{smbZIP} mutations did not differ.⁴³

Our results indicate a similar spectrum of co-mutations for *CEBPA*^{bi} and *CEBPA*^{smbZIP} (ie, significantly higher rates of *GATA2* and *WT1* mutations and a mere lack of *NPM1* alterations). *GATA2* mutations in *CEBPA*^{bi} have been described previously,⁴⁴ whereas a similar association of *CEBPA*^{smbZIP} with *GATA2* mutations has not been reported so far. The presence of *GATA2*, *TET2*, and *WT1* mutations significantly affected the outcome of patients carrying *CEBPA*^{bi}, whereas *CEBPA*^{smbZIP} mutations showed a significant effect on outcome only for mutations in *GATA2*. The prognostic impact of mutations in *GATA2*,




Figure 7. Survival analysis according to CEBPA mutation status (CEBPA^{wt}, CEBPA^{bZIP-inf}, and CEBPA^{others}) within the cohort of 4461 patients receiving intensive treatment. Kaplan-Meier plots showing OS (A) and EFS (B). (C) Frequency distribution of additional gene mutations identified in patients with CEBPA^{bZIP-inf} and CEBPA^{others} mutant genes (frequency of at least 5% in one subgroup). RAS signaling, including KRAS, NRAS, PTPN11, CBLB; spliceosome, including SF3B1, SRSF2, and ZRSR2; and methylation, including DNMT3A, IDH1, IDH2, and TET2. Numbers of patients at risk and results of log-rank tests from pairwise comparisons are provided below the x-axis.

TET2, and WT1 has been described recently,^{45,46} but was confined to biallelic CEBPA mutations. In contrast, our data indicate that these prognostic associations also apply to patients with

CEBPA^{smbZIP}, but not to those with CEBPA^{smTAD}. Clustering of co-mutations according to functional groups further highlighted a differential spectrum of co-occurring mutations, with

Table 5. Results of the multivariable analysis for CEBPA^{bZIP-inf} vs CEBPA^{other}

	OS (HR)	P	95% CI	EFS (HR)	P	95% CI	CR1 (OR)	P	95% CI
Intermediate karyotype	1			1			1		
Favorable karyotype	0.456	<.001	(0.38-0.55)	0.436	<.001	(0.38-0.51)	3.072	<.001	(2.07-4.55)
Adverse karyotype*	1.823	<.001	(1.67-2.00)	1.590	<.001	(1.46-1.73)	0.478	<.001	(0.40-0.58)
Age	1.033	<.001	(1.03-1.04)	1.025	<.001	(1.02-1.03)	0.948	<.001	(0.94-0.95)
Log ₁₀ WBC	1.159	<.001	(1.09-1.24)	1.228	<.001	(1.16-1.31)	0.753	<.001	(0.66-0.86)
De novo AML	1			1			1		
sAML	1.050	.356	(0.95-1.16)	1.020	.683	(0.93-1.12)	0.734	.003	(0.60-0.90)
tAML	1.312	<.001	(1.15-1.50)	1.056	.413	(0.93-1.20)	0.704	.012	(0.53-0.93)
No FLT3-ITD ^{mut}	1			1			1		
FLT3-ITD ^{mut}	1.179	<.001	(1.08-1.29)	1.193	<.001	(1.10-1.30)	1.032	.767	(0.84-1.27)
No NPM1 ^{mut}	1			1			1		
NPM1 ^{mut}	0.635	<.001	(0.58-0.70)	0.553	<.001	(0.51-0.60)	2.139	<.001	(1.76-2.60)
CEBPA ^{wt}	1			1			1		
CEBPA ^{bZIP-inf}	0.570	<.001	(0.46-0.71)	0.525	<.001	(0.43-0.64)	6.061	<.001	(2.78-13.23)
CEBPA ^{other}	0.897	.410	(0.69-1.16)	0.964	.762	(0.76-1.22)	1.003	.989	(0.61-1.65)
No allo-HSCT in CR1	1			1			1		
Allo-HSCT in CR1	0.671	<.001	(0.60-0.75)	0.446	<.001	(0.40-0.50)	—	—	—

The multivariable analysis performed includes the different study regimens as strata.

CR1, first complete remission.

*Adverse karyotype according to ELN 2017.

CEBPA^{smTAD} mutations showing a significantly higher prevalence of mutations affecting proteins involved in DNA methylation (ie, DNMT3A, TET2, IDH1, and IDH2) as well as RNA splicing (ie, SRSF2, SF3B1, and ZRSR2), which was not seen in patients with CEBPA^{bi} and CEBPA^{smbZIP} (Figure 4B). Additional analyses looking in more detail for the individual mutations in CEBPA in our patients suggested that the identified clinical and molecular associations as well as the association with outcome were restricted to typical in-frame mutations within the bZIP region.

The reason for this differential behavior is unclear at present, but based on previous in vitro as well as animal experiments, TAD1 and bZIP mutations have clear functional differences (reviewed in Pulikkan et al⁴⁷). The frameshift mutations typically observed in the TAD1 domain induce the consecutive translation of the shorter CEBPA^{p30} protein, instead of the CEBPA^{p42} full-length protein. In contrast, with both CEBPA^{p42} and CEBPA^{p30}, mutations in the DBD and bZIP domains result in loss of DNA binding as well as dimerization. Recent animal data modeling the disease suggest functional disparity of different bZIP mutations. Lethally irradiated mice undergoing transplant of hematopoietic stem cells homozygous for a point mutation in bZIP (designated BRM2) start to develop a myeloproliferative disease that transforms into overt AML.⁴⁸ By transplanting transgenic cells carrying the most common mutation in bZIP (K313dup; K allele), alone or in combination with a TAD1 mutation (designated L-allele), Bereshchenko et al⁴⁹ documented that cells from mice carrying either the K/K or the K/L genotype, showed similarities in their mRNA expression profiles and higher expansion of immature blasts in the BM, which was distinct from the L/L genotype, as well as CEBPA-WT cells. CEBPA-mutated AML is characterized by specific RNA,⁵⁰ as well as miRNA^{35,51} expression profiles. The expression of several key miRNAs, such as *miR-34*, *miR-182*, and *miR-223*, which are involved in stem cell self-renewal, cell migration, and granulocytic differentiation, are physiologically regulated by CEBPA (reviewed in Stavast et al⁵²). Interestingly, recent data suggest that bZIP-mutant CEBPA does not downregulate *miR-182*, leading to a block of granulocytic differentiation.⁵³ This incapability appears to be mainly restricted to typical in-frame bZIP mutations clustering around the core mutated amino acids 312 and 313 of the CEBPA protein.⁵³ In support of this, one of the genes we found most highly deregulated between CEBPA^{bi-inf}/CEBPA^{sm-inf} and CEBPA^{sm-other}, *OSTL/RNF217*, coding for a highly conserved RING-finger ubiquitin ligase overexpressed in various leukemia entities,⁵⁴ has been shown to be regulated by the miRNA cluster *miR-183-96-182*.⁵⁵

Taken together, the CEBPA^{bZIP-inf} genotype identified in our analysis describes a subgroup of CEBPA mutations predominantly found in younger adults, indicating characteristics of a more immature and proliferative disease that has an overall prevalence of 3.3% in adult patients with AML, but is found in up to 7% of patients ≤40 years, thus representing a relevant subgroup, especially in younger adults. The mere absence of *NPM1* mutations (1 of 157 patients; 0.6%) and the high prevalence of *GATA2* mutations in these patients, which was found to be associated with an even better prognosis and a long-term survival in up to 80%, further highlights the special biology of these leukemias. The 2016 World Health Organization classification defines the CEBPA mutational class by the presence of a biallelic mutation, regardless of the localization within the gene. Given that

90% of the patients with the biallelic CEBPA mutation in this analysis actually carried typical bZIP CEBPA mutations, there is obviously a high degree of overlap, which may explain why this difference was undetected in most previous analyses, although there was some evidence in another study.²⁵ However, that only the 90% of patients with biallelic CEBPA mutations containing typical bZIP mutations in fact show a better outcome indicates that only those should be assigned to the favorable risk group.

Acknowledgments

The authors thank all centers and participating physicians of the Study Alliance Leukemia who entered their patients into the study (see Appendix). They also thank M. Böhm and M. Hartwig for skillful technical assistance.

This study was supported in part by grants from the Federal Ministry of Education and Research 01GS0872 (C.T.).

Authorship

Contribution: F.T., J.A.G., and C.T. designed the study; performed the research; collected, assembled, analyzed, interpreted the data; and wrote the manuscript; F.T., J.A.G., S. Stasik., A.P., R.M.-L., and P.J.M.V., performed the molecular analyses and analyzed the data; S.H. performed the molecular analyses; M. Kramer and J.S. performed the statistical analyses; J.M.M., C.R., U.K., A.K., S. Scholl, A.H., T.H.B., R. Naumann, B.S., H.E., M.S., A.B., A.N., K.S.-E., C.S., S.W.K., M.H., R. Nopenney., U.K., C.D.B., M. Kaufmann, F.S., K.S., M.v.B., C.M.-T., U.P., W.E.B., H.S., G.E., M.B., and J.S., treated the patients and collected the clinical data; and all authors approved the final version of the manuscript.

Conflict-of-interest disclosure: C.T. is CEO and co-owner of AgenDix GmbH. The remaining authors declare no competing financial interests.

ORCID profiles: S.S., 0000-0003-0271-4894; C.R., 0000-0002-3791-0548; T.H.B., 0000-0002-9677-3723; R.M.-L., 0000-0003-0298-7948; A.B., 0000-0002-8411-8844; A.N., 0000-0002-7606-8760; S.K., 0000-0002-5259-4651; U.P., 0000-0003-1863-3239; W.E.B., 0000-0002-3030-6567; H.S., 0000-0001-8472-5516; M.B., 0000-0002-5916-3029; J.S., 0000-0002-2780-2981; C.T., 0000-0003-1241-2048.

Correspondence: Christian Thiede, Medizinische Klinik und Poliklinik I, Universitätsklinikum Carl Gustav Carus der Technischen Universität, Fetscherstr 74, 01307 Dresden, Germany; e-mail: christian.thiede@uniklinikum-dresden.de.

Footnotes

Submitted 16 February 2021; accepted 18 July 2021; prepublished online on *Blood* First Edition 28 July 2021. DOI 10.1182/blood.202009680.

*F.T. and J.A.G. contributed equally to this study.

Original data can be obtained by e-mail request to the corresponding author (christian.thiede@uniklinikum-dresden.de).

The online version of this article contains a data supplement.

There is a *Blood* Commentary on this article in this issue.

The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked "advertisement" in accordance with 18 USC section 1734.

Appendix: participating centers of the Study Alliance Leukemia (SAL)

Tim H. Brümmendorf, Universitätsklinikum Aachen der RWTH, Germany; Armin Schulz-Abelius, Klinikum Altenburger Land GmbH, Altenburg, Germany; Martin Trepel, Klinikum Augsburg, Augsburg, Germany; Martina Teichmann, Sozialstiftung Bamberg, Bamberg, Germany; Alexander Kiani, Klinikum Bayreuth GmbH, Bayreuth, Germany; Bertram Glass, Helios Klinikum Berlin-Buch, Berlin, Germany; Martin Görner, Städt Kliniken Bielefeld gGmbH, Bielefeld, Germany; Dirk Behringer, Augusta-Kranken-Anstalt gGmbH, Bochum, Germany; Johannes Kullmer, Ev Diakonie-Krankenhaus gGmbH, Bremen, Germany; Mathias Hänel, Klinikum Chemnitz gGmbH, Chemnitz, Germany; Christof Lamberti, MVZ Coburg, Coburg, Germany; Martin Schmidt-Hieber, Carl-Thiem-Klinikum Cottbus gGmbH, Cottbus, Germany; Christoph Röllig, Universitätsklinikum Dresden, Dresden, Germany; Michael Flasshove, Krankenhaus Dören gGmbH, Dören, Germany; Andreas Mackensen, Universitätsklinikum Erlangen-Nürnberg, Erlangen, Germany; Maher Hanoun, Universitätsklinikum Essen, Essen, Germany; Michael Kiehl, Klinikum Frankfurt (Oder) GmbH, Frankfurt (Oder), Germany; Hubert Serve, Klinikum der J. W. Goethe Universität, Frankfurt (Main), Germany; H.-G. Höffkes, Klinikum Fulda, Fulda, Germany;

Lutz Peter Müller, Universitätsklinikum Halle (Saale), Halle (Saale), Germany; Heinz Albert Dürk, St Barbara-Klinik Hamm, Hamm, Germany; Carsten Müller-Tidow, Universitätsklinikum Heidelberg, Heidelberg, Germany; Ulrich Kaiser, St Bernward Krankenhaus, Hildesheim, Germany; Andreas Hochhaus, Universitätsklinikum Jena, Jena, Germany; Gerhard Held, Westpfalz-Klinikum GmbH, Kaiserslautern, Germany; Roland Peter Repp, Städt Krankenhaus Kiel, Kiel, Germany; Claudia Baldus, Universitätsklinikum Schleswig-Holstein, Kiel, Germany; Jens-Marcus Chemnitz, Gemeinschaftsklinikum Mittelrhein gGmbH, Koblenz, Germany; Uwe Platzbecker, Universitätsklinikum Leipzig, Leipzig, Germany; Stefan Klein, Klinikum Mannheim gGmbH, Mannheim, Germany; Andreas Neubauer, Universitätsklinikum Marburg GmbH, Marburg, Germany; Wolfgang E. Berdel, Universitätsklinikum Münster, Münster, Germany; Martin Wilhelm, Klinikum Nürnberg Nord, Nürnberg, Germany; Jörg Schubert, Elblandklinikum Riesa, Riesa, Germany; Achim Meinhardt, Agaplesion Diakonieklinikum Rotenburg gGmbH, Rotenburg (Wümme), Germany; Thomas Geer, Diakonie-Krankenhaus, Schwäbisch-Hall, Germany; Markus Ritter, Klinikum Sindelfingen-Böblingen, Sindelfingen, Germany; Walter Erich Aulitzky, Robert-Bosch-Krankenhaus, Stuttgart, Germany; Natalia Heinz, HSK Wiesbaden, Wiesbaden, Germany; Markus Schaich, Rems-Murr-Klinikum Winnenden, Winnenden, Germany; and Hermann Einsele, Universitätsklinikum Würzburg, Würzburg, Germany.

REFERENCES

- Döhner H, Weisdorf DJ, Bloomfield CD. Acute Myeloid Leukemia. *N Engl J Med*. 2015;373(12):1136-1152.
- Nakao M, Yokota S, Iwai T, et al. Internal tandem duplication of the flt3 gene found in acute myeloid leukemia. *Leukemia*. 1996; 10(12):1911-1918.
- Kottaridis PD, Gale RE, Frew ME, et al. The presence of a FLT3 internal tandem duplication in patients with acute myeloid leukemia (AML) adds important prognostic information to cytogenetic risk group and response to the first cycle of chemotherapy: analysis of 854 patients from the United Kingdom Medical Research Council AML 10 and 12 trials. *Blood*. 2001;98(6):1752-1759.
- Meshinchi S, Woods WG, Stirewalt DL, et al. Prevalence and prognostic significance of Flt3 internal tandem duplication in pediatric acute myeloid leukemia. *Blood*. 2001;97(1): 89-94.
- Thiede C, Steudel C, Mohr B, et al. Analysis of FLT3-activating mutations in 979 patients with acute myelogenous leukemia: association with FAB subtypes and identification of subgroups with poor prognosis. *Blood*. 2002;99(12):4326-4335.
- Falini B, Mecucci C, Tiacci E, et al; GIMEMA Acute Leukemia Working Party. Cytoplasmic nucleophosmin in acute myelogenous leukemia with a normal karyotype. *N Engl J Med*. 2005;352(3):254-266.
- Verhaak RG, Goudswaard CS, van Putten W, et al. Mutations in nucleophosmin (NPM1) in acute myeloid leukemia (AML): association with other gene abnormalities and previously established gene expression signatures and their favorable prognostic significance. *Blood*. 2005;106(12):3747-3754.
- Thiede C, Koch S, Creutzig E, et al. Prevalence and prognostic impact of NPM1 mutations in 1485 adult patients with acute myeloid leukemia (AML). *Blood*. 2006; 107(10):4011-4020.
- Pabst T, Mueller BU, Zhang P, et al. Dominant-negative mutations of CEBPA, encoding CCAAT/enhancer binding protein-alpha (C/EBPalpha), in acute myeloid leukemia. *Nat Genet*. 2001;27(3):263-270.
- Gombart AF, Hofmann WK, Kawano S, et al. Mutations in the gene encoding the transcription factor CCAAT/enhancer binding protein alpha in myelodysplastic syndromes and acute myeloid leukemias. *Blood*. 2002;99(4):1332-1340.
- Preudhomme C, Sagot C, Boissel N, et al; ALFA Group. Favorable prognostic significance of CEBPA mutations in patients with de novo acute myeloid leukemia: a study from the Acute Leukemia French Association (ALFA). *Blood*. 2002;100(8): 2717-2723.
- Fröhling S, Schlenk RF, Stolze I, et al. CEBPA mutations in younger adults with acute myeloid leukemia and normal cytogenetics: prognostic relevance and analysis of cooperating mutations. *J Clin Oncol*. 2004; 22(4):624-633.
- Zhang DE, Zhang P, Wang ND, Hetherington CJ, Darlington GJ, Tenen DG. Absence of granulocyte colony-stimulating factor signaling and neutrophil development in CCAAT enhancer binding protein alpha-deficient mice. *Proc Natl Acad Sci USA*. 1997;94(2):569-574.
- Geletu M, Balkhi MY, Peer Zada AA, et al. Target proteins of C/EBPalpha30 in AML: C/EBPalpha30 enhances sumoylation of C/EBPalpha42 via up-regulation of Ubc9. *Blood*. 2007;110(9):3301-3309.
- van Waalwijk van Doorn-Khosrovani SB, Erpelinck C, Meijer J, et al. Biallelic mutations in the CEBPA gene and low CEBPA expression levels as prognostic markers in intermediate-risk AML. *Hematol J*. 2003;4(1):31-40.
- Snaddon J, Smith ML, Neat M, et al. Mutations of CEBPA in acute myeloid leukemia FAB types M1 and M2. *Genes Chromosomes Cancer*. 2003;37(1):72-78.
- Liang DC, Shih LY, Huang CF, et al. CEBPalpha mutations in childhood acute myeloid leukemia. *Leukemia*. 2005;19(3): 410-414.
- Lin LI, Chen CY, Lin DT, et al. Characterization of CEBPA mutations in acute myeloid leukemia: most patients with CEBPA mutations have biallelic mutations and show a distinct immunophenotype of the leukemic cells. *Clin Cancer Res*. 2005; 11(4):1372-1379.
- Wouters BJ, Löwenberg B, Erpelinck-Verschueren CA, van Putten WL, Valk PJ, Delwel R. Double CEBPA mutations, but not single CEBPA mutations, define a subgroup of acute myeloid leukemia with a distinctive gene expression profile that is uniquely associated with a favorable outcome. *Blood*. 2009;113(13):3088-3091.
- Dufour A, Schneider F, Metzeler KH, et al. Acute myeloid leukemia with biallelic CEBPA gene mutations and normal karyotype represents a distinct genetic entity associated with a favorable clinical outcome. *J Clin Oncol*. 2010;28(4):570-577.
- Green CL, Koo KK, Hills RK, Burnett AK, Linch DC, Gale RE. Prognostic significance of CEBPA mutations in a large cohort of younger adult patients with acute myeloid leukemia: impact of double CEBPA mutations and the interaction with FLT3 and NPM1 mutations. *J Clin Oncol*. 2010;28(16): 2739-2747.
- Arber DA, Orazi A, Hasserjian R, et al. The 2016 revision to the World Health Organization classification of myeloid

- neoplasms and acute leukemia. *Blood*. 2016; 127(20):2391-2405.
23. Döhner H, Estey E, Grimwade D, et al. Diagnosis and management of AML in adults: 2017 ELN recommendations from an international expert panel. *Blood*. 2017; 129(4):424-447.
 24. Fasan A, Haferlach C, Alpermann T, et al. The role of different genetic subtypes of CEBPA mutated AML. *Leukemia*. 2014;28(4): 794-803.
 25. El-Sharkawi D, Sproul D, Allen CG, et al. Variable outcome and methylation status according to CEBPA mutant type in double-mutated acute myeloid leukemia patients and the possible implications for treatment. *Haematologica*. 2018;103(1):91-100.
 26. Wouters BJ, Louwers I, Valk PJ, Löwenberg B, Delwel R. A recurrent in-frame insertion in a CEBPA transactivation domain is a polymorphism rather than a mutation that does not affect gene expression profiling-based clustering of AML. *Blood*. 2007;109(1): 389-390.
 27. Benthous T, Schneider F, Mellert G, et al. Rapid and sensitive screening for CEBPA mutations in acute myeloid leukaemia. *Br J Haematol*. 2008;143(2):230-239.
 28. Carnicer MJ, Lasa A, Buschbeck M, et al. K313dup is a recurrent CEBPA mutation in de novo acute myeloid leukemia (AML). *Ann Hematol*. 2008;87(10):819-827.
 29. Fuchs O, Provaznikova D, Kocova M, et al. CEBPA polymorphisms and mutations in patients with acute myeloid leukemia, myelodysplastic syndrome, multiple myeloma and non-Hodgkin's lymphoma. *Blood Cells Mol Dis*. 2008;40(3):401-405.
 30. Schaich M, Röllig C, Soucek S, et al. Cytarabine dose of 36 g/m² compared with 12 g/m² within first consolidation in acute myeloid leukemia: results of patients enrolled onto the prospective randomized AML96 study. *J Clin Oncol*. 2011;29(19): 2696-2702.
 31. Röllig C, Serve H, Hüttmann A, et al; Study Alliance Leukaemia. Addition of sorafenib versus placebo to standard therapy in patients aged 60 years or younger with newly diagnosed acute myeloid leukaemia (SORAML): a multicentre, phase 2, randomised controlled trial. *Lancet Oncol*. 2015;16(16):1691-1699.
 32. Schetelig J, Schaich M, Schäfer-Eckart K, et al; Study Alliance Leukemia. Hematopoietic cell transplantation in patients with intermediate and high-risk AML: results from the randomized Study Alliance Leukemia (SAL) AML 2003 trial. *Leukemia*. 2015;29(5): 1060-1068.
 33. Röllig C, Kramer M, Gabrecht M, et al; Study Alliance Leukemia (SAL). Intermediate-dose cytarabine plus mitoxantrone versus standard-dose cytarabine plus daunorubicin for acute myeloid leukemia in elderly patients. *Ann Oncol*. 2018;29(4):973-978.
 34. Stasik S, Middeke JM, Kramer M, et al; Study Alliance Leukemia (SAL). EZH2 mutations and impact on clinical outcome: an analysis in 1,604 patients with newly diagnosed acute myeloid leukemia. *Haematologica*. 2020;105(5):e228-e231.
 35. Marcucci G, Maharry K, Radmacher MD, et al. Prognostic significance of, and gene and microRNA expression signatures associated with, CEBPA mutations in cytogenetically normal acute myeloid leukemia with high-risk molecular features: a Cancer and Leukemia Group B Study. *J Clin Oncol*. 2008; 26(31):5078-5087.
 36. Pabst T, Eyholzer M, Haefliger S, Schardt J, Mueller BU. Somatic CEBPA mutations are a frequent second event in families with germline CEBPA mutations and familial acute myeloid leukemia. *J Clin Oncol*. 2008; 26(31):5088-5093.
 37. Schlenk RF, Döhner K, Krauter J, et al; German-Austrian Acute Myeloid Leukemia Study Group. Mutations and treatment outcome in cytogenetically normal acute myeloid leukemia. *N Engl J Med*. 2008; 358(18):1909-1918.
 38. Pabst T, Eyholzer M, Fos J, Mueller BU. Heterogeneity within AML with CEBPA mutations; only CEBPA double mutations, but not single CEBPA mutations are associated with favourable prognosis. *Br J Cancer*. 2009;100(8):1343-1346.
 39. Fos J, Pabst T, Petkovic V, Ratschiller D, Mueller BU. Deficient CEBPA DNA binding function in normal karyotype AML patients is associated with favorable prognosis. *Blood*. 2011;117(18):4881-4884.
 40. Shen Y, Zhu YM, Fan X, et al. Gene mutation patterns and their prognostic impact in a cohort of 1185 patients with acute myeloid leukemia. *Blood*. 2011; 118(20):5593-5603.
 41. Taskesen E, Bullinger L, Corbacioglu A, et al. Prognostic impact, concurrent genetic mutations, and gene expression features of AML with CEBPA mutations in a cohort of 1182 cytogenetically normal AML patients: further evidence for CEBPA double mutant AML as a distinctive disease entity. *Blood*. 2011;117(8):2469-2475.
 42. Schlenk RF, Taskesen E, van Norden Y, et al. The value of allogeneic and autologous hematopoietic stem cell transplantation in prognostically favorable acute myeloid leukemia with double mutant CEBPA. *Blood*. 2013;122(9):1576-1582.
 43. Tarlock K, Lambie A, Wang J, et al. CEBPA bZip mutations are associated with favorable prognosis in de novo AML: a report from the Children's Oncology Group [published online ahead of print 21 May 2021]. *Blood*.
 44. Fasan A, Eder C, Haferlach C, et al. GATA2 mutations are frequent in intermediate-risk karyotype AML with biallelic CEBPA mutations and are associated with favorable prognosis. *Leukemia*. 2013;27(2): 482-485.
 45. Grossmann V, Haferlach C, Nadarajah N, et al. CEBPA double-mutated acute myeloid leukaemia harbours concomitant molecular mutations in 76.8% of cases with TET2 and GATA2 alterations impacting prognosis. *Br J Haematol*. 2013;161(5):649-658.
 46. Theis F, Corbacioglu A, Gaidzik VI, et al. Clinical impact of GATA2 mutations in acute myeloid leukemia patients harboring CEBPA mutations: a study of the AML study group. *Leukemia*. 2016;30(11):2248-2250.
 47. Pulikkan JA, Tenen DG, Behre G. C/EBPα deregulation as a paradigm for leukemogenesis. *Leukemia*. 2017;31(11): 2279-2285.
 48. Porse BT, Bryder D, Theilgaard-Mönch K, et al. Loss of C/EBP alpha cell cycle control increases myeloid progenitor proliferation and transforms the neutrophil granulocyte lineage. *J Exp Med*. 2005; 202(1):85-96.
 49. Bereshchenko O, Mancini E, Moore S, et al. Hematopoietic stem cell expansion precedes the generation of committed myeloid leukemia-initiating cells in C/EBPα mutant AML. *Cancer Cell*. 2009;16(5):390-400.
 50. Valk PJ, Verhaak RG, Beijnen MA, et al. Prognostically useful gene-expression profiles in acute myeloid leukemia. *N Engl J Med*. 2004;350(16):1617-1628.
 51. Pulikkan JA, Peramangalam PS, Dengler V, et al. C/EBPα regulated microRNA-34a targets E2F3 during granulopoiesis and is down-regulated in AML with CEBPA mutations. *Blood*. 2010;116(25): 5638-5649.
 52. Stavast CJ, Leenen PJM, Erkeland SJ. The interplay between critical transcription factors and microRNAs in the control of normal and malignant myelopoiesis. *Cancer Lett*. 2018;427:28-37.
 53. Wurm AA, Zjablovskaja P, Kardosova M, et al. Disruption of the C/EBPα-miR-182 balance impairs granulocytic differentiation. *Nat Commun*. 2017;8(1):46.
 54. Fontanari Krause LM, Japp AS, Krause A, et al. Identification and characterization of OSTL (RNF217) encoding a RING-IBR-RING protein adjacent to a translocation breakpoint involving ETV6 in childhood ALL. *Sci Rep*. 2014;4(1):6565.
 55. Zhang CJ, Xiang L, Chen XJ, et al. Ablation of mature miR-183 leads to retinal dysfunction in mice. *Invest Ophthalmol Vis Sci*. 2020;61(3):12.