

Introduction to Markov Chain Monte Carlo

Chris Men

chrismen4@gmail.com

October 14, 2017

Outline

Motivation

Examples

Summary

Why Markov Chain Monte Carlo (MCMC)?

- ▶ Last time we studied Gibbs sampler, which is a special case of MCMC.

Why Markov Chain Monte Carlo (MCMC)?

- ▶ Last time we studied Gibbs sampler, which is a special case of MCMC.
- ▶ Today, we will study MCMC and show why and how it works.

Why Markov Chain Monte Carlo (MCMC)?

- ▶ Last time we studied Gibbs sampler, which is a special case of MCMC.
- ▶ Today, we will study MCMC and show why and how it works.
- ▶ MCMC is very powerful in fitting very complicated models such as overly parameterized models.

Why Markov Chain Monte Carlo (MCMC)?

- ▶ Last time we studied Gibbs sampler, which is a special case of MCMC.
- ▶ Today, we will study MCMC and show why and how it works.
- ▶ MCMC is very powerful in fitting very complicated models such as overly parameterized models.
- ▶ When usual MLE and Newton-Raphson methods do not work properly, MCMC will be the best choice.

Introduction to Markov Process

- ▶ Let X_t denote the value of a random variable at time t .

Introduction to Markov Process

- ▶ Let X_t denote the value of a random variable at time t .
- ▶ Let the state space S refer to the range of the possible X values.

Introduction to Markov Process

- ▶ Let X_t denote the value of a random variable at time t .
- ▶ Let the state space S refer to the range of the possible X values.
- ▶ The random variable X is a Markov process if the transition probabilities between difference values in the state space depend only on the random variable's current state.

$$Pr(X_{t+1} = s_j | X_t = s_i, \dots, X_0 = s_k) = Pr(X_{t+1} = s_j | X_t = s_i)$$

Introduction to Markov Process

- ▶ Let X_t denote the value of a random variable at time t .
- ▶ Let the state space S refer to the range of the possible X values.
- ▶ The random variable X is a Markov process if the transition probabilities between difference values in the state space depend only on the random variable's current state.

$$Pr(X_{t+1} = s_j | X_t = s_i, \dots, X_0 = s_k) = Pr(X_{t+1} = s_j | X_t = s_i)$$

- ▶ A Markov Chain refers to a sequence of random variables (X_0, \dots, X_n, \dots) generated by a Markov process.

Introduction to Markov Chain

- ▶ A particular chain is defined most critically by its transition probabilities (or transition kernel), $P(i, j) = P(i \rightarrow j)$, which is the single step move of the process from state i to state j ,

$$P(i, j) = p(i \rightarrow j) = P(X_{t+1} = s_j | X_t = s_i)$$

Introduction to Markov Chain

- ▶ A particular chain is defined most critically by its transition probabilities (or transition kernel), $P(i, j) = P(i \rightarrow j)$, which is the single step move of the process from state i to state j ,

$$P(i, j) = p(i \rightarrow j) = P(X_{t+1} = s_j | X_t = s_i)$$

- ▶ Let $\pi_j(t) = P(X_t = j)$ denote the probability that the chain is in state j at time t , and $\pi(t)$ denote the row vector of the state space probabilities at step t .

Introduction to Markov Chain

- ▶ A particular chain is defined most critically by its transition probabilities (or transition kernel), $P(i, j) = P(i \rightarrow j)$, which is the single step move of the process from state i to state j ,

$$P(i, j) = p(i \rightarrow j) = P(X_{t+1} = s_j | X_t = s_i)$$

- ▶ Let $\pi_j(t) = P(X_t = j)$ denote the probability that the chain is in state j at time t , and $\pi(t)$ denote the row vector of the state space probabilities at step t .
- ▶ We start the chain by specifying a starting vector $\pi(0)$. Often all the elements of $\pi(0)$ are zero except for a single element of 1.

Introduction to Markov Chain

- ▶ A particular chain is defined most critically by its transition probabilities (or transition kernel), $P(i, j) = P(i \rightarrow j)$, which is the single step move of the process from state i to state j ,

$$P(i, j) = p(i \rightarrow j) = P(X_{t+1} = s_j | X_t = s_i)$$

- ▶ Let $\pi_j(t) = P(X_t = j)$ denote the probability that the chain is in state j at time t , and $\pi(t)$ denote the row vector of the state space probabilities at step t .
- ▶ We start the chain by specifying a starting vector $\pi(0)$. Often all the elements of $\pi(0)$ are zero except for a single element of 1.
- ▶ As the chain progresses, the probability values get spread out over the possible states space.

Introduction to Markov Chains (Cont'd)

- ▶ The probability that the chain has state value s_i at time or (step) $t + 1$ is given by the **Chapman – Kolomogrov** equation,

$$\begin{aligned}\pi_i(t + 1) &= P(X_{t+1} = s_i) \\ &= \sum_k P(x_{t+1} = s_i | X_t = s_k) P(X_t = s_k) \\ &= \sum_k P(k \rightarrow i) \pi_k(t) = \sum_k P(k, i) \pi_k(t)\end{aligned}$$

Introduction to Markov Chains (Cont'd)

- ▶ The probability that the chain has state value s_i at time or (step) $t + 1$ is given by the **Chapman – Kolomogrov** equation,

$$\begin{aligned}\pi_i(t + 1) &= P(X_{t+1} = s_i) \\ &= \sum_k P(x_{t+1} = s_i | X_t = s_k) P(X_t = s_k) \\ &= \sum_k P(k \rightarrow i) \pi_k(t) = \sum_k P(k, i) \pi_k(t)\end{aligned}$$

- ▶ Successive iteration of the **Chapman – Kolomogrov** describes the evolution of the chain.

Introduction to Markov Chains (Cont'd)

- ▶ The probability that the chain has state value s_i at time or (step) $t + 1$ is given by the **Chapman – Kolomogrov** equation,

$$\begin{aligned}\pi_i(t + 1) &= P(X_{t+1} = s_i) \\ &= \sum_k P(x_{t+1} = s_i | X_t = s_k) P(X_t = s_k) \\ &= \sum_k P(k \rightarrow i) \pi_k(t) = \sum_k P(k, i) \pi_k(t)\end{aligned}$$

- ▶ Successive iteration of the **Chapman – Kolomogrov** describes the evolution of the chain.
- ▶ Define the probability transition matrix **P** as the matrix whose (i, j) -th element is $P(i, j)$, the probability of moving from state i to state j , $P(i \rightarrow j)$. Which implies that

$$\sum_j P(i, j) = \sum_j P(i \rightarrow j) = 1$$

Introduction to Markov Chains (Cont'd)

- ▶ The **Chapman – Kolomogrov** equation becomes

$$\pi(t + 1) = \pi(t)\mathbf{P}$$

Introduction to Markov Chains (Cont'd)

- ▶ The **Chapman – Kolomogrov** equation becomes

$$\pi(t + 1) = \pi(t)\mathbf{P}$$

- ▶ Using the matrix form, we get

$$\pi(t) = \pi(t - 1)\mathbf{P} = (\pi(t - 2)\mathbf{P})\mathbf{P} = \pi(t - 2)\mathbf{P}^2 = \dots = \pi(0)\mathbf{P}^t$$

Introduction to Markov Chains (Cont'd)

- ▶ The **Chapman – Kolomogrov** equation becomes

$$\pi(t+1) = \pi(t)\mathbf{P}$$

- ▶ Using the matrix form, we get

$$\pi(t) = \pi(t-1)\mathbf{P} = (\pi(t-2)\mathbf{P})\mathbf{P} = \pi(t-2)\mathbf{P}^2 = \dots = \pi(0)\mathbf{P}^t$$

- ▶ Defining the n -step transition probability $P_{i,j}^{(n)}$,

$$p_{i,j}^{(n)} = P(X_{t+n} = s_j | X_t = s_i)$$

then $p_{i,j}^{(n)}$ is just the (i,j) -th element of \mathbf{P}^n .

Introduction to Markov Chains (Cont'd)

- ▶ A Markov Chain is said to be **irreducible** if there exists a positive integer such that $p_{i,j}^{(n_{ij})} > 0$ for all i, j .

Introduction to Markov Chains (Cont'd)

- ▶ A Markov Chain is said to be **irreducible** if there exists a positive integer such that $p_{i,j}^{(n_{ij})} > 0$ for all i, j .
- ▶ All states can communicate each other, as one can always go from any state to any other states.

Introduction to Markov Chains (Cont'd)

- ▶ A Markov Chain is said to be **irreducible** if there exists a positive integer such that $p_{ij}^{(n_{ij})} > 0$ for all i, j .
- ▶ All states can communicate each other, as one can always go from any state to any other states.
- ▶ A chain is said to be **aperiodic** when the number of steps required to move between two states (say x and y) is not required to be multiple of some integer.

Introduction to Markov Chains (Cont'd)

- ▶ A Markov Chain is said to be **irreducible** if there exists a positive integer such that $p_{ij}^{(n_{ij})} > 0$ for all i, j .
- ▶ All states can communicate each other, as one can always go from any state to any other states.
- ▶ A chain is said to be **aperiodic** when the number of steps required to move between two states (say x and y) is not required to be multiple of some integer.
- ▶ The conditions for a stationary distribution is that the chain is **irreducible** and **aperiodic**.

Introduction to Markov Chains (Cont'd)

- ▶ A chain may reach a stationary distribution π^* , where the vector of probabilities of being in any particular given state is independent of the initial condition.

$$\pi^* = \pi^* \mathbf{P}$$

Introduction to Markov Chains (Cont'd)

- ▶ A chain may reach a stationary distribution π^* , where the vector of probabilities of being in any particular given state is independent of the initial condition.

$$\pi^* = \pi^* \mathbf{P}$$

- ▶ π^* is the left eigenvector associated with the eigenvalue $\lambda = 1$ of \mathbf{P} .

Introduction to Markov Chains (Cont'd)

- ▶ A sufficient condition for a unique stationary distribution is that the **detailed balance** equation holds (for all i and j),

$$P(j \rightarrow k)\pi_j^* = P(k \rightarrow j)\pi_k^*$$

Introduction to Markov Chains (Cont'd)

- ▶ A sufficient condition for a unique stationary distribution is that the **detailed balance** equation holds (for all i and j),

$$P(j \rightarrow k)\pi_j^* = P(k \rightarrow j)\pi_k^*$$

- ▶ The Markov Chain is said to be **reversible**, and hence the above equation is called the **reversibility condition**.

Introduction to Markov Chains (Cont'd)

- ▶ A sufficient condition for a unique stationary distribution is that the **detailed balance** equation holds (for all i and j),

$$P(j \rightarrow k)\pi_j^* = P(k \rightarrow j)\pi_k^*$$

- ▶ The Markov Chain is said to be **reversible**, and hence the above equation is called the **reversibility condition**.
- ▶ This condition implies $\pi = \pi\mathbf{P}$, as the j -th element of $\pi\mathbf{P}$ is

$$(\pi\mathbf{P})_j = \sum_i \pi_i P(i \rightarrow j) = \sum_i \pi_j P(j \rightarrow i) = \pi_j \sum_i P(j \rightarrow i) = \pi_j$$

Introduction to Markov Chains (Cont'd)

- ▶ A sufficient condition for a unique stationary distribution is that the **detailed balance** equation holds (for all i and j),

$$P(j \rightarrow k)\pi_j^* = P(k \rightarrow j)\pi_k^*$$

- ▶ The Markov Chain is said to be **reversible**, and hence the above equation is called the **reversibility condition**.
- ▶ This condition implies $\pi = \pi \mathbf{P}$, as the j -th element of $\pi \mathbf{P}$ is

$$(\pi \mathbf{P})_j = \sum_i \pi_i P(i \rightarrow j) = \sum_i \pi_j P(j \rightarrow i) = \pi_j \sum_i P(j \rightarrow i) = \pi_j$$

- ▶ The discrete-state Markov chain can be generalized to a continuous state Markov process by having a probability kernel $P(x, y)$ that satisfies

$$\int P(x, y) dy = 1$$

Introduction to Markov Chains (Cont'd)

- ▶ And the continuous extension of the **Chapman – Kologronvo** equation becomes,

$$\pi_t(y) = \int \pi_{t-1}(x)P(x, y)dx$$

Introduction to Markov Chains (Cont'd)

- ▶ And the continuous extension of the **Chapman – Kologronvo** equation becomes,

$$\pi_t(y) = \int \pi_{t-1}(x)P(x, y)dx$$

- ▶ At equilibrium, that the stationary distribution satisfies,

$$\pi^*(y) = \int \pi^*(x)P(x, y)dx$$

The Metropolis-Hastings (MH) Algorithm

- ▶ The goal of Monte Carlo is to obtain samples from some complex probability distribution $p(\theta)$, θ may be a vector of variables.

The Metropolis-Hastings (MH) Algorithm

- ▶ The goal of Monte Carlo is to obtain samples from some complex probability distribution $p(\theta)$, θ may be a vector of variables.
- ▶ Suppose that we want to draw samples from some distribution $p(\theta)$, where $p(\theta) = f(\theta)/k$, where k may not be known or is very difficult to obtain.

The Metropolis-Hastings (MH) Algorithm

- ▶ The goal of Monte Carlo is to obtain samples from some complex probability distribution $p(\theta)$, θ may be a vector of variables.
- ▶ Suppose that we want to draw samples from some distribution $p(\theta)$, where $p(\theta) = f(\theta)/k$, where k may not be known or is very difficult to obtain.
- ▶ This is usually the case in Bayesian statistics, for example, $f(\theta) = \prod_{i=1}^n f_i(\theta), f_i(\theta) > 0$.

The Metropolis-Hastings (MH) Algorithm

- ▶ The goal of Monte Carlo is to obtain samples from some complex probability distribution $p(\theta)$, θ may be a vector of variables.
- ▶ Suppose that we want to draw samples from some distribution $p(\theta)$, where $p(\theta) = f(\theta)/k$, where k may not be known or is very difficult to obtain.
- ▶ This is usually the case in Bayesian statistics, for example, $f(\theta) = \prod_{i=1}^n f_i(\theta), f_i(\theta) > 0$.
- ▶ In my next talk, we will study the slice sampler proposed by Neal (2013) to simulate those types of posterior distributions.

The Metropolis Algorithm

- 1 Start with any initial value $\theta^{(0)}$ satisfying $f(\theta^{(0)}) > 0$. Set $k = 0$.
- 2 Using the current values $\theta^{(k)}$, generate a value θ^* from some jumping (candidate, or proposal) distribution, $q(\theta^{(k)}, \theta^*)$, which is the probability of returning a values of θ^* given a previous value of $\theta^{(k)}$. Here the jump distribution is symmetric such that $q(\theta^{(k)}, \theta^*) = q(\theta^*, \theta^{(k)})$.

- 3 Calculate the ratio of the density at the candidate θ^* and current $\theta^{(k)}$ points,

$$\alpha = \frac{p(\theta^*)}{p(\theta^{(k-1)})} = \frac{f(\theta^*)}{f(\theta^{(k-1)})}$$

- 4 If $\alpha \geq 1$ then set $\theta^{(k+1)} = \theta^*$ and go to step 2. If $\alpha < 1$ then with probability α accept the candidate point, otherwise reject θ^* and set $\theta^{(k+1)} = \theta^{(k)}$ then go to step 2.
- 4' Generate a uniform distribution $u = U(0, 1)$. if $u < \min(\alpha, 1)$ then set $\theta^{(k+1)} = \theta^*$; Otherwise set $\theta^{(k+1)} = \theta^{(k)}$ and go step 2.

The Metropolis-Hastings Algorithm (Cont'd)

- ▶ Hastings (1970) generalized the Metropolis algorithm by using an arbitrary transition probability function $q(\theta_1, \theta_2) = \Pr(\theta_1 \rightarrow \theta_2)$, and setting the acceptance probability for a candidate point as

$$\alpha = \frac{f(\theta^*)q(\theta^*|\theta^{(k-1)})}{f(\theta^{(k-1)})q(\theta^{(k-1)}|\theta^*)}$$

- ▶ Chib and Greenberg (1995) said that finding a jumping function appears to be a search for the proverbial needle in a haystack.

Example 1 – fit a Student $-t$ distribution

- ▶ Suppose that we have a data set $\mathbf{y} = \{y_i, i = 1, \dots, n\}$, which is generated from a Student- t distribution $y \sim t(\nu)$ with pdf

$$f(y) = \frac{\nu^{\nu/2} \Gamma((\nu+1)/2)}{\Gamma(\nu/2) \Gamma(1/2)} (\nu + y^2)^{-(\nu+1)/2}.$$

Example 1 – fit a Student - t distribution

- ▶ Suppose that we have a data set $\mathbf{y} = \{y_i, i = 1, \dots, n\}$, which is generated from a Student- t distribution $y \sim t(\nu)$ with pdf

$$f(y) = \frac{\nu^{1/2} \Gamma((\nu+1)/2)}{\Gamma(\nu/2) \Gamma(1/2)} (\nu + y^2)^{-(\nu+1)/2}.$$

- ▶ The likelihood of the data based on the model is

$$\begin{aligned} f(\mathbf{y}|\nu) &= \prod_{i=1}^n f(y_i|\nu) \\ &\propto \prod_{i=1}^n \frac{\nu^{1/2} \Gamma((\nu+1)/2)}{\Gamma(\nu/2)} (\nu + y_i^2)^{-(\nu+1)/2} \end{aligned}$$

Example 1 – fit a Student - t distribution

- ▶ Suppose that we have a data set $\mathbf{y} = \{y_i, i = 1, \dots, n\}$, which is generated from a Student- t distribution $y \sim t(v)$ with pdf

$$f(y) = \frac{v^{v/2} \Gamma((v+1)/2)}{\Gamma(v/2) \Gamma(1/2)} (v + y^2)^{-(v+1)/2}.$$

- ▶ The likelihood of the data based on the model is

$$\begin{aligned} f(\mathbf{y}|v) &= \prod_{i=1}^n f(y_i|v) \\ &\propto \prod_{i=1}^n \frac{v^{v/2} \Gamma((v+1)/2)}{\Gamma(v/2)} (v + y_i^2)^{-(v+1)/2} \end{aligned}$$

- ▶ The posterior distribution of v is

$$f(v|\mathbf{y}) \propto f(v) \prod_{i=1}^n \frac{v^{v/2} \Gamma((v+1)/2)}{\Gamma(v/2)} (v + y_i^2)^{-(v+1)/2}$$

where $f(v)$ is the prior distribution of v .

Example 1 – fit a Student $-t$ distribution (Cont'd)

- ▶ Since this full conditional is an unknown distribution, we use a random-walk Metropolis-Hastings algorithm, in which the proposal density is a standard Gaussian density.




Figure: The training data was drawn from $t(10)$ with sample size = 1000. The estimate of v is $v = 12.57$ with standard deviation $sd = 2.73$.

- ▶ The MH algorithm was iterated 10,000 times. After the first 5000 iterations were discarded as burn in, parameter was estimated by sample mean.

Example 2 – fit a logistic regression model

- ▶ Let y be a Bernoulli random variable taking value either 0 or 1.

Example 2 – fit a logistic regression model

- ▶ Let y be a Bernoulli random variable taking value either 0 or 1.
- ▶ Define $X \in \mathbb{R}^m$ be the associated vector of covariates.

Example 2 – fit a logistic regression model

- ▶ Let y be a Bernoulli random variable taking value either 0 or 1.
- ▶ Define $X \in \mathbb{R}^m$ be the associated vector of covariates.
- ▶ Supposed that we obtain a sample $\{y_i, X_i\}_{i=1}^n$, where n is the sample size.

Example 2 – fit a logistic regression model

- ▶ Let y be a Bernoulli random variable taking value either 0 or 1.
- ▶ Define $X \in \mathbb{R}^m$ be the associated vector of covariates.
- ▶ Supposed that we obtain a sample $\{y_i, X_i\}_{i=1}^n$, where n is the sample size.
- ▶ The relationship between y and X can be modeled by a generalized linear model (GLM),

$$f(y_i|X_i, \theta) = \beta_0 + X_i^T \beta + \epsilon_i, \quad i = 1, \dots, n$$

where ϵ_i are *iid* white noises with mean 0 and variance σ^2 .

Logistic regression (LR) model (Cont'd)

- ▶ In logistic regression, we use a logistic function $p(\cdot)$, defined below,

$$p(y|X, \theta) = \frac{e^{\beta_0 + X^T \beta}}{1 + e^{\beta_0 + X^T \beta}}.$$

Logistic regression (LR) model (Cont'd)

- ▶ In logistic regression, we use a logistic function $p(\cdot)$, defined below,

$$p(y|X, \theta) = \frac{e^{\beta_0 + X^T \beta}}{1 + e^{\beta_0 + X^T \beta}}.$$

- ▶ After a bit of manipulation, we obtain the following

$$\frac{p(y|X, \theta)}{1 - p(y|X, \theta)} = e^{\beta_0 + X^T \beta}, \theta^T = (\beta_0, \beta_1, \dots, \beta_m)$$

The quantity of $p(y|X, \theta)/(1 - p(y|X, \theta))$ is called the odds, which can take any values in $(0, \infty)$.

Logistic regression (LR) model (Cont'd)

- ▶ In logistic regression, we use a logistic function $p(\cdot)$, defined below,

$$p(y|X, \theta) = \frac{e^{\beta_0 + X^T \beta}}{1 + e^{\beta_0 + X^T \beta}}.$$

- ▶ After a bit of manipulation, we obtain the following

$$\frac{p(y|X, \theta)}{1 - p(y|X, \theta)} = e^{\beta_0 + X^T \beta}, \theta^T = (\beta_0, \beta_1, \dots, \beta_m)$$

The quantity of $p(y|X, \theta)/(1 - p(y|X, \theta))$ is called the odds, which can take any values in $(0, \infty)$.

- ▶ The usual logistic regression model is,

$$\log \left(\frac{p(y|X, \theta)}{1 - p(y|X, \theta)} \right) = \beta_0 + X^T \beta.$$

Markov Chain Monte Carlo (MCMC) estimation

- The likelihood function of the model is

$$\begin{aligned} L(y|X, \theta) &= \prod_{i=1}^n p(X_i; \theta)^{y_i} (1 - p(X_i; \theta))^{1-y_i} \\ &= \prod_{i=1}^n \left(\frac{e^{\beta_0 + X_i^T \beta}}{1 + e^{\beta_0 + X_i^T \beta}} \right)^{y_i} \left(1 - \frac{e^{\beta_0 + X_i^T \beta}}{1 + e^{\beta_0 + X_i^T \beta}} \right)^{1-y_i} \end{aligned}$$

Note: All the parameters in the model are treated as random variables with prior distributions.

Metropolis-Hastings (MH) sampler

- 0 Let γ be a parameter in θ , and define $\theta_{-\gamma}$ as all parameters in θ except γ .





Metropolis-Hastings (MH) sampler

- 0 Let γ be a parameter in θ , and define $\theta_{-\gamma}$ as all parameters in θ except γ .
1. Initialize $\gamma^{(0)} \sim q(\cdot)$, where $q(\cdot)$ is a candidate distribution, which is a normal distribution, $N(\mu, \sigma^2)$.
2. For $k=1, \dots, K$, repeat:
 - (a) Sample $\gamma^* \sim q(\gamma | \gamma^{(k-1)})$ such that $\gamma^* \sim N(\gamma^{(k-1)}, \sigma^2)$.
 - (b) Calculate acceptance probability:

$$\begin{aligned}\alpha &= \min \left\{ 1, \frac{L(y|x, \theta_{-\gamma}, \gamma^{(*)}) q(\gamma^{(k-1)} | \gamma^{(*)})}{L(y|x, \theta_{-\gamma}, \gamma^{(k-1)}) q(\gamma^{(*)} | \gamma^{(k-1)})} \right\} \\ &= \min \left\{ 1, \frac{L(y|x, \theta_{-\gamma}, \gamma^{(*)})}{L(y|x, \theta_{-\gamma}, \gamma^{(k-1)})} \right\}, \text{ as } q(\cdot) \text{ is symmetric.}\end{aligned}$$

- (c) Sample $u \sim \text{Uniform}(0, 1)$.
- (d) If $u < \alpha$ then accept the proposal $\gamma^{(k)} = \gamma^*$; Otherwise, reject the proposal and set $\gamma^{(k)} = \gamma^{(k-1)}$.

Example 2 – fit a logistic regression model – sampled time series



- The MH algorithm was iterated 10,000 times. After the first 5000 iterations were discarded, parameters were estimated by sample means.

Example 2 – fit a logistic regression model (Cont'd)

Table: Fit a logistic regression model.

Parameter	True	Estimate	SD
β_0	1	1.10	0.12
β_1	1.5	1.62	0.13
β_2	2.2	2.45	0.17
β_3	3	3.36	0.20

Summary

- ▶ Introduced Markov process and Markov Chain.

Summary

- ▶ Introduced Markov process and Markov Chain.
- ▶ Introduced MCMC, Metropolis, and Metropolis-Hastings samplers.

Summary

- ▶ Introduced Markov process and Markov Chain.
- ▶ Introduced MCMC, Metropolis, and Metropolis-Hastings samplers.
- ▶ Two examples were given to show how MCMC works.

Summary

- ▶ Introduced Markov process and Markov Chain.
- ▶ Introduced MCMC, Metropolis, and Metropolis-Hastings samplers.
- ▶ Two examples were given to show how MCMC works.
- ▶ The usual logistic regression model can be extended to,

$$\log \left(\frac{p(y|X, \theta)}{1 - p(y|X, \theta)} \right) = \beta_0 + X^T \beta + \epsilon,$$

where ϵ is white noises with mean 0 and variance σ^2 .

Thank you!

References

- [1] Robert, C. and G., Casella. 2013. Monte Carlo Statistical Methods: Edition 2. London: Chapman and Hall.

- [2] Gilks, W. R., Richardson, S., and Spiegelhalter, D. J. 1995. Markov Chain Monte Carlo in Practice. London: Chapman and Hall.

- [3] Neal, R. N. 2003. Slice sampling. *The Annals of Statistics* 31: 705-767.

- [4] Chib, S. and Greenberg E. 1995. Understanding the Metropolis-Hastings Algorithm. *The American Statistician* Vol. 49, No. 4: 327-335.