

Mixed method for linear elastic thin-plate flexure: variational form

PROBLEM DESCRIPTION

Consider a thin plate loaded with some force $f(x, y)$. Let \mathbf{M} denote the moment tensor and $\eta = \eta(x, y)$ represent the vertical deflection in response to the load $f = f(x, y)$. With λ_1 and λ_2 some appropriately-defined scalar-valued functions in x and y , the governing equation and constitutive relation on the plate are:

$$-\nabla \cdot (\nabla \cdot \mathbf{M}) = f \quad (1a)$$

$$\mathbf{M} = \lambda_1 \nabla \nabla \eta + \lambda_2 \text{tr}(\nabla \nabla \eta) \vec{I} \quad (1b)$$

The goal is to separately put Equations 1a and 1b into variational form.

USEFUL IDENTITIES

Some useful identities which will help in attaining the variational forms for Equations 1a and 1b. In the identities below, α and β will denote scalar quantities, \vec{a} and \vec{b} are vectors, and \mathbf{A} and \mathbf{B} are tensors. Ω represents the domain of the problem, Γ is the boundary, and \hat{n} is the unit, outward-pointing vector normal to Γ .

$$\nabla \cdot (\alpha \mathbf{A}) = \alpha (\nabla \cdot \mathbf{A}) + \mathbf{A} \cdot \nabla \alpha \quad (2)$$

$$\nabla \cdot (\alpha \vec{a}) = \alpha (\nabla \cdot \vec{a}) + \vec{a} \cdot \nabla \alpha \quad (3)$$

$$\int_{\Omega} \nabla \cdot \vec{a} = \int_{\Gamma} \vec{a} \cdot \hat{n} \quad (4)$$

14 PUTTING EQUATION 1a INTO VARIATIONAL FORM

15 Multiply both sides of the governing equation by arbitrary scalar function $w = w(x, y)$. Then

$$-w \nabla \cdot (\nabla \cdot \mathbf{M}) = wf$$

16 Note that the left-hand side is the product of the scalar variable $-w$ with the divergence of the vector
17 $(\nabla \cdot \mathbf{M})$. Therefore, by the product rule of Equation 3, the equation above can be rewritten so that

$$(\nabla \cdot \mathbf{M}) \cdot \nabla w - \nabla \cdot [w(\nabla \cdot \mathbf{M})] = wf.$$

18 Integrating over the domain Ω , we have that

$$\int_{\Omega} [(\nabla \cdot \mathbf{M}) \cdot \nabla w] - \int_{\Omega} \nabla \cdot [w(\nabla \cdot \mathbf{M})] = \int_{\Omega} wf.$$

19 Finally, by the Divergence Theorem of Equation 4, it follows that

$$\int_{\Omega} [(\nabla \cdot \mathbf{M}) \cdot \nabla w] - \int_{\Gamma} [w(\nabla \cdot \mathbf{M}) \cdot \hat{n}] = \int_{\Omega} wf \quad (5)$$

20 PUTTING EQUATION 1b INTO VARIATIONAL FORM

21 Let \mathbf{N} be an arbitrary tensor function. Then the constitutive equation of Equation 1b requires that

$$\int_{\Omega} \mathbf{N} : \mathbf{M} = \int_{\Omega} \lambda_1 \mathbf{N} : \nabla \nabla \eta + \int_{\Omega} \lambda_2 \mathbf{N} : \text{tr}(\nabla \nabla \eta) \vec{I}$$

22 I'll split the right-hand side up into two pieces:

23 The first term

24 First, we'll try writing the expression

$$\int_{\Omega} \lambda_1 \mathbf{N} : \nabla \nabla \eta \quad (6)$$

25 in terms of only first derivatives...

26 **The second term**

27 Finally, put the expression

$$\int_{\Omega} \lambda_2 \mathbf{N} : tr(\nabla \nabla \eta) \vec{I} \tag{7}$$

28 in terms of only first derivatives...