

ABSTRACT

Permutation polynomials over finite fields are important in many applications, for example in cryptography. We want to provide families of polynomials that are rich in permutation polynomials. In particular we study polynomials of the form $F_{a,b}(x) = x^{\frac{q-1}{2}} + ax^{\frac{q+d-1}{d}} + bx$, where $a, b \in \mathbb{F}_q^*$, $q = p^r$, p prime, and $d \mid (q - 1)$.

PRELIMINARIES

Definition. A *permutation* of a set A is an ordering of the elements of A . A function $f : A \rightarrow A$ gives a permutation of A if and only if f is one to one and onto.

Definition. A *finite field* \mathbb{F}_q , $q = p^r$, p prime, is a field with $q = p^r$ elements.

Definition. A *primitive root* $\alpha \in \mathbb{F}_q$ is a generator for the multiplicative group \mathbb{F}_q^\times

Example 1. Consider the finite field \mathbb{F}_7 . We have that: $3^1 = 3, 3^2 = 2, 3^3 = 6, 3^4 = 4, 3^5 = 5, 3^6 = 1$, so 3 is a primitive root of \mathbb{F}_7 .

Definition. Let $f(x)$ be a polynomial defined over a finite field \mathbb{F}_q . Then the *value set* of f is defined as $V_f = \{f(a) \mid a \in \mathbb{F}_q\}$

Note that a polynomial $f(x)$ defined over \mathbb{F}_q is a permutation polynomial if and only if $V_f = \mathbb{F}_q$.

Example 2. Consider the polynomial $f(x) = x + 3$ defined over \mathbb{F}_7 . We have that $f(0) = 3, f(1) = 4, f(2) = 5, f(3) = 6, f(4) = 0, f(5) = 1, f(6) = 2$, so $f(x)$ is a permutation polynomial over \mathbb{F}_7

MOTIVATION

Binomials that produce permutations have been studied extensively. The next case to be studied are trinomials. We have found that within the family of polynomials of the form

$$F_{a,b}(x) = x^{\frac{q-1}{2}} + ax^{\frac{q+d-1}{d}} + bx,$$

where $d \mid (q - 1)$ there are many permutation polynomials. We want to find conditions in $[a, b]$ that guarantee that $F_{a,b}(X)$ is a permutation polynomial and count how many permutation polynomials exist in each family.

PROBLEM

Determine conditions in a, b such that polynomials of the form

$$F_{a,b}(x) = x^{\frac{q-1}{2}} + ax^{\frac{q+d-1}{d}} + bx,$$

where $a, b \in \mathbb{F}_q^*$, and $d \mid (q - 1)$ give permutations of \mathbb{F}_q , and determine how many pairs exist for each d .

RESULTS

The following theorem gives information on the amount of polynomials with the same value set.

APPLICATIONS

An example of applications of permutation polynomials over finite fields are RSA-type cryptosystems. In some of these systems secret messages are encoded as elements of a field \mathbb{F}_q with a sufficiently large q . The encryption operator used for these systems is a permutation of the field \mathbb{F}_q and needs to be efficiently computable. It is easy to see that expressing this operator in terms of a permutation polynomial is simple and efficient.

ONGOING WORK

Knowing a permutation polynomial we can construct d or $2d$ of them (depending on the parity of d). We still need to characterize which

REFERENCES

- Lidl, Rudolf, and Harald Niederreiter. *Finite Fields*. Reading, Mass.: Addison-Wesley Pub. Co., Advanced Book Program/World Science Division, 1983. Print.
- Wan, D., Lidl, R. *Permutation Polynomials of the Form $x^r f(x^{\frac{q-1}{d}})$ and Their Group Structure*. Mh. Math. 112, 149-163 (1991).
- Mullen, G., Stevens H. *Polynomial Functions (mod m)*. Acta Math. Hung. 44(3-4) (1984), 237-241.