

On a Class of Permutation Polynomials over Finite Fields

September 30, 2013

Abstract

Permutation polynomials over finite fields are important in many applications, for example in cryptography. We want to provide families of polynomials that are rich in permutation polynomials. In particular we study polynomials of the form $F_{a,b}(x) = x^{\frac{q+1}{2}} + ax^{\frac{q+d-1}{d}} + bx$, where $a, b \in \mathbb{F}_q$, $q = p^r$, p prime, and $d \mid (q-1)$.

1 Introduction

Permutation polynomials over finite fields are important in many applications, for example in cryptography. Binomials that produce permutations have been studied extensively. The next case to be studied are trinomials. We want to provide families of polynomials that are rich in permutation polynomials. We have found that within the family of polynomials of the form

$$F_{a,b}(x) = x^{\frac{q+1}{2}} + ax^{\frac{q+d-1}{d}} + bx,$$

where $d \mid (q-1)$ there are many permutation polynomials. We want to find conditions in $[a, b]$ that guarantee that $F_{a,b}(X)$ is a permutation polynomial and count how many permutation polynomials exist in each family.

An example of applications of permutation polynomials over finite fields are RSA-type cryptosystems. In some of these systems secret messages are encoded as elements of a field \mathbb{F}_q with a sufficiently large q . The encryption operator used for these systems is a permutation of the field \mathbb{F}_q and needs to be efficiently computable. It is easy to see that expressing this operator in terms of a permutation polynomial is simple and efficient.

2 Preliminaries

Definition 2.1. A **permutation** of a set A is an ordering of the elements of A . A function $f : A \rightarrow A$ gives a permutation of A if and only if f is one to one and onto.

Definition 2.2. A **finite field** \mathbb{F}_q , $q = p^r$, p prime, is a field with $q = p^r$ elements.

Definition 2.3. A **primitive root** $\alpha \in \mathbb{F}_q$ is a generator for the multiplicative group \mathbb{F}_q^\times

Example 2.4. Consider the finite field \mathbb{F}_7 . We have that: $3^1 = 3, 3^2 = 2, 3^3 = 6, 3^4 = 4, 3^5 = 5, 3^6 = 1$, so 3 is a primitive root of \mathbb{F}_7 .

Definition 2.5. Let $f(x)$ be a polynomial defined over a finite field \mathbb{F}_q . Then the **value set** of f is defined as $V_f = \{f(a) \mid a \in \mathbb{F}_q\}$

Note that a polynomial $f(x)$ defined over \mathbb{F}_q is a permutation polynomial if and only if $V_f = \mathbb{F}_q$.

Example 2.6. Consider the polynomial $f(x) = x + 3$ defined over \mathbb{F}_7 . We have that $f(0) = 3, f(1) = 4, f(2) = 5, f(3) = 6, f(4) = 0, f(5) = 1, f(6) = 2$, so $f(x)$ is a permutation polynomial over \mathbb{F}_7

3 Results

3.1 Counting permutation polynomials

The following theorem gives information on the amount of polynomials with the same value set.

Theorem 3.1. Fix $n \in \mathbb{N}$, $n \leq q$. The number of polynomials of the form $F_{a,b}(x)$ with $|V_{F_{a,b}}| = n$ is a multiple of d if d is even, or a multiple of $2d$ if d is odd.

Corollary 3.2. The number of permutation polynomials over \mathbb{F}_q of the form $F_{a,b}(x)$ is a multiple of d if d is even, or a multiple of $2d$ if d is odd.

Given coefficients $[a, b]$ for which $F_{a,b}(x)$ is a permutation polynomial of \mathbb{F}_q , we can construct a list of d or $2d$ coefficients $[a', b']$ such that $F_{a',b'}(x)$ is also permutation polynomial of \mathbb{F}_q as follows:

Construction 3.3. Let $d \mid (q-1)$, d odd, and $F_{a,b}(x) = x^{\frac{q+1}{2}} + ax^{\frac{q+d-1}{d}} + bx$ be a permutation polynomial of \mathbb{F}_q , where $a = \alpha^i$, $b = \alpha^j$. Then $F_{a',b'}(x)$ is also a permutation polynomial for $[a', b'] \in \left\{ \alpha^{i+k(d+2)\frac{q-1}{2d}}, \alpha^{j+k\frac{q-1}{2}} \mid k = 1, \dots, 2d-1 \right\}$

Example 3.4. Fix $d = 3$ and $q = 43$. There exists 48 pairs $[a, b]$ such that $F_{a,b}(x) = x^{22} + ax^{15} + bx$. In particular, we know that $F_{1,17}(x) = x^{22} + 1x^{15} + 17x$ is a permutation polynomial over \mathbb{F}_{43} . Using $1 = 3^{42} = \alpha^{42}$, $17 = 3^{38} = \alpha^{38}$, we obtain 5 other pairs $[a', b']$ and new permutation polynomials $F_{a',b'}(x)$ using our construction:

$$\begin{aligned}
& [7 = \alpha^{42+5 \cdot 7}, 26 = \alpha^{38+21}], [6 = \alpha^{42+2(5 \cdot 7)}, 17 = \alpha^{38+2(21)}], \\
& [42 = \alpha^{42+3(5 \cdot 7)}, 26 = \alpha^{38+3(21)}], \\
& [36 = \alpha^{42+4(5 \cdot 7)}, 17 = \alpha^{38+4(21)}], \\
& [37 = \alpha^{42+5(5 \cdot 7)}, 26 = \alpha^{38+5(21)}]
\end{aligned}$$

Knowing a permutation polynomial we can construct d or $2d$ of them (depending on the parity of d). We still need to characterize which polynomials are permutation polynomials. For this, we are studying the size of the value sets of $F_{a,b}(x)$.

3.2 Size of the value set

We divide the value set into subsets:

Definition 3.5. Let $F_{a,b}(x) = x^{\frac{q+1}{2}} + ax^{\frac{q+d-1}{d}} + bx$ be a polynomial defined over \mathbb{F}_q , where $d \mid (q-1)$. We define the sets $A_l = \{F_{a,b}(\alpha^{dk+l}) \mid k = 0, \dots, \frac{q-1}{d}\}$ for $l = 0, \dots, d-1$, where α is a primitive root of \mathbb{F}_q .

For these subsets we have proved the following lemmas

Lemma 3.6. Let $F_{a,b}(x)$ be defined over \mathbb{F}_q and A_l be defined as above. We have that $|A_l| = \frac{q-1}{d}$ or $A_l = \{0\}$

Lemma 3.7. Let $F_{a,b}(x)$ be defined over \mathbb{F}_q . The sets A_l defined above are such that, for $l \neq k$, $A_l \cap A_k = \emptyset$ or $A_l = A_k$.

Proposition 3.8. Let $F_{a,b}(x)$ be defined over \mathbb{F}_q and A_l be defined as above. $F_{a,b}(x)$ is a permutation polynomial if and only if $A_l \neq \{0\}$ and $A_l \cap A_k = \emptyset$ for $0 \leq l, k \leq d-1$.

3.3 Future work

- Find necessary and sufficient conditions on the coefficients $a = \alpha^i$, $b = \alpha^j$ such that $A_l \neq \{0\}$ and $A_l \cap A_k = \emptyset$
- Study our results on the family of polynomials of the form $F_{a,b}(x) = x^{\frac{q-1}{2}+m} + ax^{\frac{q-1}{d}+m} + bx^m$