

Construction of Families of Permutation Trinomials over Finite Fields

Christian A. Rodriguez
Alex D. Santos

Department of Computer Science
University of Puerto Rico, Río Piedras

March 14, 2014

Table of Contents

- 1 Introduction
- 2 Our Problem
- 3 Results

Table of Contents

1 Introduction

2 Our Problem

3 Results

Finite Fields

Definition

A **finite field** \mathbb{F}_q is a field with $q = p^r$ elements where p is prime.

Example

$$\mathbb{F}_5 = \{0, 1, 2, 3, 4\}$$

Addition:

$$2 + 2 = 4$$

$$4 + 4 = 8$$

$$(\text{mod } 5) = 3$$

Multiplication:

$$2 \cdot 2 = 4$$

$$4 \cdot 4 = 16$$

$$(\text{mod } 5) = 1$$

Value Sets

Definition

Let $f(x)$ be a polynomial defined over a finite field \mathbb{F}_q . Then the **value set** of f is defined as $V(f) = \{f(a) \mid a \in \mathbb{F}_q\}$

Example

Consider $f(x) = x^2$ defined over \mathbb{F}_5 .

Note: $f(0) = 0, f(1) = 1, f(2) = 4, f(3) = 4, f(4) = 1$

$V(f) = \{0, 1, 4\}$.

Permutation Polynomials

Definition

A polynomial $f(x)$ defined over \mathbb{F}_q is a **permutation polynomial** if and only if $V(f) = \mathbb{F}_q$.

Permutation Polynomials

Definition

A polynomial $f(x)$ defined over \mathbb{F}_q is a **permutation polynomial** if and only if $V(f) = \mathbb{F}_q$.

Example

Let $f(x) = x^3$ over \mathbb{F}_5 . Note: $V(f) = \{0, 1, 3, 2, 4\}$ so $f(x)$ is a permutation polynomial over \mathbb{F}_5

Permutation Polynomials

Definition

A polynomial $f(x)$ defined over \mathbb{F}_q is a **permutation polynomial** if and only if $V(f) = \mathbb{F}_q$.

Example

Let $f(x) = x^3$ over \mathbb{F}_5 . Note: $V(f) = \{0, 1, 3, 2, 4\}$ so $f(x)$ is a permutation polynomial over \mathbb{F}_5

Example

Let $f(x) = x^2$ over \mathbb{F}_5 . We have that $V(f) = \{0, 1, 4\}$ so $f(x)$ is not a permutation polynomial over \mathbb{F}_5 .

Primitive Roots

Definition

A **primitive root** $\alpha \in \mathbb{F}_q$ is a generator for the multiplicative group \mathbb{F}_q^\times

Primitive Roots

Definition

A **primitive root** $\alpha \in \mathbb{F}_q$ is a generator for the multiplicative group \mathbb{F}_q^\times

$$\mathbb{F}_5$$

Primitive Roots

Definition

A **primitive root** $\alpha \in \mathbb{F}_q$ is a generator for the multiplicative group \mathbb{F}_q^\times

$$\mathbb{F}_5$$

$$\begin{array}{rcl} 3^1 & = & 3 \\ 3^2 & = & 4 \\ 3^3 & = & 2 \\ 3^4 & = & 1 \end{array}$$

Primitive Roots

Definition

A **primitive root** $\alpha \in \mathbb{F}_q$ is a generator for the multiplicative group \mathbb{F}_q^\times

$$\mathbb{F}_5$$

$$\begin{aligned} 3^1 &= 3 \\ 3^2 &= 4 \\ 3^3 &= 2 \\ 3^4 &= 1 \end{aligned}$$

$$\begin{aligned} 4^1 &= 4 \\ 4^2 &= 1 \\ 4^3 &= 4 \\ 4^4 &= 1 \end{aligned}$$

Table of Contents

1 Introduction

2 Our Problem

3 Results

Permutation Polynomials

- Everything is known about Permutation Monomials
- Permutation Binomials have been studied extensively
- The next case is to study Permutation Trinomials

Permutation trinomials of the form $X^{\frac{q+1}{2}} + aX^{\frac{q-1}{d}+1} + bX$

$$f_{a,b} = X \left(X^{\frac{p-1}{2}} + aX^{\frac{p-1}{d}} + b \right)$$

Permutation trinomials of the form $X^{\frac{q+1}{2}} + aX^{\frac{q-1}{d}+1} + bX$

$$f_{a,b} = X \left(X^{\frac{p-1}{2}} + aX^{\frac{p-1}{d}} + b \right), N(p, d) = \text{number of permutations}$$

Permutation trinomials of the form $X^{\frac{q+1}{2}} + aX^{\frac{q-1}{d}+1} + bX$

$$f_{a,b} = X \left(X^{\frac{p-1}{2}} + aX^{\frac{p-1}{d}} + b \right), N(p, d) = \text{number of permutations}$$

p	$N(p, 3)$	$N(p, 4)$	$N(p, 6)$		p	$N(p, 3)$	$N(p, 4)$	$N(p, 6)$
13	—	8	18		61	60	304	30
17	—	16	—		67	78	—	108
19	0	—	0		73	54	440	54
29	—	48	—		79	96	—	48
31	0	—	18		89	—	680	—
37	12	132	12		97	174	840	102
41	—	140	—		101	—	940	—
43	48	—	36		103	162	—	72
53	—	244	—					

Our Polynomial

Let $d_1, d_2 \in \mathbb{N}$ such that $d_1 \mid (q-1)$ y $d_2 \mid (q-1)$. We are interested in the polynomial:

$$f_{a,b}(X) = X^r \left(X^{\frac{q-1}{d_1}} + aX^{\frac{q-1}{d_2}} + b \right)$$

with $a, b \in \mathbb{F}_q^\times$.

Problem

Our Problem

Study the value set of polynomials of the form

$$f_{a,b}(X) = X^r \left(X^{\frac{q-1}{d_1}} + aX^{\frac{q-1}{d_2}} + b \right)$$

and determine conditions in a, b such that they are permutation polynomials.

Table of Contents

1 Introduction

2 Our Problem

3 Results

The class of equivalence $[a, b]$

$$f_{a,b}(X) = X^r(X^{\frac{q-1}{d_1}} + aX^{\frac{q-1}{d_2}} + b)$$

$a = \alpha^i, b = \alpha^j, \alpha$ a primitive root in \mathbb{F}_q

The class of equivalence $[a, b]$

$$f_{a,b}(X) = X^r(X^{\frac{q-1}{d_1}} + aX^{\frac{q-1}{d_2}} + b)$$

$a = \alpha^i, b = \alpha^j, \alpha$ a primitive root in \mathbb{F}_q

$$(a, b) \sim (a', b') \iff$$

The class of equivalence $[a, b]$

$$f_{a,b}(X) = X^r(X^{\frac{q-1}{d_1}} + aX^{\frac{q-1}{d_2}} + b)$$

$a = \alpha^i, b = \alpha^j, \alpha$ a primitive root in \mathbb{F}_q

$$(a, b) \sim (a', b') \iff$$

$$a' = \alpha^{i+(\frac{q-1}{d_1} - \frac{q-1}{d_2})}$$

The class of equivalence $[a, b]$

$$f_{a,b}(X) = X^r(X^{\frac{q-1}{d_1}} + aX^{\frac{q-1}{d_2}} + b)$$

$a = \alpha^i, b = \alpha^j, \alpha$ a primitive root in \mathbb{F}_q

$$(a, b) \sim (a', b') \iff$$

$$a' = \alpha^{i+(\frac{q-1}{d_1} - \frac{q-1}{d_2})}$$

$$b' = \alpha^{j+(\frac{q-1}{d_1})}$$

The class of equivalence $[a, b]$

$$f_{a,b}(X) = X^r(X^6 + aX^4 + b)$$

$$q = 13, d_1 = 2, d_2 = 3, \alpha = 2$$

The class of equivalence $[a, b]$

$$f_{a,b}(X) = X^r(X^6 + aX^4 + b)$$

$$q = 13, d_1 = 2, d_2 = 3, \alpha = 2$$

$$a = 4 = 2^2, b = 8 = 2^3$$

The class of equivalence $[a, b]$

$$f_{a,b}(X) = X^r(X^6 + aX^4 + b)$$

$$q = 13, d_1 = 2, d_2 = 3, \alpha = 2$$

$$a = 4 = 2^2, b = 8 = 2^3$$

$$(2^2, 2^3) \sim (a', b') \iff a' = 2^{2+(6-4)}, b' = 2^{3+6}$$

The class of equivalence $[a, b]$

$$f_{a,b}(X) = X^r(X^6 + aX^4 + b)$$

$$q = 13, d_1 = 2, d_2 = 3, \alpha = 2$$

$$a = 4 = 2^2, b = 8 = 2^3$$

$$(2^2, 2^3) \sim (a', b') \iff a' = 2^{2+(6-4)}, b' = 2^{3+6}$$

$$(2^2, 2^3) \sim (2^4, 2^9)$$

The class of equivalence $[a, b]$

$$f_{a,b}(X) = X^r(X^{\frac{q-1}{d_1}} + aX^{\frac{q-1}{d_2}} + b)$$


Lemma

The relation \sim defined previously is an equivalence relation.

$f_{a,b}$ with equivalence classes:

$$[f_{a,b}] = [f_{\alpha^i, \alpha^j}] = \{f_{a',b'} \mid (a, b) \sim (a', b')\}$$

Polynomial Results



Value set correspondence

$$f_{a,b}(X) = X^r(X^{\frac{q-1}{d_1}} + aX^{\frac{q-1}{d_2}} + b)$$

Theorem

Suppose that $f_{a,b} \sim f_{a',b'}$ then $|V(f_{a,b})| = |V(f_{a',b'})|$.

Example

Let $q = 13$, $d_1 = 2$, $d_2 = 3$, $a = 4$, $b = 8$. Since $(2^2, 2^3) \sim (2^4, 2^9)$ we have that $|V(f_{2^2,2^3})| = |V(f_{2^4,2^9})| = 7$

Permutation Polynomial correspondence

$$f_{a,b}(X) = X^r(X^{\frac{q-1}{d_1}} + aX^{\frac{q-1}{d_2}} + b)$$

Corollary

Suppose that $f_{a,b}$ is a permutation polynomial and $f_{a,b} \sim f_{a',b'}$, then $f_{a',b'}$ is also a permutation polynomial.

Size of equivalence classes

$$f_{a,b}(X) = X^r(X^{\frac{q-1}{d_1}} + aX^{\frac{q-1}{d_2}} + b)$$

Proposition

$|[f_{a,b}]| = \text{lcm}(d_1, d_2)$ where $\text{lcm}(x, y)$ is the least common multiple of x and y .


Example

Let $q = 13$, $d_1 = 2$, $d_2 = 3$, $a = 4$, $b = 8$. Note that $\text{lcm}(2, 3) = 6$. These are the elements of (a, b) :

$$\begin{array}{ccccccc} (2^2, 2^3) & (2^4, 2^9) & (2^6, 2^3) & (2^8, 2^9) & (2^{10}, 2^3) & (2^{12}, 2^9) & (2^2, 2^3) \\ (4, 8) & (3, 5) & (12, 8) & (9, 5) & (10, 8) & (1, 5) & (4, 8) \end{array}$$

Polynomials Results

Number of Permutation Polynomials



Polynomial Results

Proposition

The number of polynomials of the form $f_{a,b}(X)$ with $|V(f_{a,b})| = n$ is a multiple of $\text{lcm}(d_1, d_2)$

Corollary

The number of permutation polynomials of the form $f_{a,b}(X)$ is a multiple of $\text{lcm}(d_1, d_2)$

Future Work

- Find necessary and sufficient conditions such that $V(f_{a,b}) = \mathbb{F}_q$
- Generalize results to polynomials with more terms and with exponents not divisors of $q - 1$:

$$f_{a,b}(X) = X^r(X^{d_1} + aX^{d_2} + b)$$