

# Construction of Families of Permutation Trinomials over Finite Fields

Christian A. Rodriguez  
Alex D. Santos

Department of Computer Science  
University of Puerto Rico, Río Piedras

March 4, 2014

# Table of Contents

- 1 Introduction
- 2 Our Problem
- 3 Results

# Table of Contents

1 Introduction

2 Our Problem

3 Results

# Finite Fields

## Definition

A **finite field**  $\mathbb{F}_q$  is a field with  $q = p^r$  elements where  $p$  is prime.

## Example

$$\mathbb{F}_7 = \{0, 1, 2, 3, 4, 5, 6\}$$

### Addition:

$$\begin{aligned} 2 + 2 &= 4 \\ 4 + 4 &= 8 \\ (\text{mod } 7) &= 1 \end{aligned}$$

### Multiplication:

$$\begin{aligned} 2 \cdot 2 &= 4 \\ 4 \cdot 4 &= 16 \\ (\text{mod } 7) &= 2 \end{aligned}$$

# Value Sets

## Definition

Let  $f(x)$  be a polynomial defined over a finite field  $\mathbb{F}_q$ . Then the **value set** of  $f$  is defined as  $V(f) = \{f(a) \mid a \in \mathbb{F}_q\}$

## Example

Consider  $f(x) = x^2$  defined over  $\mathbb{F}_5$ . Note:  
 $f(0) = 0, f(1) = 1, f(2) = 4, f(3) = 4, f(4) = 1$ , so  
 $V(f) = \{0, 1, 4\}$ .

# Permutation Polynomials

## Definition

A polynomial  $f(x)$  defined over  $\mathbb{F}_q$  is a **permutation polynomial** if and only if  $V(f) = \mathbb{F}_q$ .

# Permutation Polynomials

## Definition

A polynomial  $f(x)$  defined over  $\mathbb{F}_q$  is a **permutation polynomial** if and only if  $V(f) = \mathbb{F}_q$ .

## Example

Let  $f(x) = x^3$  over  $\mathbb{F}_5$ . Note:  $V(f) = \{0, 1, 3, 2, 4\}$  so  $f(x)$  is a permutation polynomial over  $\mathbb{F}_5$

# Permutation Polynomials

## Definition

A polynomial  $f(x)$  defined over  $\mathbb{F}_q$  is a **permutation polynomial** if and only if  $V(f) = \mathbb{F}_q$ .

## Example

Let  $f(x) = x^3$  over  $\mathbb{F}_5$ . Note:  $V(f) = \{0, 1, 3, 2, 4\}$  so  $f(x)$  is a permutation polynomial over  $\mathbb{F}_5$

## Example

Let  $f(x) = x^2$  over  $\mathbb{F}_5$ . We have that  $V(f) = \{0, 1, 4\}$  so  $f(x)$  is not a permutation polynomial over  $\mathbb{F}_5$ .



# Primitive Roots

## Definition

A **primitive root**  $\alpha \in \mathbb{F}_q$  is a generator for the multiplicative group  $\mathbb{F}_q^\times$

# Primitive Roots

## Definition

A **primitive root**  $\alpha \in \mathbb{F}_q$  is a generator for the multiplicative group  $\mathbb{F}_q^\times$

$$\mathbb{F}_7$$

# Primitive Roots

## Definition

A **primitive root**  $\alpha \in \mathbb{F}_q$  is a generator for the multiplicative group  $\mathbb{F}_q^\times$

$$\mathbb{F}_7$$

$$\begin{array}{rcl} 3^1 & = & 3 \\ 3^2 & = & 2 \\ 3^3 & = & 6 \\ 3^4 & = & 4 \\ 3^5 & = & 5 \\ 3^6 & = & 1 \end{array}$$

# Primitive Roots

## Definition

A **primitive root**  $\alpha \in \mathbb{F}_q$  is a generator for the multiplicative group  $\mathbb{F}_q^\times$

$$\mathbb{F}_7$$

$$\begin{aligned} 3^1 &= 3 \\ 3^2 &= 2 \\ 3^3 &= 6 \\ 3^4 &= 4 \\ 3^5 &= 5 \\ 3^6 &= 1 \end{aligned}$$

$$\begin{aligned} 2^1 &= 2 \\ 2^2 &= 4 \\ 2^3 &= 1 \\ 2^4 &= 2 \\ 2^5 &= 4 \\ 2^6 &= 1 \end{aligned}$$

# Table of Contents

1 Introduction

2 Our Problem

3 Results

# Our Problem

- Everything is known about Permutation Monomials
- Permutation Binomials have been studied extensively
- The next case is to study Permutation Trinomials

# Our Polynomial

Let  $d_1, d_2 \in \mathbb{N}$  such that  $d_1 \mid (q-1)$  y  $d_2 \mid (q-1)$ . We are interested in the polynomial:

$$f_{a,b}(X) = X^r \left( X^{\frac{q-1}{d_1}} + aX^{\frac{q-1}{d_2}} + b \right)$$

with  $a, b \in \mathbb{F}_q^\times$ .

Denote the value set of this polynomial  $V(f_{a,b})$ .

# Our Polynomial

Let  $d_1, d_2 \in \mathbb{N}$  such that  $d_1 \mid (q-1)$  y  $d_2 \mid (q-1)$ . We are interested in the polynomial:

$$f_{a,b}(X) = X^r \left( X^{\frac{q-1}{d_1}} + aX^{\frac{q-1}{d_2}} + b \right)$$

with  $a, b \in \mathbb{F}_q^\times$ .

Denote the value set of this polynomial  $V(f_{a,b})$ .



# Problem

## Our Problem

*Study the value set of polynomials of the form*

*$f_{a,b}(X) = X^r(X^{\frac{q-1}{d_1}} + aX^{\frac{q-1}{d_2}} + b)$  and determine conditions in  $a, b$  such that they are permutation polynomials.*

# Table of Contents

1 Introduction

2 Our Problem

3 Results

# The class of equivalence $[a, b]$

$$f_{a,b}(X) = X^r(X^{\frac{q-1}{d_1}} + aX^{\frac{q-1}{d_2}} + b)$$

$a = \alpha^i, b = \alpha^j, \alpha$  a primitive root in  $\mathbb{F}_q$

# The class of equivalence $[a, b]$

$$f_{a,b}(X) = X^r(X^{\frac{q-1}{d_1}} + aX^{\frac{q-1}{d_2}} + b)$$

$a = \alpha^i, b = \alpha^j$ ,  $\alpha$  a primitive root in  $\mathbb{F}_q$

$\sim$  defined as  $(a, b) \sim (a', b') \iff$

# The class of equivalence $[a, b]$

$$f_{a,b}(X) = X^r(X^{\frac{q-1}{d_1}} + aX^{\frac{q-1}{d_2}} + b)$$

$a = \alpha^i, b = \alpha^j$ ,  $\alpha$  a primitive root in  $\mathbb{F}_q$

$\sim$  defined as  $(a, b) \sim (a', b') \iff$   
 $a' = \alpha^{i+h(\frac{q-1}{d_1} - \frac{q-1}{d_2})}$

# The class of equivalence $[a, b]$

$$f_{a,b}(X) = X^r(X^{\frac{q-1}{d_1}} + aX^{\frac{q-1}{d_2}} + b)$$

$a = \alpha^i, b = \alpha^j, \alpha$  a primitive root in  $\mathbb{F}_q$

$\sim$  defined as  $(a, b) \sim (a', b') \iff$

$$a' = \alpha^{i+h(\frac{q-1}{d_1} - \frac{q-1}{d_2})}$$

$$b' = \alpha^{j+h(\frac{q-1}{d_1})}$$

# The class of equivalence $[a, b]$

$$q = 13$$

$$d_1 = 2$$

$$d_2 = 3$$

$$\alpha = 2$$

$$a = 4 = 2^2$$

$$b = 8 = 2^3$$

# The class of equivalence $[a, b]$

$$q = 13$$

$$d_1 = 2$$

$$d_2 = 3$$

$$\alpha = 2$$

$$a = 4 = 2^2$$

$$b = 8 = 2^3$$

$$(2^2, 2^3) \sim (a', b') \iff a' = 2^{2+2h}, b' = 2^{3+6h}$$



# The class of equivalence $[a, b]$

$$q = 13$$

$$d_1 = 2$$

$$d_2 = 3$$

$$\alpha = 2$$

$$a = 4 = 2^2$$

$$b = 8 = 2^3$$

$$(2^2, 2^3) \sim (a', b') \iff a' = 2^{2+2h}, b' = 2^{3+6h}$$

$$\text{Example } (h = 1): (2^2, 2^3) \sim (2^4, 2^9).$$

# The class of equivalence $[a, b]$


## Lemma

*The relation  $\sim$  defined previously is an equivalence relation.*

$f_{a,b}$  with equivalence classes:

$$[f_{a,b}] = [f_{\alpha^i, \alpha^j}] = \{f_{a',b'} \mid (a,b) \sim (a',b')\}$$

# Polynomial Results



# Value set correspondence

$$f_{a,b}(X) = X^r(X^{\frac{q-1}{d_1}} + aX^{\frac{q-1}{d_2}} + b)$$

## Theorem

*Suppose that  $f_{a,b} \sim f_{a',b'}$  then  $|V(f_{a,b})| = |V(f_{a',b'})|$ .*

## Example

Let  $q = 13$ ,  $d_1 = 2$ ,  $d_2 = 3$ ,  $a = 4$ ,  $b = 8$ . Since  $(2^2, 2^3) \sim (2^4, 2^9)$  we have that  $|V(f_{2^2, 2^3})| = |V(f_{2^4, 2^9})|$

# Size of equivalence classes

$$f_{a,b}(X) = X^r \left( X^{\frac{q-1}{d_1}} + aX^{\frac{q-1}{d_2}} + b \right)$$

## Proposition

$|[f_{a,b}]| = \text{lcm}(d_1, d_2)$  where  $\text{lcm}(x, y)$  is the least common multiple of  $x$  and  $y$ .


## Example

Let  $q = 13$ ,  $d_1 = 2$ ,  $d_2 = 3$ ,  $a = 4$ ,  $b = 8$ . Note that  $\text{lcm}(2, 3) = 6$ . These are the elements of  $(a, b)$ :

$$(2^2, 2^3), (2^4, 2^9), (2^6, 2^3), (2^8, 2^9), (2^{10}, 2^3), (2^{12}, 2^9), (2^2, 2^3)$$

# Polynomials Results

Number of Permutation Polynomials



\* Note that the number of polynomials in each cell is  $6 = \text{lcm}(2, 3)$

\* The polynomials within each cell have value sets of the same size. The size of the value sets associated to different cells might or might not be equal.

# Polynomial Results

## Proposition

*The number of polynomials of the form  $f_{a,b}(X)$  with  $|V(f_{a,b})| = n$  is a multiple of  $\text{lcm}(d_1, d_2)$*

## Corollary

*The number of permutation polynomials of the form  $f_{a,b}(X)$  is a multiple of  $\text{lcm}(d_1, d_2)$*

# Future Work

- Find necessary and sufficient conditions such that  $V(f_{a,b}) = \mathbb{F}_q$
- Collect data on number of permutation polynomials of the form  $f_{a,b}$  for different values of  $d_1$  and  $d_2$  and compare results with number of permutation polynomials.