

Imaging Geometry

Computer Vision

Modelling from 3D to 2D world



Perspective Matters!!

Taj Mahal










Paris town hall
“Anamorphosis”




Cricket



Cameras, Multiple Views, and Motion

-  Imaging Geometry
 - Image transforms like scaling, rotation etc
-  Perspective Transformation
-  Projective Transformation
-  Camera Model
-  Camera Calibration

Camera Dimensionality Reduction Machine (3D to 2D)



How to recover knowledge about 3D from 2D...?? ✓

Common transformations



Original

Transformed



Translation



Rotation



Scaling




Affine



Perspective

✓ Parametric (global) transformations



Transformation T is a coordinate-changing machine:

$$p' = T(p)$$

What does it mean that T is global?

– T is the same for any point p ✓

T can be described by just a few numbers (parameters) ✓

①

For linear transformations, we can represent T as a matrix

$$p' = Tp$$

$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \mathbf{T} \begin{bmatrix} x \\ y \end{bmatrix}$$

Common transformations



Original

Transformed



Translation



Rotation



Scaling




Affine



Perspective


Scaling

- *Scaling* a coordinate means multiplying each of its components by a scalar ✓
- *Uniform scaling* means this scalar is the same for all components:




Scaling


- *Non-uniform scaling*: different scalars per component:



Scaling



$$\begin{aligned} x' &= ax \\ y' &= by \end{aligned}$$



- Scaling operation:


$$x' = ax$$

$$y' = by$$

$$I(x, y) = I(x', y')$$

how to compute (x', y')

- Or, in matrix form:




$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} a & 0 \\ 0 & b \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

scaling matrix S

$$\begin{aligned} x' &= ax + 0y \\ y' &= 0x + by \\ x' &= ax \\ y' &= by \end{aligned}$$

2-D Rotation



Polar coordinates...

$$x = r \cos(\phi)$$

$$y = r \sin(\phi)$$

$$x' = r \cos(\phi + \theta)$$

$$y' = r \sin(\phi + \theta)$$

Trig Identity...

$$x' = r \cos(\phi) \cos(\theta) - r \sin(\phi) \sin(\theta)$$

$$y' = r \sin(\phi) \cos(\theta) + r \cos(\phi) \sin(\theta)$$

Substitute...

$$x' = x \cos(\theta) - y \sin(\theta)$$

$$y' = x \sin(\theta) + y \cos(\theta)$$

$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

2-D Rotation

This is easy to capture in matrix form:

$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \underbrace{\begin{bmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{bmatrix}}_{\mathbf{R}} \begin{bmatrix} x \\ y \end{bmatrix}$$

Even though $\sin(\theta)$ and $\cos(\theta)$ are nonlinear functions of θ ,

- x' is a linear combination of x and y
- y' is a linear combination of x and y

What is the inverse transformation?

- Rotation by $-\theta$
- For rotation matrices $\mathbf{R}^{-1} = \mathbf{R}^T$


Basic 2D transformations

$$\begin{aligned} x' &= x + \alpha_x y \\ y' &= a_y x + y \end{aligned}$$

①


$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} s_x & 0 \\ 0 & s_y \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

Scale



$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} 1 & \alpha_x \\ \alpha_y & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

Shear



②

$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} \cos \Theta & -\sin \Theta \\ \sin \Theta & \cos \Theta \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

Rotate

③

$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

Translate

$$\begin{aligned} x' &= x + t_x \\ y' &= y + t_y \end{aligned}$$

④

$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} a & b & c \\ d & e & f \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

Affine

Affine is any combination of translation, scale, rotation, and shear

Affine Transformations

- ① Affine transformations are combinations of
- Linear transformations, and
 - Translations *(S, T, R, Sh)*







Properties of affine transformations:

- Lines map to lines
- Parallel lines remain parallel
- Ratios are preserved
- Closed under composition

$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} a & b & c \\ d & e & f \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$


or

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} a & b & c \\ d & e & f \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

Transformation Name	Affine Matrix, A	Coordinate Equations	Example
Identity	$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$	$\begin{aligned} x' &= x \\ y' &= y \end{aligned}$	
Scaling/Reflection (For reflection, set one scaling factor to -1 and the other to 0)	$\begin{bmatrix} c_x & 0 & 0 \\ 0 & c_y & 0 \\ 0 & 0 & 1 \end{bmatrix}$	$\begin{aligned} x' &= c_x x \\ y' &= c_y y \end{aligned}$	
Rotation (about the origin)	$\begin{bmatrix} \cos \theta & -\sin \theta & 0 \\ \sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{bmatrix}$	$\begin{aligned} x' &= x \cos \theta - y \sin \theta \\ y' &= x \sin \theta + y \cos \theta \end{aligned}$	
Translation	$\begin{bmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \\ 0 & 0 & 1 \end{bmatrix}$	$\begin{aligned} x' &= x + t_x \\ y' &= y + t_y \end{aligned}$	
Shear (vertical)	$\begin{bmatrix} 1 & s_v & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$	$\begin{aligned} x' &= x + s_v y \\ y' &= y \end{aligned}$	
Shear (horizontal)	$\begin{bmatrix} 1 & 0 & 0 \\ s_h & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$	$\begin{aligned} x' &= x \\ y' &= s_h x + y \end{aligned}$	

basic
image
transform

2D image transformations (reference table)



Transformation	Matrix	# DoF	Preserves	Icon
① translation	$\begin{bmatrix} I & t \end{bmatrix}_{2 \times 3}$	2	orientation	
② rigid (Euclidean)	$\begin{bmatrix} R & t \end{bmatrix}_{2 \times 3}$	3	lengths	
③ similarity	$\begin{bmatrix} sR & t \end{bmatrix}_{2 \times 3}$	4	angles	
④ affine	$\begin{bmatrix} A \end{bmatrix}_{2 \times 3}$	6	parallelism	
projective	$\begin{bmatrix} \tilde{H} \end{bmatrix}_{3 \times 3}$	8	straight lines	

‘Homography’

Table 2.1 Hierarchy of 2D coordinate transformations. Each transformation also preserves the properties listed in the rows below it, i.e., similarity preserves not only angles but also parallelism and straight lines. The 2×3 matrices are extended with a third $[0^T \ 1]$ row to form a full 3×3 matrix for homogeneous coordinate transformations.

Projective Transformations

Projective transformations are combos of

- ① Affine transformations, and $(S+R+T+Sh)$
- ② Projective warps


$$\begin{bmatrix} x' \\ y' \\ w' \end{bmatrix} = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} \begin{bmatrix} x \\ y \\ w \end{bmatrix}$$

$\cdot 3 \times 3$

Properties of projective transformations:

- Lines map to lines ✓
- ✓ Parallel lines do not necessarily remain parallel
- ✓ Ratios are not preserved
- Models change of basis
- ✓ Projective matrix is defined up to a scale (8 DOF)

we use projective transforms to create a
360 panorama



- In order to figure this out, we need to learn what a **camera** is

The Geometry of Image Formation

Szeliski 2.1, parts of 2.2

To understand projective transfer

Mapping between image and world coordinates

- Pinhole camera model ✓
- Projective geometry ✓
 - Vanishing points and lines
- Projection matrix ✓

Image Formation: Orthographic Projection

- Means of representing 3-dimensional objects in 2-Dimensions.
- It is a form of parallel projection, in which all the projection lines are orthogonal to the projection plane




FIGURE 1.6: Orthographic projection. Unlike other geometric models of the image formation process, orthographic projection does not involve a reversal of image features.

Orthographic Projections

- A simple orthographic projection onto the plane $z = 0$ can be defined by the following matrix:

$$P = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

- For each point $\checkmark v = (v_x, v_y, v_z)$, the transformed point Pv would be

$$Pv = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} v_x \\ v_y \\ v_z \end{bmatrix} = \begin{bmatrix} v_x \\ v_y \\ 0 \end{bmatrix}$$

$$\begin{aligned} Pv_x &= v_x + 0 + 0 \\ Pv_y &= 0 + v_y + 0 \\ Pv_z &= 0 + 0 + 0 \end{aligned} \quad Pv = \begin{bmatrix} Pv_x = x \\ Pv_y = y \end{bmatrix}$$

- Often, it is more useful to use homogeneous coordinates. The transformation in homogeneous coordinates

$$(v_x, v_y, v_z) \rightarrow (v_x, v_y, v_z, 1)$$

- For each homogeneous vector $v = (v_x, v_y, v_z, 1)$, the transformed vector Pv would be


$$P = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \quad 4 \times 4$$

$$Pv = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} v_x \\ v_y \\ v_z \\ 1 \end{bmatrix} = \begin{bmatrix} v_x \\ v_y \\ 0 \\ 1 \end{bmatrix}$$

$\rightarrow v_x' = v_x$
 $\rightarrow v_y' = v_y$
 $\rightarrow 1$

$$(v_x', v_y', 1)$$


Pinhole camera model



- ✓ f = Focal length
- ✓ c = Optical center of the camera

Perspective Projection

Projection: world coordinates \rightarrow image coordinates



$$\underline{\underline{\mathbf{p}}} = \begin{bmatrix} \underline{\underline{u}} \\ \underline{\underline{v}} \end{bmatrix}$$

\mathbf{p} = distance from image center

$$\underline{\underline{\mathbf{P}}} = \begin{bmatrix} X \\ Y \\ Z \end{bmatrix}$$

$$\frac{f}{Z} = -\frac{v}{Y} \therefore v = -\frac{f}{Z} Y$$

$$\therefore v = -X \cdot \frac{f}{Z}$$

$$\frac{f}{Z} = -\frac{u}{X}$$

$$\underline{\underline{u}} = -X * \frac{f}{Z}$$

x comp of 3D point
comp of the img 2D

$$\underline{\underline{v}} = -Y * \frac{f}{Z}$$


y of the 3D pt
y comp of 2D

What is the effect if f and Z are equal?

Taj Mahal




Taj Mahal




Perspective Transforms

- Parallel lines in the world intersect in the projected image at a “vanishing point”.
- Parallel lines on the same plane in the world converge to vanishing points on a “vanishing line”.



Perspective Projection:

- (<https://www.youtube.com/watch?v=17kqhGRDHc8>)



Basic

Scale \rightarrow 2D

Rot \rightarrow 2D

Shear \rightarrow 2D

Trans

(1) Affine = S + R + S + T

(2) Euclidean \Rightarrow R + T

(3) Similarity \rightarrow R + T + S

(4) Projective Transf.

3D \rightarrow 2D.


↓
Ortho
|||

↓
Persp
X

3D Rigid Body Transform

Euclidean (R+T)

3D object
→ R.
T.



Transform

Object in the 3D?

How the 3D world point
3x3 of the object changes
when the object R&T.

$$\begin{bmatrix} X' \\ Y' \\ Z' \end{bmatrix} = R \begin{bmatrix} X \\ Y \\ Z \end{bmatrix} + T = \begin{bmatrix} r_{11} & r_{12} & r_{13} \\ r_{21} & r_{22} & r_{23} \\ r_{31} & r_{32} & r_{33} \end{bmatrix} \begin{bmatrix} X \\ Y \\ Z \end{bmatrix} + \begin{bmatrix} T_X \\ T_Y \\ T_Z \end{bmatrix}$$


New 3D point old 3D p.

Rotation matrix (9 unknowns)

Translation (3 unknowns)


Rotation

$$R = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$




Rotation

$$R = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$
$$R = \begin{bmatrix} \cos\Theta & -\sin\Theta & 0 \\ \sin\Theta & \cos\Theta & 0 \\ 0 & 0 & 1 \end{bmatrix}$$



Rotation

- About z axis



Homogeneous $\left\{ \begin{matrix} x' \\ y' \\ z' \\ 1 \end{matrix} \right\} = \begin{bmatrix} \cos \theta & -\sin \theta & 0 & 0 \\ \sin \theta & \cos \theta & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \\ 1 \end{bmatrix}$

Rotation

- About x axis:


$$\begin{bmatrix} x' \\ y' \\ z' \\ 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \underline{\cos} \theta & -\underline{\sin} \theta & 0 \\ 0 & \underline{\sin} \theta & \underline{\cos} \theta & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \\ 1 \end{bmatrix}$$

- About y axis:

$$\begin{bmatrix} x' \\ y' \\ z' \\ 1 \end{bmatrix} = \begin{bmatrix} \underline{\cos} \theta & 0 & \underline{\sin} \theta & 0 \\ 0 & 1 & 0 & 0 \\ -\underline{\sin} \theta & 1 & \underline{\cos} \theta & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \\ 1 \end{bmatrix}$$


✓ Euler Angles

$\begin{bmatrix} x' & y' & z' \end{bmatrix}$
 0°




$\alpha: \begin{bmatrix} \cdot & 0 & \cdot \end{bmatrix}$

$\begin{bmatrix} x' & y' & z' \end{bmatrix}$
 0°



$\beta: \begin{bmatrix} \cdot & 0 & \cdot \end{bmatrix}$


$\begin{bmatrix} x' & y' & z' \end{bmatrix}$
 0°



$\gamma: \begin{bmatrix} \cdot & 0 & \cdot \end{bmatrix}$


$$\underline{\underline{R}} = \underline{\underline{R}}_x(0^\circ) \underline{\underline{R}}_y(0^\circ) \underline{\underline{R}}_z(0^\circ) = \begin{bmatrix} 1.000 & 0.000 & 0.000 \\ 0.000 & 1.000 & 0.000 \\ 0.000 & 0.000 & 1.000 \end{bmatrix}$$

Euler Angles




$$\mathbf{R} = \mathbf{R}_x(20^\circ) \mathbf{R}_y(0^\circ) \mathbf{R}_z(0^\circ) = \begin{bmatrix} 1.000 & 0.000 & 0.000 \\ 0.000 & 0.940 & -0.342 \\ 0.000 & 0.342 & 0.940 \end{bmatrix}$$


Euler Angles



$\alpha: \langle 20 \rangle$




$\beta: \langle \underline{30} \rangle$




$\gamma: \langle 0 \rangle$

$$\mathbf{R} = \underline{\mathbf{R}_x(20^\circ)} \underline{\mathbf{R}_y(30^\circ)} \underline{\mathbf{R}_z(0^\circ)} = \begin{bmatrix} 0.866 & 0.000 & 0.500 \\ 0.171 & 0.940 & -0.296 \\ -0.470 & 0.342 & 0.814 \end{bmatrix}$$


Euler Angles



$\alpha: < 20 >$



$\beta: < 30 >$



$\gamma: < 40 >$

$$R = R_x(20^\circ) R_y(30^\circ) R_z(40^\circ) = \begin{bmatrix} 0.663 & -0.557 & 0.500 \\ 0.735 & 0.610 & -0.296 \\ -0.140 & 0.564 & 0.814 \end{bmatrix}$$

Euler Angles


$$R = R_z^\alpha R_y^\beta R_x^\gamma = \begin{matrix} R_z(\alpha) \\ \left[\begin{array}{ccc} \cos \alpha & -\sin \alpha & 0 \\ \sin \alpha & \cos \alpha & 0 \\ 0 & 0 & 1 \end{array} \right] \end{matrix} \begin{matrix} R_y(\beta) \\ \left[\begin{array}{ccc} \cos \beta & 0 & \sin \beta \\ 0 & 1 & 0 \\ -\sin \beta & 0 & \cos \beta \end{array} \right] \end{matrix} \begin{matrix} R_x(\gamma) \\ \left[\begin{array}{ccc} 1 & 0 & 0 \\ 0 & \cos \gamma & -\sin \gamma \\ 0 & \sin \gamma & \cos \gamma \end{array} \right] \end{matrix}$$

$$R = R_z^\alpha R_y^\beta R_x^\gamma = \begin{bmatrix} \cos \alpha \cos \beta & \cos \alpha \sin \beta \sin \gamma - \sin \alpha \cos \gamma & \cos \alpha \sin \beta \cos \gamma + \sin \alpha \sin \gamma \\ \sin \alpha \cos \beta & \sin \alpha \sin \beta \sin \gamma + \cos \alpha \cos \gamma & \sin \alpha \sin \beta \cos \gamma - \cos \alpha \sin \gamma \\ -\sin \beta & \cos \beta \sin \gamma & \cos \beta \cos \gamma \end{bmatrix}$$

$$R = \begin{bmatrix} 1 & -\alpha & \beta \\ \alpha & 1 & -\gamma \\ -\beta & \gamma & 1 \end{bmatrix}$$

if angles are small $\cos \Theta \approx 1$ $\sin \Theta \approx \Theta$

Important Definitions



- **Frame of reference:** a measurements are made with respect to a particular coordinate system called the frame of reference.
- **World Frame:** a fixed coordinate system for representing objects (points, lines, surfaces, etc.) in the world.
- **Camera Frame:** coordinate system that uses the camera center as its origin (and the optic axis as the Z-axis)
- **Image or retinal plane:** plane on which the image is formed, note that the image plane is measured in camera frame coordinates (mm)
- **Image Frame:** coordinate system that measures pixel locations in the image plane.
- **Intrinsic Parameters:** Camera parameters that are internal and fixed to a particular camera/digitization setup
- **Extrinsic Parameters:** Camera parameters that are external to the camera and may change with respect to the world frame.

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} f \\ c \end{bmatrix} \begin{bmatrix} x' \\ y' \\ z' \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} R \\ T \end{bmatrix} \begin{bmatrix} x' \\ y' \\ z' \end{bmatrix}$$

3D point $\begin{bmatrix} x \\ y \\ z \end{bmatrix}$ = 2D $\begin{bmatrix} x' \\ y' \end{bmatrix}$!