


First Order Differential Equations

2.1

5.(a)




(b) If $y(0) > -2$, solutions eventually have positive slopes, and hence increase without bound. If $y(0) \leq -2$, solutions have negative slopes and decrease without bound.

(c) The integrating factor is $\mu(t) = e^{-\int 3dt} = e^{-3t}$. The differential equation can be written as $e^{-3t}y' - 3e^{-3t}y = 4e^{-2t}$, that is, $(e^{-3t}y)' = 4e^{-2t}$. Integration of both sides of the equation results in the general solution $y(t) = -2e^t + ce^{3t}$. It follows that all solutions will increase exponentially if $c > 0$ and will decrease exponentially

if $c \leq 0$. Letting $c = 0$ and then $t = 0$, we see that the boundary of these behaviors is at $y(0) = -2$.


9.(a)



(b) All solutions eventually have positive slopes, and hence increase without bound.

(c) The integrating factor is $\mu(t) = e^{\int (1/2) dt} = e^{t/2}$. The differential equation can be written as $e^{t/2}y' + e^{t/2}y/2 = 4te^{t/2}/2$, that is, $(e^{t/2}y/2)' = 2te^{t/2}$. Integration of both sides of the equation results in the general solution $y(t) = 4t - 8 + ce^{-t/2}$. All solutions approach the specific solution $y_0(t) = 4t - 8$.


10.(a)



(b) For $y > 0$, the slopes are all positive, and hence the corresponding solutions increase without bound. For $y < 0$, almost all solutions have negative slopes, and hence solutions tend to decrease without bound.

(c) First divide both sides of the equation by t ($t > 0$). From the resulting standard form, the integrating factor is $\mu(t) = e^{-\int (1/t) dt} = 1/t$. The differential equation can be written as $y'/t - y/t^2 = te^{-2t}$, that is, $(y/t)' = te^{-2t}$. Integration leads to the general solution $y(t) = -te^{-2t}/2 + ct$. For $c \neq 0$, solutions diverge, as implied by the direction field. For the case $c = 0$, the specific solution is $y(t) = -te^{-2t}/2$, which evidently approaches zero as $t \rightarrow \infty$.

12.(a)



(b) All solutions eventually have positive slopes, and hence increase without bound.

(c) The integrating factor is $\mu(t) = e^{t/2}$. The differential equation can be written as $e^{t/2}y' + e^{t/2}y/2 = 4e^{t/2}t^2/2$, that is, $(e^{t/2}y/2)' = 2e^{t/2}t^2$. Integration of both sides of the equation results in the general solution $y(t) = 4t^2 - 16t + 32 + ce^{-t/2}$. It follows that all solutions converge to the specific solution $4t^2 - 16t + 32$.


14. The integrating factor is $\mu(t) = e^{3t}$. After multiplying both sides by $\mu(t)$, the equation can be written as $(e^{3t}y)' = t$. Integrating both sides of the equation results in the general solution $y(t) = t^2e^{-3t}/2 + ce^{-3t}$. Invoking the specified condition, we require that $e^{-3}/2 + ce^{-3} = 0$. Hence $c = -1/2$, and the solution to the initial value problem is $y(t) = (t^2 - 1)e^{-3t}/2$.

16. The integrating factor is $\mu(t) = e^{\int (3/t) dt} = t^3$. Multiplying both sides by $\mu(t)$, the equation can be written as $(t^3y)' = \cos t$. Integrating both sides of the equation results in the general solution $y(t) = \sin t/t^3 + ct^{-3}$. Substituting $t = \pi$ and setting the value equal to zero gives $c = 0$. Hence the specific solution is $y(t) = \sin t/t^3$.

17. The integrating factor is $\mu(t) = e^{-4t}$, and the differential equation can be written as $(e^{-4t}y)' = 1$. Integrating, we obtain $e^{-4t}y(t) = t + c$. Invoking the specified initial condition results in the solution $y(t) = (t + 2)e^{4t}$.

19. After writing the equation in standard form, we find that the integrating factor is $\mu(t) = e^{\int (5/t) dt} = t^5$. Multiplying both sides by $\mu(t)$, the equation can be written as $(t^5y)' = te^{-t}$. Integrating both sides results in $t^5y(t) = -(t + 1)e^{-t} + c$. Letting $t = -1$ and setting the value equal to zero gives $c = 0$. Hence the specific solution of the initial value problem is $y(t) = -(t^{-4} + t^{-5})e^{-t}$.

22.(a)




The solutions eventually increase or decrease, depending on the initial value a . The critical value seems to be $a_0 = -2$.

(b) The integrating factor is $\mu(t) = e^{-t/2}$, and the general solution of the differential equation is $y(t) = -2e^{t/4} + ce^{t/4}$. Invoking the initial condition $y(0) = a$, the solution may also be expressed as $y(t) = -2e^{t/4} + (a + 2)e^{t/2}$. The critical value is $a_0 = -2$.

(c) For $a_0 = -2$, the solution is $y(t) = -2e^{t/4}$, which diverges to $-\infty$ as $t \rightarrow \infty$.

23.(a)




Solutions appear to grow infinitely large in absolute value, with signs depending on the initial value $y(0) = a_0$. The direction field appears horizontal for $a_0 \approx -1/8$.

(b) Dividing both sides of the given equation by 3, the integrating factor is $\mu(t) = e^{-2t/3}$. Multiplying both sides of the original differential equation by $\mu(t)$ and integrating results in $y(t) = (2e^{2t/3} - 2e^{-\pi t/2} + a(4 + 3\pi)e^{2t/3})/(4 + 3\pi)$. The qualitative behavior of the solution is determined by the terms containing $e^{2t/3}$: $2e^{2t/3} + a(4 + 3\pi)e^{2t/3}$. The nature of the solutions will change when $2 + a(4 + 3\pi) = 0$. Thus the critical initial value is $a_0 = -2/(4 + 3\pi)$.

(c) In addition to the behavior described in part (a), when $y(0) = -2/(4 + 3\pi)$, the solution is $y(t) = (-2e^{-\pi t/2})/(4 + 3\pi)$, and that specific solution will converge to $y = 0$.

24.(a)



As $t \rightarrow 0$, solutions increase without bound if $y(1) = a > 0.8$, and solutions decrease without bound if $y(1) = a < 0.8$.

(b) The integrating factor is $\mu(t) = e^{\int (t+1)/t dt} = t e^t$. The general solution of the differential equation is $y(t) = 2t e^{-t} + c e^{-t}/t$. Since $y(1) = a$, we have that $2 + c = ae$. That is, $c = ae - 2$. Hence the solution can also be expressed as $y(t) = 2t e^{-t} + (ae - 2) e^{-t}/t$. For small values of t , the second term is dominant. Setting $ae - 2 = 0$, the critical value of the parameter is $a_0 = 2/e$.

(c) When $a = 2/e$, the solution is $y(t) = 2t e^{-t}$, which approaches 0 as $t \rightarrow 0$.

27. The integrating factor is $\mu(t) = e^{\int (1/2) dt} = e^{t/2}$. Therefore the general solution is $y(t) = (4 \cos t + 8 \sin t)/5 + c e^{-t/2}$. Invoking the initial condition, the specific solution is $y(t) = (4 \cos t + 8 \sin t - 9 e^{-t/2})/5$. Differentiating, it follows that $y'(t) = (-4 \sin t + 8 \cos t + 4.5 e^{-t/2})/5$ and $y''(t) = (-4 \cos t - 8 \sin t - 2.25 e^{-t/2})/5$. Setting $y'(t) = 0$, the first solution is $t_1 = 1.3643$, which gives the location of the first stationary point. Since $y''(t_1) < 0$, the first stationary point is a local maximum. The coordinates of the point are $(1.3643, 0.82008)$.

28. The integrating factor is $\mu(t) = e^{\int (2/3) dt} = e^{2t/3}$, and the differential equation can be written as $(e^{2t/3} y)' = e^{2t/3} - t e^{2t/3}/2$. The general solution is $y(t) = (21 - 6t)/8 + c e^{-2t/3}$. Imposing the initial condition, we have $y(t) = (21 - 6t)/8 + (y_0 - 21/8) e^{-2t/3}$. Since the solution is smooth, the desired intersection will be a point of tangency. Taking the derivative, $y'(t) = -3/4 - (2y_0 - 21/4) e^{-2t/3}/3$. Setting $y'(t) = 0$, the solution is $t_1 = (3/2) \ln[(21 - 8y_0)/9]$. Substituting into the solution, the respective value at the stationary point is $y(t_1) = 3/2 + (9/4) \ln 3 - (9/8) \ln(21 - 8y_0)$. Setting this result equal to zero, we obtain the required initial value $y_0 = (21 - 9 e^{4/3})/8 \approx -1.643$.

29.(a) The integrating factor is $\mu(t) = e^{t/4}$, and the differential equation can be written as $(e^{t/4} y)' = 3 e^{t/4} + 2 e^{t/4} \cos 2t$. After integration, we get that the general solution is $y(t) = 12 + (8 \cos 2t + 64 \sin 2t)/65 + c e^{-t/4}$. Invoking the initial condition, $y(0) = 0$, the specific solution is $y(t) = 12 + (8 \cos 2t + 64 \sin 2t - 788 e^{-t/4})/65$. As $t \rightarrow \infty$, the exponential term will decay, and the solution will oscillate about

an average value of 12, with an amplitude of $8/\sqrt{65}$.

(b) Solving $y(t) = 12$, we obtain the desired value $t \approx 10.0658$.

31. The integrating factor is $\mu(t) = e^{-2t}$, and the differential equation can be written as $(e^{-2t} y)' = 3t e^{-2t} + 2 e^{-t}$. The general solution is $y(t) = -3t/2 - 3/4 - 2e^t + c e^{2t}$. Imposing the initial condition, $y(t) = -3t/2 - 3/4 - 2e^t + (y_0 + 11/4)e^{2t}$. Now as $t \rightarrow \infty$, the term containing e^{2t} will dominate the solution. Its sign will determine the divergence properties. Hence the critical value of the initial condition is $y_0 = -11/4$. The corresponding solution, $y(t) = -3t/2 - 3/4 - 2e^t$, will also decrease without bound.

Note on Problems 34-37:

Let $g(t)$ be given, and consider the function $y(t) = y_1(t) + g(t)$, in which $y_1(t) \rightarrow 0$ as $t \rightarrow \infty$. Differentiating, $y'(t) = y_1'(t) + g'(t)$. Letting a be a constant, it follows that $y'(t) + ay(t) = y_1'(t) + ay_1(t) + g'(t) + ag(t)$. Note that the hypothesis on the function $y_1(t)$ will be satisfied, if $y_1'(t) + ay_1(t) = 0$. That is, $y_1(t) = c e^{-at}$. Hence $y(t) = c e^{-at} + g(t)$, which is a solution of the equation $y' + ay = g'(t) + ag(t)$. For convenience, choose $a = 1$.

34. Here $g(t) = 5$, and we consider the linear equation $y' + y = 5$. The integrating factor is $\mu(t) = e^t$, and the differential equation can be written as $(e^t y)' = 5e^t$. The general solution is $y(t) = 5 + c e^{-t}$.

36. Here $g(t) = 3t + 1$. Consider the linear equation $y' + y = 3 + 3t + 1$. The integrating factor is $\mu(t) = e^t$, and the differential equation can be written as $(e^t y)' = (3t + 4)e^t$. The general solution is $y(t) = 3t + 1 + c e^{-t}$.

37. $g(t) = 1 - t^2$. Consider the linear equation $y' + y = 1 - 2t - t^2$. The integrating factor is $\mu(t) = e^t$, and the equation can be written as $(e^t y)' = (1 - 2t - t^2)e^t$. The general solution is $y(t) = 1 - t^2 + c e^{-t}$.

38.(a) Differentiating y and using the fundamental theorem of calculus we obtain that $y' = A e^{-\int p(t)dt} \cdot (-p(t))$, and then $y' + p(t)y = 0$.

(b) Differentiating y we obtain that

$$y' = A'(t)e^{-\int p(t)dt} + A(t)e^{-\int p(t)dt} \cdot (-p(t)).$$

If this satisfies the differential equation then

$$y' + p(t)y = A'(t)e^{-\int p(t)dt} = g(t)$$

and the required condition follows.

(c) Let us denote $\mu(t) = e^{\int p(t)dt}$. Then clearly $A(t) = \int \mu(t)g(t)dt$, and after substitution $y = \int \mu(t)g(t)dt \cdot (1/\mu(t))$, which is just Eq. (33).

40. We assume a solution of the form $y = A(t)e^{-\int(1/t)dt} = A(t)e^{-\ln t} = A(t)t^{-1}$, where $A(t)$ satisfies $A'(t) = 2t \sin 2t$. This implies that

$$A(t) = -t \cos 2t + \frac{\sin 2t}{2} + c$$

and the solution is

$$y = -\cos 2t + \frac{\sin 2t}{2t} + \frac{c}{t}.$$

41. First rewrite the differential equation as

$$y' + \frac{2}{t}y = \frac{2 \sin t}{t}.$$

Assume a solution of the form $y = A(t)e^{-\int(2/t)dt} = A(t)t^{-2}$, where $A(t)$ satisfies the ODE $A'(t) = 2t \sin t$. It follows that $A(t) = 2 \sin t - 2t \cos t + c$ and thus $y = (2 \sin t - 2t \cos t + c)/t^2$.

2.2

Problems 1 through 20 follow the pattern of the examples worked in this section. The first eight problems, however, do not have an initial condition, so the integration constant c cannot be found.

2. For $x \neq -1$, the differential equation may be written as $y dy = [3x^2/(1+x^3)] dx$. Integrating both sides, with respect to the appropriate variables, we obtain the relation $y^2/2 = \ln |1+x^3| + c$. That is, $y(x) = \pm \sqrt{2 \ln |1+x^3| + c}$.

3. The differential equation may be written as $y^{-2} dy = -\cos x dx$. Integrating both sides of the equation, with respect to the appropriate variables, we obtain the relation $-y^{-1} = -\sin x + c$. That is, $(c + \sin x)y = 1$, in which c is an arbitrary constant. Solving for the dependent variable, explicitly, $y(x) = 1/(c + \sin x)$.


5. Write the differential equation as $\cos^{-2} 4y dy = \cos^2 x dx$, which also can be written as $\sec^2 4y dy = \cos^2 x dx$. Integrating both sides of the equation, with respect to the appropriate variables, we obtain the relation $\tan 4y = 2 \sin x \cos x + 2x + c$.

7. The differential equation may be written as $(2y + e^y)dy = (3x - e^{-x})dx$. Integrating both sides of the equation, with respect to the appropriate variables, we obtain the relation $y^2 + e^y = 3x^2/2 + e^{-x} + c$.

8. Write the differential equation as $(2 + y^2)dy = x^3 dx$. Integrating both sides of the equation, we obtain the relation $2y + y^3/3 = x^4/4 + c$.

9.(a) The differential equation is separable, with $y^{-2} dy = (1 - 2x)dx$. Integration yields $-y^{-1} = x - x^2 + c$. Substituting $x = 0$ and $y = -1/12$, we find that $c = 12$. Hence the specific solution is $y = 1/(x^2 - x - 12)$.


(b)



(c) Note that $x^2 - x - 12 = (x + 3)(x - 4)$. Hence the solution becomes singular at $x = -3$ and $x = 4$, so the interval of existence is $(-3, 4)$.

11.(a) Rewrite the differential equation as $x e^x dx = -2y dy$. Integrating both sides of the equation results in $x e^x - e^x = -y^2 + c$. Invoking the initial condition, we obtain $c = -3/4$. Hence $y^2 = e^x - x e^x - 3/4$. The explicit form of the solution is $y(x) = \sqrt{e^x - x e^x - 3/4}$. The positive sign is chosen, since $y(0) = 1/2$.


(b)



(c) The function under the radical becomes negative near $x \approx -0.96$ and $x \approx 0.58$.

12.(a) Write the differential equation as $r^{-2} dr = \theta^{-1} d\theta$. Integrating both sides of the equation results in the relation $-r^{-1} = \ln \theta + c$. Imposing the condition $r(1) = 3$, we obtain $c = -1/3$. The explicit form of the solution is $r = 3/(1 - 3 \ln \theta)$.


(b)



(c) Clearly, the solution makes sense only if $\theta > 0$. Furthermore, the solution becomes singular when $\ln \theta = 1/3$, that is, $\theta = \sqrt[3]{e}$.

14.(a) Write the differential equation as $y^{-3}dy = x(1+x^2)^{-1/2}dx$. Integrating both sides of the equation, with respect to the appropriate variables, we obtain the relation $-y^{-2}/2 = \sqrt{1+x^2} + c$. Imposing the initial condition, we obtain $c = -3/2$. Hence the specific solution can be expressed as $y^{-2} = 3 - 2\sqrt{1+x^2}$. The explicit form of the solution is $y(x) = 1/\sqrt{3 - 2\sqrt{1+x^2}}$. The positive sign is chosen to satisfy the initial condition.


(b)



(c) The solution becomes singular when $2\sqrt{1+x^2} = 3$. That is, at $x = \pm\sqrt{5}/2$.


16.(a) Rewrite the differential equation as $4y^3dy = x(x^2+1)dx$. Integrating both sides of the equation results in $y^4 = (x^2+1)^2/4 + c$. Imposing the initial condition, we obtain $c = 0$. Hence the solution may be expressed as $(x^2+1)^2 - 4y^4 = 0$. The explicit form of the solution is $y(x) = -\sqrt{(x^2+1)/2}$. The sign is chosen based on $y(0) = -1/\sqrt{2}$.

(b)

(c) The solution is valid for all $x \in \mathbb{R}$.

18.(a) Write the differential equation as $(3 + 4y)dy = (e^{-x} - e^x)dx$. Integrating both sides of the equation, with respect to the appropriate variables, we obtain the relation $3y + 2y^2 = -(e^x + e^{-x}) + c$. Imposing the initial condition, $y(0) = 2$, we obtain $c = 16$. Thus, the solution can be expressed as $3y + 2y^2 = -(e^x + e^{-x}) + 16$. Now by completing the square on the left hand side, $2(y + 3/4)^2 = -(e^x + e^{-x}) + 137/8$. Hence the explicit form of the solution is $y(x) = -3/4 + \sqrt{137/16 - \cosh x}$.


(b)



(c) Note the $137 - 16 \cosh x \geq 0$ as long as $|x| > 2.8371$ (approximately). Hence the solution is valid on the interval $-2.8371 < x < 2.8371$.

20.(a) Rewrite the differential equation as $y^2 dy = \arcsin x / \sqrt{1 - x^2} dx$. Integrating both sides of the equation results in $y^3/3 = (\arcsin x)^2/2 + c$. Imposing the condition $y(0) = 1$, we obtain $c = 1/3$. The explicit form of the solution is $y(x) = (3(\arcsin x)^2/2 + 1)^{1/3}$.

(b)



(c) Since $\arcsin x$ is defined for $-1 \leq x \leq 1$, this is the interval of existence.

22. The differential equation can be written as $(3y^2 - 6)dy = 3x^2 dx$. Integrating both sides, we obtain $y^3 - 6y = x^3 + c$. Imposing the initial condition, the specific solution is $y^3 - 6y = x^3 - 1$. Referring back to the differential equation, we find that $y' \rightarrow \infty$ as $y \rightarrow \pm\sqrt{2}$. The respective values of the abscissas are $x \approx -1.67, 1.88$. Hence the solution is valid for $-1.67 < x < 1.88$.

24. Write the differential equation as $(3 + 2y)dy = (2 - e^x)dx$. Integrating both sides, we obtain $3y + y^2 = 2x - e^x + c$. Based on the specified initial condition, the solution can be written as $3y + y^2 = 2x - e^x + 1$. Completing the square, it follows that $y(x) = -3/2 + \sqrt{2x - e^x + 13/4}$. The solution is defined if $2x - e^x + 13/4 \geq 0$, that is, $-1.5 \leq x \leq 2$ (approximately). In that interval, $y' = 0$ for $x = \ln 2$. It can be verified that $y''(\ln 2) < 0$. In fact, $y''(x) < 0$ on the interval of definition. Hence the solution attains a global maximum at $x = \ln 2$.

26. The differential equation can be written as $(1 + y^2)^{-1}dy = 4(1 + x)dx$. Integrating both sides of the equation, we obtain $\arctan y = 4x + 2x^2 + c$. Imposing the given initial condition, the specific solution is $\arctan y = 4x + 2x^2$. Therefore, $y = \tan(4x + 2x^2)$. The solution is valid on the interval $-0.537 < x < 0.336$. Referring back to the differential equation, the solution is stationary at $x = -1$. This is not on the interval of existence, and there is no global minimum for the solution.

28.(a) Write the differential equation as $y^{-1}(4 - y)^{-1}dy = t(1 + t)^{-1}dt$. Integrating both sides of the equation, we obtain $\ln |y| - \ln |y - 4| = 4t - 4 \ln |1 + t| + c$. Taking the exponential of both sides $|y/(y - 4)| = ce^{4t}/(1 + t)^4$. It follows that as $t \rightarrow \infty$, $|y/(y - 4)| = |1 + 4/(y - 4)| \rightarrow \infty$. That is, $y(t) \rightarrow 4$.


(b) Setting $y(0) = 2$, we obtain that $c = 1$. Based on the initial condition, the solution may be expressed as $y/(y - 4) = -e^{4t}/(1 + t)^4$. Note that $y/(y - 4) < 0$, for all $t \geq 0$. Hence $y < 4$ for all $t \geq 0$. Referring back to the differential equation, it follows that y' is always positive. This means that the solution is monotone increasing. We find that the root of the equation $e^{4t}/(1 + t)^4 = 399$ is near $t = 2.844$.

(c) Note the $y(t) = 4$ is an equilibrium solution. Examining the local direction field we see that if $y(0) > 0$, then the corresponding solutions converge to $y = 4$. Referring back to part (a), we have $y/(y-4) = [y_0/(y_0-4)] e^{4t}/(1+t)^4$, for $y_0 \neq 4$. Setting $t = 2$, we obtain $y_0/(y_0-4) = (3/e^2)^4 y(2)/(y(2)-4)$. Now since the function $f(y) = y/(y-4)$ is monotone for $y < 4$ and $y > 4$, we need only solve the equations $y_0/(y_0-4) = -399(3/e^2)^4$ and $y_0/(y_0-4) = 401(3/e^2)^4$. The respective solutions are $y_0 = 3.6622$ and $y_0 = 4.4042$.

32.(a) Observe that $(2x^2 + 3y^2)/2xy = (y/x)^{-1} + (3/2)(y/x)$. Hence the differential equation is homogeneous.

(b) The substitution $y = xv$ results in $v + xv' = (2x^2 + 3x^2v^2)/2x^2v$. The transformed equation is $v' = (2 + v^2)/2xv$. This equation is separable, with general solution $v^2 + 2 = cx$. In terms of the original dependent variable, the solution is $2x^2 + y^2 = cx^3$.


(c) The integral curves are symmetric with respect to the origin.



34.(a) Observe that $-(4x + 3y)/(2x + y) = -2 - (y/x)[2 + (y/x)]^{-1}$. Hence the differential equation is homogeneous.

(b) The substitution $y = xv$ results in $v + xv' = -2 - v/(2 + v)$. The transformed equation is $v' = -(v^2 + 5v + 4)/(2 + v)x$. This equation is separable, with general solution $(v + 4)^2 |v + 1| = c/x^3$. In terms of the original dependent variable, the solution is $(4x + y)^2 |x + y| = c$.


(c) The integral curves are symmetric with respect to the origin.



36.(a) Divide by x^2 to see that the equation is homogeneous. Substituting $y = xv$, we obtain $xv' = 1 + 2v + 2v^2$. The resulting differential equation is separable.

(b) Write the equation as $dv/(1 + 2v + 2v^2) = x^{-1}dx$. Integrating both sides of the equation, we obtain the general solution $\arctan(2v + 1) = \ln|x| + c$. In terms of the original dependent variable, the solution is $\arctan((2y + x)/x) - \ln|x| = c$.


(c) The integral curves are symmetric with respect to the origin.



37.(a) The differential equation can be expressed as $y' = (1/2)(y/x)^{-1} - 2(y/x)$. Hence the equation is homogeneous. The substitution $y = xv$ results in $xv' = (1 - 6v^2)/2v$. Separating variables, we have $2v dv/(1 - 6v^2) = dx/x$.

(b) Integrating both sides of the transformed equation yields $-(\ln|1 - 6v^2|)/6 = \ln|x| + c$, that is, $1 - 6v^2 = c/|x|^6$. In terms of the original dependent variable, the general solution is $6y^2 = x^2 - c/|x|^4$.


(c) The integral curves are symmetric with respect to the origin.



38.(a) The differential equation can be expressed as $y' = (3/2)(y/x) - (y/x)^{-1}$. Hence the equation is homogeneous. The substitution $y = xv$ results in $xv' = (v^2 - 2)/2v$, that is, $2v dv/(v^2 - 2) = dx/x$.

(b) Integrating both sides of the transformed equation yields $\ln |v^2 - 2| = \ln |x| + c$, that is, $v^2 - 2 = c|x|$. In terms of the original dependent variable, the general solution is $y^2 = cx^2|x| + 2x^2$.

(c) The integral curves are symmetric with respect to the origin.



2.3

1. Let $Q(t)$ be the amount of dye in the tank at time t . Clearly, $Q(0) = 150$ g. The differential equation governing the amount of dye is $Q'(t) = -2Q(t)/150$. The solution of this separable equation is $Q(t) = Q(0)e^{-t/75} = 150e^{-t/75}$. We need the time T such that $Q(T) = 1.5$ g. This means we have to solve $1.5 = 150e^{-T/75}$ and we obtain that $T = -75 \ln(1/100) = 75 \ln 100 \approx 345.4$ min.


5.(a) Let Q be the amount of salt in the tank. Salt enters the tank of water at a rate of $2(1/4)(1 + (1/2)\sin t) = 1/2 + (1/4)\sin t$ dkg/min. It leaves the tank at a

rate of $2Q/100$ dkg/min. Hence the differential equation governing the amount of salt at any time is

$$\frac{dQ}{dt} = \frac{1}{2} + \frac{1}{4} \sin t - \frac{Q}{50}.$$

The initial amount of salt is $Q_0 = 50$ dkg. The governing differential equation is linear, with integrating factor $\mu(t) = e^{t/50}$. Write the equation as $(e^{t/50}Q)' = e^{t/50}(1/2 + (1/4) \sin t)$. The specific solution is $Q(t) = 25 + (12.5 \sin t - 625 \cos t + 63150 e^{-t/50})/2501$ dkg.

(b)



(c) The amount of salt approaches a steady state, which is an oscillation of approximate amplitude $1/4$ about a level of 25 dkg.

6.(a) Using the Principle of Conservation of Energy, the speed v of a particle falling from a height h is given by

$$\frac{1}{2}mv^2 = mgh.$$

(b) The outflow rate is (outflow cross-section area) \times (outflow velocity): $\alpha a \sqrt{2gh}$. At any instant, the volume of water in the tank is $V(h) = \int_0^h A(u) du$. The time rate of change of the volume is given by $dV/dt = (dV/dh)(dh/dt) = A(h)dh/dt$. Since the volume is decreasing, $dV/dt = -\alpha a \sqrt{2gh}$.

(c) With $A(h) = 4\pi$, $a = 0.01\pi$, $\alpha = 0.6$, the differential equation for the water level h is $4\pi(dh/dt) = -0.006\pi\sqrt{2gh}$, with solution $h(t) = 0.00000125gt^2 - 0.0015\sqrt{2gh(0)}t + h(0)$. Setting $h(0) = 4$ and $g = 9.8$, $h(t) = 0.000011025t^2 - 0.0132816t + 4$, resulting in $h(t) = 0$ for $t \approx 602.3$ s.

7.(a) The equation governing the value of the investment is $dS/dt = rS$. The value of the investment, at any time, is given by $S(t) = S_0 e^{rt}$. Setting $S(T) = 2S_0$, the required time is $T = \ln(2)/r$.

(b) For the case $r = .08$, $T \approx 8.66$ yr.

(c) Referring to part (a), $r = \ln(2)/T$. Setting $T = 8$, the required interest rate is to be approximately $r = 8.66\%$.

12.(a) Using Eq.(15) we have $dS/dt - 0.005S = -(800 + 10t)$, $S(0) = 150,000$. Using an integrating factor and integration by parts we obtain that $S(t) = 560,000 - 410,000e^{0.005t} + 2000t$. Setting $S(t) = 0$ and solving numerically for t yields $t = 146.54$ months.


(b) The solution we obtained in part (a) with a general initial condition $S(0) = S_0$ is $S(t) = 560,000 - 560,000e^{0.005t} + S_0e^{0.005t} + 2000t$. Solving the equation $S(240) = 0$ yields $S_0 = 246,758$.

13.(a) Let $Q' = -rQ$. The general solution is $Q(t) = Q_0e^{-rt}$. Based on the definition of half-life, consider the equation $Q_0/2 = Q_0e^{-5730r}$. It follows that $-5730r = \ln(1/2)$, that is, $r = 1.2097 \times 10^{-4}$ per year.

(b) The amount of carbon-14 is given by $Q(t) = Q_0e^{-1.2097 \times 10^{-4}t}$.

(c) Given that $Q(T) = Q_0/4$, we have the equation $1/4 = e^{-1.2097 \times 10^{-4}T}$. Solving for the decay time, the apparent age of the remains is approximately $T = 11,460$ years.

15.(a) The differential equation $dy/dt = r(t)y - k$ is linear, with integrating factor $\mu(t) = e^{-\int r(t)dt}$. Write the equation as $(\mu y)' = -k\mu(t)$. Integration of both sides yields the general solution $y = [-k \int \mu(\tau)d\tau + y_0\mu(0)]/\mu(t)$. In this problem, the integrating factor is $\mu(t) = e^{(\cos t - t)/5}$.



(b) The population becomes extinct, if $y(t^*) = 0$, for some $t = t^*$. Referring to part (a), we find that $y(t^*) = 0$ when

$$\int_0^{t^*} e^{(\cos \tau - \tau)/5} d\tau = 5e^{1/5}y_c.$$

It can be shown that the integral on the left hand side increases monotonically, from zero to a limiting value of approximately 5.0893. Hence extinction can happen only if $5e^{1/5}y_0 < 5.0893$. Solving $5e^{1/5}y_c = 5.0893$ yields $y_c = 0.8333$.

(c) Repeating the argument in part (b), it follows that $y(t^*) = 0$ when


$$\int_0^{t^*} e^{(\cos \tau - \tau)/5} d\tau = \frac{1}{k} e^{1/5} y_c.$$

Hence extinction can happen only if $e^{1/5} y_0/k < 5.0893$, so $y_c = 4.1667 k$.

(d) Evidently, y_c is a linear function of the parameter k .

17.(a) The solution of the governing equation satisfies $u^3 = u_0^3/(3\alpha u_0^3 t + 1)$. With the given data, it follows that $u(t) = 2000/\sqrt[3]{6t/125 + 1}$.

(b)




(c) Numerical evaluation results in $u(t) = 600$ for $t \approx 750.77$ s.

22.(a) The differential equation for the upward motion is $mdv/dt = -\mu v^2 - mg$, in which $\mu = 1/1325$. This equation is separable, with $m/(\mu v^2 + mg) dv = -dt$. Integrating both sides and invoking the initial condition, $v(t) = 44.133 \tan(0.425 - 0.222t)$. Setting $v(t_1) = 0$, the ball reaches the maximum height at $t_1 = 1.916$ s. Integrating $v(t)$, the position is given by $x(t) = 198.75 \ln[\cos(0.222t - 0.425)] + 48.57$. Therefore the maximum height is $x(t_1) = 48.56$ m.

(b) The differential equation for the downward motion is $mdv/dt = +\mu v^2 - mg$. This equation is also separable, with $m/(mg - \mu v^2) dv = -dt$. For convenience, set $t = 0$ at the top of the trajectory. The new initial condition becomes $v(0) = 0$. Integrating both sides and invoking the initial condition, we obtain $\ln((44.13 - v)/(44.13 + v)) = t/2.25$. Solving for the velocity, $v(t) = 44.13(1 - e^{t/2.25})/(1 + e^{t/2.25})$. Integrating $v(t)$, we obtain $x(t) = 99.29 \ln(e^{t/2.25}/(1 + e^{t/2.25})^2) + 186.2$. To estimate the duration of the downward motion, set $x(t_2) = 0$, resulting in $t_2 = 3.276$ s. Hence the total time that the ball spends in the air is $t_1 + t_2 = 5.192$ s.

(c)



24.(a) Setting $-\mu v^2 = v(dv/dx)$, we obtain $dv/dx = -\mu v$.

(b) The speed v of the sled satisfies $\ln(v/v_0) = -\mu x$. Noting that the unit conversion factors cancel, solution of $\ln(24/240) = -600\mu$ results in $\mu = \ln(10)/600 \text{ m}^{-1} \approx 0.0038376 \text{ m}^{-1} \approx 3.8376 \text{ km}^{-1}$.

(c) Solution of $dv/dt = -\mu v^2$ can be expressed as $1/v - 1/v_0 = \mu t$. The elapsed time is

$$t = (1/24 - 1/240)/(\ln(10)/600) \cdot 3600 \approx 35.18 \text{ s}.$$

25.(a) Measure the positive direction of motion upward. The equation of motion is given by $mdv/dt = -kv - mg$. The initial value problem is $dv/dt = -kv/m - g$, with $v(0) = v_0$. The solution is $v(t) = -mg/k + (v_0 + mg/k)e^{-kt/m}$. Setting $v(t_m) = 0$, the maximum height is reached at time $t_m = (m/k) \ln[(mg + kv_0)/mg]$. Integrating the velocity, the position of the body is

$$x(t) = -mgt/k + \left[\left(\frac{m}{k}\right)^2 g + \frac{mv_0}{k} \right] (1 - e^{-kt/m}).$$

Hence the maximum height reached is

$$x_m = x(t_m) = \frac{mv_0}{k} - g\left(\frac{m}{k}\right)^2 \ln \left[\frac{mg + kv_0}{mg} \right].$$

(b) Recall that for $\delta \ll 1$, $\ln(1 + \delta) = \delta - \delta^2/2 + \delta^3/3 - \delta^4/4 + \dots$


(c) The dimensions of the quantities involved are $[k] = MT^{-1}$, $[v_0] = LT^{-1}$, $[m] = M$ and $[g] = LT^{-2}$. This implies that kv_0/mg is dimensionless.

31.(a) Both equations are linear and separable. Initial conditions: $v(0) = u \cos A$ and $w(0) = u \sin A$. We obtain the solutions $v(t) = (u \cos A)e^{-rt}$ and $w(t) = -g/r + (u \sin A + g/r)e^{-rt}$.

(b) Integrating the solutions in part (a), and invoking the initial conditions, the coordinates are $x(t) = u \cos A(1 - e^{-rt})/r$ and

$$y(t) = -\frac{gt}{r} + \frac{g + ur \sin A + hr^2}{r^2} - \left(\frac{u}{r} \sin A + \frac{g}{r^2}\right)e^{-rt}.$$

(c)



(d) Let T be the time that it takes the ball to go 350 ft horizontally. Then from above, $e^{-T/5} = (u \cos A - 70)/u \cos A$. At the same time, the height of the ball is given by

$$y(T) = -160T + 803 + 5u \sin A - \frac{(800 + 5u \sin A)(u \cos A - 70)}{u \cos A}.$$

Hence A and u must satisfy the equality

$$800 \ln \left[\frac{u \cos A - 70}{u \cos A} \right] + 803 + 5u \sin A - \frac{(800 + 5u \sin A)(u \cos A - 70)}{u \cos A} = 10$$

for the ball to touch the top of the wall. To find the optimal values for u and A , consider u as a function of A and use implicit differentiation in the above equation to find that

$$\frac{du}{dA} = -\frac{u(u^2 \cos A - 70u - 11200 \sin A)}{11200 \cos A}.$$

Solving this equation simultaneously with the above equation yields optimal values for u and A : $u \approx 145.3$ ft/s, $A \approx 0.644$ rad.

32.(a) Solving equation (i), $y'(x) = [(k^2 - y)/y]^{1/2}$. The positive answer is chosen, since y is an increasing function of x .

(b) Let $y = k^2 \sin^2 t$. Then $dy = 2k^2 \sin t \cos t dt$. Substituting into the equation in part (a), we find that

$$\frac{2k^2 \sin t \cos t dt}{dx} = \frac{\cos t}{\sin t}.$$

Hence $2k^2 \sin^2 t dt = dx$.

(c) Setting $\theta = 2t$, we further obtain $k^2 \sin^2(\theta/2) d\theta = dx$. Integrating both sides of the equation and noting that $t = \theta = 0$ corresponds to the origin, we obtain the solutions $x(\theta) = k^2(\theta - \sin \theta)/2$ and (from part (b)) $y(\theta) = k^2(1 - \cos \theta)/2$.

(d) Note that $y/x = (1 - \cos \theta)/(\theta - \sin \theta)$. Setting $x = 1$, $y = 2$, the solution of the equation $(1 - \cos \theta)/(\theta - \sin \theta) = 2$ is $\theta \approx 1.401$. Substitution into either of the expressions yields $k \approx 2.193$.

2.4

2. Rewrite the differential equation as $y' + 1/(t(t-5))y = 0$. It is evident that the coefficient $1/(t(t-5))$ is continuous everywhere except at $t = 0, 5$. Since the initial condition is specified at $t = 3$, Theorem 2.4.1 assures the existence of a unique solution on the interval $0 < t < 5$.

3. The function $\tan t$ is discontinuous at odd multiples of $\pi/2$. Since $3\pi/2 < 2\pi < 5\pi/2$, the initial value problem has a unique solution on the interval $(3\pi/2, 5\pi/2)$.

5. $p(t) = 2t/(16 - t^2)$ and $g(t) = 3t^2/(16 - t^2)$. These functions are discontinuous at $x = \pm 4$. The initial value problem has a unique solution on the interval $(-4, 4)$.

6. The function $\ln t$ is defined and continuous on the interval $(0, \infty)$. At $t = 1$, $\ln t = 0$, so the normal form of the differential equation has a singularity there. Also, $\cot t$ is not defined at integer multiples of π , so the initial value problem will have a solution on the interval $(1, \pi)$.

7. The function $f(t, y)$ is continuous everywhere on the plane, except along the straight line $y = -2t/5$. The partial derivative $\partial f/\partial y = -16t/(2t + 5y)^2$ has the same region of continuity.

9. The function $f(t, y)$ is discontinuous along the coordinate axes, and on the hyperbola $t^2 - y^2 = 1$. Furthermore,

$$\frac{\partial f}{\partial y} = \frac{\pm 1}{y(1 - t^2 + y^2)} - 2 \frac{y \ln |ty|}{(1 - t^2 + y^2)^2}$$

has the same points of discontinuity.

10. $f(t, y)$ is continuous everywhere on the plane. The partial derivative $\partial f/\partial y$ is also continuous everywhere.

12. The function $f(t, y)$ is discontinuous along the lines $t = \pm k\pi$ for $k = 0, 1, 2, \dots$ and $y = -1$. The partial derivative $\partial f/\partial y = \cot t(2y + y^2)/(1 + y)^2$ has the same region of continuity.


14. The equation is separable, with $dy/y^2 = 4t dt$. Integrating both sides, the solution is given by $y(t) = y_0/(1 - y_0 2t^2)$. For $y_0 > 0$, solutions exist as long as $t^2 < 1/2y_0$. For $y_0 \leq 0$, solutions are defined for all t .

15. The equation is separable, with $dy/y^5 = -dt$. Integrating both sides and invoking the initial condition, $y(t) = y_0/\sqrt[4]{4y_0^4 t + 1}$. Solutions exist as long as

$4y_0^4 t + 1 > 0$, that is, $4y_0^4 t > -1$. If $y_0 \neq 0$, solutions exist for $t > -1/4y_0^4$. If $y_0 = 0$, then the solution $y(t) = 0$ exists for all t .


16. The function $f(t, y)$ is discontinuous along the straight lines $t = -1$ and $y = 0$. The partial derivative $\partial f / \partial y$ is discontinuous along the same lines. The equation is separable, with $y dy = t^2 dt / (1 + t^3)$. Integrating and invoking the initial condition, the solution is $y(t) = [(2/3) \ln |1 + t^3| + y_0^2]^{1/2}$. Solutions exist as long as $(2/3) \ln |1 + t^3| + y_0^2 \geq 0$, that is, $y_0^2 \geq -(2/3) \ln |1 + t^3|$. For all y_0 (it can be verified that $y_0 = 0$ yields a valid solution, even though Theorem 2.4.2 does not guarantee one), solutions exist as long as $|1 + t^3| \geq e^{-3y_0^2/2}$. From above, we must have $t > -1$. Hence the inequality may be written as $t^3 \geq e^{-3y_0^2/2} - 1$. It follows that the solutions are valid for $(e^{-3y_0^2/2} - 1)^{1/3} < t < \infty$.

18.




Based on the direction field, and the differential equation, for $y_0 < 0$, the slopes eventually become negative, and hence solutions tend to $-\infty$. For $y_0 > 0$, solutions increase without bound if $t_0 < 0$. Otherwise, the slopes eventually become negative, and solutions tend to zero. Furthermore, $y_0 = 0$ is an equilibrium solution. Note that slopes are zero along the curves $y = 0$ and $ty = 4$.

19.



For initial conditions (t_0, y_0) satisfying $ty < 4$, the respective solutions all tend to zero. For $y_0 \leq 16$, the solutions tend to 0; for $y_0 > 16$, the solutions tend to ∞ . Also, $y_0 = 0$ is an equilibrium solution.

20.



Solutions with $t_0 < 0$ all tend to $-\infty$. Solutions with initial conditions (t_0, y_0) to the right of the parabola $t = 1 + y^2$ asymptotically approach the parabola as $t \rightarrow \infty$. Integral curves with initial conditions above the parabola (and $y_0 > 0$) also approach the curve. The slopes for solutions with initial conditions below the parabola (and $y_0 < 0$) are all negative. These solutions tend to $-\infty$.

21.(a) No. There is no value of $t_0 \geq 0$ for which $(2/3)(t - t_0)^{2/3}$ satisfies the condition $y(1) = 1$.

(b) Yes. Let $t_0 = 1/2$ in Eq.(19).

(c) For $t_0 > 0$, $|y(2)| \leq (4/3)^{3/2} \approx 1.54$.

24. The assumption is $\phi'(t) + p(t)\phi(t) = 0$. But then $c\phi'(t) + p(t)c\phi(t) = 0$ as well.

26.(a) Recalling Eq.(33) in Section 2.1,

$$y = \frac{1}{\mu(t)} \int_{t_0}^t \mu(s)g(s) ds + \frac{c}{\mu(t)}.$$

It is evident that $y_1(t) = 1/\mu(t)$ and $y_2(t) = (1/\mu(t)) \int_{t_0}^t \mu(s)g(s) ds$.

(b) By definition, $1/\mu(t) = e^{-\int p(t)dt}$. Hence $y_1' = -p(t)/\mu(t) = -p(t)y_1$. That is, $y_1' + p(t)y_1 = 0$.

(c) $y_2' = (-p(t)/\mu(t)) \int_{t_0}^t \mu(s)g(s) ds + \mu(t)g(t)/\mu(t) = -p(t)y_2 + g(t)$. This implies that $y_2' + p(t)y_2 = g(t)$.

30. Since $n = 3$, set $v = y^{-2}$. It follows that $v' = -2y^{-3}y'$ and $y' = -(y^3/2)v'$. Substitution into the differential equation yields $-(y^3/2)v' - \varepsilon y = -\sigma y^3$, which further results in $v' + 2\varepsilon v = 2\sigma$. The latter differential equation is linear, and can be written as $(ve^{2\varepsilon t})' = 2\sigma e^{2\varepsilon t}$. The solution is given by $v(t) = \sigma/\varepsilon + ce^{-2\varepsilon t}$. Converting back to the original dependent variable, $y = \pm v^{-1/2} = \pm(\sigma/\varepsilon + ce^{-2\varepsilon t})^{-1/2}$.

31. Since $n = 3$, set $v = y^{-2}$. It follows that $v' = -2y^{-3}y'$ and $y' = -(y^3/2)v'$. The differential equation is written as $-(y^3/2)v' - (\Gamma \cos t + T)y = \sigma y^3$, which upon

further substitution is $v' + 2(\Gamma \cos t + T)v = 2$. This ODE is linear, with integrating factor $\mu(t) = e^{2\int(\Gamma \cos t + T)dt} = e^{2\Gamma \sin t + 2Tt}$. The solution is

$$v(t) = 2e^{-(2\Gamma \sin t + 2Tt)} \int_0^t e^{2\Gamma \sin \tau + 2T\tau} d\tau + ce^{-(2\Gamma \sin t + 2Tt)}.$$

Converting back to the original dependent variable, $y = \pm v^{-1/2}$.

33. The solution of the initial value problem $y_1' + 4y_1 = 0$, $y_1(0) = 1$ is $y_1(t) = e^{-4t}$. Therefore $y(1^-) = y_1(1) = e^{-4}$. On the interval $(1, \infty)$, the differential equation is $y_2' + y_2 = 0$, with $y_2(t) = ce^{-t}$. Therefore $y(1^+) = y_2(1) = ce^{-1}$. Equating the limits $y(1^-) = y(1^+)$, we require that $c = e^{-3}$. Hence the global solution of the initial value problem is


$$y(t) = \begin{cases} e^{-4t}, & 0 \leq t \leq 1 \\ e^{-3-t}, & t > 1 \end{cases}.$$


Note the discontinuity of the derivative

$$y'(t) = \begin{cases} -4e^{-4t}, & 0 < t < 1 \\ -e^{-3-t}, & t > 1 \end{cases}.$$

2.5



1.







For $y_0 \geq 0$, the only equilibrium point is $y^* = 0$, and $y' = ay + by^2 > 0$ when $y > 0$, hence the equilibrium solution $y = 0$ is unstable.

2.




The equilibrium points are $y^* = -a/b$ and $y^* = 0$, and $y' > 0$ when $y > 0$ or $y < -a/b$, and $y' < 0$ when $-a/b < y < 0$, therefore the equilibrium solution $y = -a/b$ is asymptotically stable and the equilibrium solution $y = 0$ is unstable.


4.



The only equilibrium point is $y^* = 0$, and $y' > 0$ when $y > 0$, $y' < 0$ when $y < 0$, hence the equilibrium solution $y = 0$ is unstable.


6.







The only equilibrium point is $y^* = 0$, and $y' > 0$ when $y < 0$, $y' < 0$ when $y > 0$, hence the equilibrium solution $y = 0$ is asymptotically stable.

8.




The only equilibrium point is $y^* = 2$, and $y' < 0$ for $y \neq 2$. As long as $y_0 \neq 2$, the corresponding solution is monotone decreasing. Hence the equilibrium solution $y = 2$ is semistable.


10.



The equilibrium points are $y^* = 0, \pm 2$, and $y' > 0$ for $y < -2$ or $0 < y < 2$ and $y' < 0$ for $-2 < y < 0$ or $y > 2$. The equilibrium solution $y = 0$ is unstable, and the remaining two are asymptotically stable.



12.





The equilibrium points are $y^* = 0, \pm 3$, and $y' < 0$ when $y < -3$ or $y > 3$, and $y' > 0$ for $-3 < y < 0$ or $0 < y < 3$. The equilibrium solutions $y = -3$ and $y = 3$ are unstable and asymptotically stable, respectively. The equilibrium solution $y = 0$ is semistable.

13.



The equilibrium points are $y^* = 0, 3$. $y' > 0$ for all y except $y = 0$ and $y = 3$. Both equilibrium solutions are semistable.

15.(a) Inverting Eq.(11), Eq.(13) shows t as a function of the population y and the carrying capacity K . With $y_0 = K/3$,

$$t = -\frac{1}{r} \ln \left| \frac{(1/3) [1 - (y/K)]}{(y/K) [1 - (1/3)]} \right|.$$

Setting $y = 2y_0$,

$$\tau = -\frac{1}{r} \ln \left| \frac{(1/3)[1 - (2/3)]}{(2/3)[1 - (1/3)]} \right|.$$

That is, $\tau = (\ln 4)/r$. If $r = 0.03$ per year, $\tau \approx 46.21$ years.

(b) In Eq.(13), set $y_0/K = \alpha$ and $y/K = \beta$. As a result, we obtain

$$T = -\frac{1}{r} \ln \left| \frac{\alpha[1 - \beta]}{\beta[1 - \alpha]} \right|.$$

Given $\alpha = 0.1$, $\beta = 0.9$ and $r = 0.03$ per year, $\tau \approx 146.48$ years.

19.(a) The rate of increase of the volume is given by rate of flow in – rate of flow out. That is, $dV/dt = k - \alpha a \sqrt{2gh}$. Since the cross section is constant, $dV/dt = Adh/dt$. Hence the governing equation is $dh/dt = (k - \alpha a \sqrt{2gh})/A$.

(b) Setting $dh/dt = 0$, the equilibrium height is $h_e = (1/2g)(k/\alpha a)^2$. Furthermore, since $dh/dt < 0$ for $h > h_e$ and $dh/dt > 0$ for $h < h_e$, it follows that the equilibrium height is asymptotically stable.

22.(a) The equilibrium points are at $y^* = 0$ and $y^* = 1$. Since $f'(y) = \alpha - 2\alpha y$, the equilibrium solution $y = 0$ is unstable and the equilibrium solution $y = 1$ is asymptotically stable.

(b) The differential equation is separable, with $[y(1 - y)]^{-1} dy = \alpha dt$. Integrating both sides and invoking the initial condition, the solution is

$$y(t) = \frac{y_0 e^{\alpha t}}{1 - y_0 + y_0 e^{\alpha t}} = \frac{y_0}{y_0 + (1 - y_0)e^{-\alpha t}}.$$

It is evident that (independent of y_0) $\lim_{t \rightarrow -\infty} y(t) = 0$ and $\lim_{t \rightarrow \infty} y(t) = 1$.


23.(a) $y(t) = y_0 e^{-\beta t}$.

(b) From part (a), $dx/dt = -\alpha x y_0 e^{-\beta t}$. Separating variables, $dx/x = -\alpha y_0 e^{-\beta t} dt$. Integrating both sides, the solution is $x(t) = x_0 e^{-\alpha y_0 (1 - e^{-\beta t})/\beta}$.


(c) As $t \rightarrow \infty$, $y(t) \rightarrow 0$ and $x(t) \rightarrow x_0 e^{-\alpha y_0/\beta}$. Over a long period of time, the proportion of carriers vanishes. Therefore the proportion of the population that escapes the epidemic is the proportion of susceptibles left at that time, $x_0 e^{-\alpha y_0/\beta}$.

26.(a) For $a < 0$, the only critical point is at $y = 0$, which is asymptotically stable. For $a = 0$, the only critical point is at $y = 0$, which is asymptotically stable. For $a > 0$, the three critical points are at $y = 0, \pm\sqrt{a}$. The critical point at $y = 0$ is unstable, whereas the other two are asymptotically stable.

(b) Below, we graph solutions in the case $a = -1$, $a = 0$ and $a = 1$ respectively.






(c)




27.(a) $f(y) = y(a - y)$; $f'(y) = a - 2y$. For $a < 0$, the critical points are at $y = a$ and $y = 0$. Observe that $f'(a) > 0$ and $f'(0) < 0$. Hence $y = a$ is unstable and $y = 0$ asymptotically stable. For $a = 0$, the only critical point is at $y = 0$, which is semistable since $f(y) = -y^2$ is concave down. For $a > 0$, the critical points are at $y = 0$ and $y = a$. Observe that $f'(0) > 0$ and $f'(a) < 0$. Hence $y = 0$ is unstable and $y = a$ asymptotically stable.

(b) Below, we graph solutions in the case $a = -1$, $a = 0$ and $a = 1$ respectively.





(c)



2.6

1. $M(x, y) = 4x + 3$ and $N(x, y) = 6y - 1$. Since $M_y = N_x = 0$, the equation is exact. Integrating M with respect to x , while holding y constant, yields $\psi(x, y) = 2x^2 + 3x + h(y)$. Now $\psi_y = h'(y)$, and equating with N results in the possible function $h(y) = 3y^2 - y$. Hence $\psi(x, y) = 2x^2 + 3x + 3y^2 - y$, and the solution is defined implicitly as $2x^2 + 3x + 3y^2 - y = c$.

2. $M(x, y) = 3x - y$ and $N(x, y) = x - 3y$. Note that $M_y \neq N_x$, and hence the differential equation is not exact.

4. First divide both sides by $(4xy + 4)$. We now have $M(x, y) = y$ and $N(x, y) = x$. Since $M_y = N_x = 0$, the resulting equation is exact. Integrating M with respect to x , while holding y constant, results in $\psi(x, y) = xy + h(y)$. Differentiating with respect to y , $\psi_y = x + h'(y)$. Setting $\psi_y = N$, we find that $h'(y) = 0$, and hence $h(y) = 0$ is acceptable. Therefore the solution is defined implicitly as $xy = c$. Note that if $xy + 1 = 0$, the equation is trivially satisfied.

6. Write the equation as $(ax - by)dx + (bx - cy)dy = 0$. Now $M(x, y) = ax - by$ and $N(x, y) = bx - cy$. Since $M_y \neq N_x$, the differential equation is not exact.

8. $M(x, y) = e^x \sin y + 2y$ and $N(x, y) = -2x + e^x \sin y$. Note that $M_y \neq N_x$, and hence the differential equation is not exact.

10. $M(x, y) = y/x + 4x$ and $N(x, y) = \ln x - 3$. Since $M_y = N_x = 1/x$, the given equation is exact. Integrating N with respect to y , while holding x constant, results in $\psi(x, y) = y \ln x - 3y + h(x)$. Differentiating with respect to x , $\psi_x = y/x + h'(x)$. Setting $\psi_x = M$, we find that $h'(x) = 4x$, and hence $h(x) = 2x^2$. Therefore the solution is defined implicitly as $2x^2 + y \ln x - 3y = c$.

11. $M(x, y) = x \ln y + xy$ and $N(x, y) = y \ln x + xy$. Note that $M_y \neq N_x$, and hence the differential equation is not exact.

13. $M(x, y) = 2x - y$ and $N(x, y) = 2y - x$. Since $M_y = N_x = -1$, the equation is exact. Integrating M with respect to x , while holding y constant, yields $\psi(x, y) = x^2 - xy + h(y)$. Now $\psi_y = -x + h'(y)$. Equating ψ_y with N results in $h'(y) = 2y$, and hence $h(y) = y^2$. Thus $\psi(x, y) = x^2 - xy + y^2$, and the solution is given implicitly as $x^2 - xy + y^2 = c$. Invoking the initial condition $y(1) = 4$, the specific solution is $x^2 - xy + y^2 = 13$. The explicit form of the solution is $y(x) = (x + \sqrt{52 - 3x^2})/2$. Hence the solution is valid as long as $3x^2 \leq 52$.

16. $M(x, y) = y e^{2xy} + 3x$ and $N(x, y) = b x e^{2xy}$. Note that $M_y = e^{2xy} + 2xy e^{2xy}$, and $N_x = b e^{2xy} + 2bxy e^{2xy}$. The given equation is exact, as long as $b = 1$. Integrating N with respect to y , while holding x constant, results in $\psi(x, y) = e^{2xy}/2 + h(x)$. Now differentiating with respect to x , $\psi_x = y e^{2xy} + h'(x)$. Setting $\psi_x = M$, we find that $h'(x) = 3x$, and hence $h(x) = 3x^2/2$. We conclude that $\psi(x, y) = e^{2xy}/2 + 3x^2/2$. Hence the solution is given implicitly as $e^{2xy} + 3x^2 = c$.

17. Note that ψ is of the form $\psi(x, y) = f(x) + g(y)$, since each of the integrands is a function of a single variable. It follows that $\psi_x = f'(x)$ and $\psi_y = g'(y)$. That is, $\psi_x = M(x, y_0)$ and $\psi_y = N(x_0, y)$. Furthermore,

$$\frac{\partial^2 \psi}{\partial x \partial y}(x_0, y_0) = \frac{\partial M}{\partial y}(x_0, y_0) \quad \text{and} \quad \frac{\partial^2 \psi}{\partial y \partial x}(x_0, y_0) = \frac{\partial N}{\partial x}(x_0, y_0),$$

based on the hypothesis and the fact that the point (x_0, y_0) is arbitrary, $\psi_{xy} = \psi_{yx}$ and $M_y(x, y) = N_x(x, y)$.

18. Observe that $(M(x))_y = (N(y))_x = 0$.

20. $M_y = y^{-1} \cos y - y^{-2} \sin y$ and $N_x = -3e^{-x}(\cos x + \sin x)/y$. Multiplying both sides by the integrating factor $\mu(x, y) = y e^x$, the given equation can be written as $(e^x \sin y - 3y \sin x)dx + (e^x \cos y + 3 \cos x)dy = 0$. Let $\tilde{M} = \mu M$ and $\tilde{N} = \mu N$. Observe that $\tilde{M}_y = \tilde{N}_x$, and hence the latter ODE is exact. Integrating \tilde{N} with respect to y , while holding x constant, results in $\psi(x, y) = e^x \sin y + 3y \cos x + h(x)$. Now differentiating with respect to x , $\psi_x = e^x \sin y - 3y \sin x + h'(x)$. Setting $\psi_x = \tilde{M}$, we find that $h'(x) = 0$, and hence $h(x) = 0$ is feasible. Hence the solution of the given equation is defined implicitly by $e^x \sin y + 3y \cos x = c$.

21. $M_y = 1$ and $N_x = 2$. Multiply both sides by the integrating factor $\mu(x, y) = y$ to obtain $y^2 dx + (2xy - 3y^2 e^y) dy = 0$. Let $\tilde{M} = yM$ and $\tilde{N} = yN$. It is easy to see that $\tilde{M}_y = \tilde{N}_x$, and hence the latter ODE is exact. Integrating \tilde{M} with respect to x yields $\psi(x, y) = xy^2 + h(y)$. Equating ψ_y with \tilde{N} results in $h'(y) = -3y^2 e^y$, and hence $h(y) = -3e^y(y^2 - 2y + 2)$. Thus $\psi(x, y) = xy^2 - 3e^y(y^2 - 2y + 2)$, and the solution is defined implicitly by $xy^2 - 3e^y(y^2 - 2y + 2) = c$.

24. The equation $\mu M + \mu N y' = 0$ has an integrating factor if $(\mu M)_y = (\mu N)_x$, that is, $\mu_y M - \mu_x N = \mu N_x - \mu M_y$. Suppose that $N_x - M_y = R(xM - yN)$, in which R is some function depending only on the quantity $z = xy$. It follows that the modified form of the equation is exact, if $\mu_y M - \mu_x N = \mu R(xM - yN) = R(\mu x M - \mu y N)$. This relation is satisfied if $\mu_y = (\mu x)R$ and $\mu_x = (\mu y)R$. Now consider $\mu = \mu(xy)$. Then the partial derivatives are $\mu_x = \mu' y$ and $\mu_y = \mu' x$. Note that $\mu' = d\mu/dz$. Thus μ must satisfy $\mu'(z) = R(z)$. The latter equation is separable, with $d\mu = R(z)dz$, and $\mu(z) = \int R(z)dz$. Therefore, given $R = R(xy)$, it is possible to determine $\mu = \mu(xy)$ which becomes an integrating factor of the differential equation.

28. The equation is not exact, since $N_x - M_y = 2y - 1$. However, $(N_x - M_y)/M = (2y - 1)/y$ is a function of y alone. Hence there exists $\mu = \mu(y)$, which is a solution of the differential equation $\mu' = (2 - 1/y)\mu$. The latter equation is separable, with $d\mu/\mu = 2 - 1/y$. One solution is $\mu(y) = e^{2y - \ln y} = e^{2y}/y$. Now rewrite the given ODE as $e^{2y} dx + (2x e^{2y} - 1) dy = 0$. This equation is exact, and it is easy to see that $\psi(x, y) = x e^{2y} - y$. Therefore the solution of the given equation is defined implicitly by $x e^{2y} - y = c$.

30. The given equation is not exact, since $N_x - M_y = 8x^3/y^3 + 6/y^2$. But note that $(N_x - M_y)/M = 2/y$ is a function of y alone, and hence there is an integrating factor $\mu = \mu(y)$. Solving the equation $\mu' = (2/y)\mu$, an integrating factor is $\mu(y) = y^2$. Now rewrite the differential equation as $(4x^3 + 3y)dx + (3x + 2y^3)dy = 0$. By inspection, $\psi(x, y) = x^4 + 3xy + y^4/2$, and the solution of the given equation is defined implicitly by $x^4 + 3xy + y^4/2 = c$.

32. Multiplying both sides of the ODE by $\mu = [xy(2x + y)]^{-1}$, the given equation is equivalent to $[(3x + y)/(2x^2 + xy)] dx + [(x + y)/(2xy + y^2)] dy = 0$. Rewrite the differential equation as

$$\left[\frac{2}{x} + \frac{2}{2x + y} \right] dx + \left[\frac{1}{y} + \frac{1}{2x + y} \right] dy = 0.$$

It is easy to see that $M_y = N_x$. Integrating M with respect to x , while keeping y constant, results in $\psi(x, y) = 2 \ln |x| + \ln |2x + y| + h(y)$. Now taking the partial derivative with respect to y , $\psi_y = (2x + y)^{-1} + h'(y)$. Setting $\psi_y = N$, we find that $h'(y) = 1/y$, and hence $h(y) = \ln |y|$. Therefore $\psi(x, y) = 2 \ln |x| + \ln |2x + y| + \ln |y|$, and the solution of the given equation is defined implicitly by $2x^3 y + x^2 y^2 = c$.

2.7

2. The Euler formula is given by $y_{n+1} = y_n + h(3y_n - 1) = (1 + 3h)y_n - h$.


(a) 1.2, 1.46, 1.798, 2.2374

(b) 1.215, 1.49934, 1.87537, 2.37268

(c) 1.22365, 1.52232, 1.92119, 2.45386


(d) The differential equation is linear with solution $y(t) = (1 + 2e^{3t})/3$. The values are 1.23324, 1.54808, 1.97307, 2.54674.

5.




All solutions seem to converge to $y = 25/4$.

7.




All solutions seem to converge to a specific function.

8.



Solutions with initial conditions $|y(0)| > 2.5$ seem to diverge. On the other hand, solutions with initial conditions $|y(0)| < 2.5$ seem to converge to zero. Also, $y = 0$ is an equilibrium solution.

10.



Solutions with positive initial conditions increase without bound. Solutions with negative initial conditions decrease without bound. Note that $y = 0$ is an equilibrium solution.

11. The Euler formula is $y_{n+1} = y_n - 2h\sqrt{y_n} + 5h$. The initial value is $y_0 = 2$.

(a) 2.95148, 3.65042, 4.18248, 4.59561, 4.92048, 5.17816

(b) 2.93808, 3.63167, 4.16190, 4.57499, 4.90074, 5.15979

(c) 2.93159, 3.62253, 4.15183, 4.56486, 4.89102, 5.15072

(d) 2.92776, 3.73397, 4.14586, 4.55884, 4.88523, 5.14531

12. The Euler formula is $y_{n+1} = (1 + 4h)y_n - ht_n y_n^2$. The initial value is $(t_0, y_0) = (0, 0.5)$.

(a) 2.42831, 4.49378, 3.13540, 2.26560, 1.76850, 1.44951

(b) 2.65748, 4.46004, 3.14322, 2.27392, 1.77288, 1.45195

(c) 2.78710, 4.44811, 3.14607, 2.27804, 1.77516, 1.45322

(d) 2.86988, 4.44232, 3.14756, 2.28049, 1.77655, 1.45399

14. The Euler formula is $y_{n+1} = (1 - ht_n)y_n + hy_n^3/10$, with $(t_0, y_0) = (0, 1)$.

(a) 0.950517, 0.687550, 0.369188, 0.145990, 0.0421429, 0.00872877


(b) 0.938298, 0.672145, 0.362640, 0.147659, 0.0454100, 0.0104931

(c) 0.932253, 0.664778, 0.359567, 0.148416, 0.0469514, 0.0113722

(d) 0.928649, 0.660463, 0.357783, 0.148848, 0.0478492, 0.0118978

17. The Euler formula is $y_{n+1} = y_n + h(y_n^2 + 2t_n y_n)/(3 + t_n^2)$. The initial point is $(t_0, y_0) = (1, 2)$. Using this iteration formula with the specified h values, the value of the solution at $t = 2.5$ is somewhere between 18 and 19. At $t = 3$ there is no reliable estimate.

19.(a)



(b) The iteration formula is $y_{n+1} = y_n + h y_n^2 - h t_n^2$. The critical value α_0 appears to be between 0.67 and 0.68. For $y_0 > \alpha_0$, the iterations diverge.

20.(a) The ODE is linear, with general solution $y(t) = t + ce^t$. Invoking the specified initial condition, $y(t_0) = y_0$, we have $y_0 = t_0 + ce^{t_0}$. Hence $c = (y_0 - t_0)e^{-t_0}$. Thus the solution is given by $\phi(t) = (y_0 - t_0)e^{t-t_0} + t$.

(b) The Euler formula is $y_{n+1} = (1 + h)y_n + h - ht_n$. Now set $k = n + 1$.

(c) We have $y_1 = (1 + h)y_0 + h - ht_0 = (1 + h)y_0 + (t_1 - t_0) - ht_0$. Rearranging the terms, $y_1 = (1 + h)(y_0 - t_0) + t_1$. Now suppose that $y_k = (1 + h)^k(y_0 - t_0) + t_k$, for some $k \geq 1$. Then $y_{k+1} = (1 + h)y_k + h - ht_k$. Substituting for y_k , we find

that

$$y_{k+1} = (1+h)^{k+1}(y_0 - t_0) + (1+h)t_k + h - ht_k = (1+h)^{k+1}(y_0 - t_0) + t_k + h.$$

Noting that $t_{k+1} = t_k + h$, the result is verified.

(d) Substituting $h = (t - t_0)/n$, with $t_n = t$, $y_n = (1 + (t - t_0)/n)^n(y_0 - t_0) + t$. Taking the limit of both sides, and using the fact that $\lim_{n \rightarrow \infty} (1 + a/n)^n = e^a$, pointwise convergence is proved.

21. The exact solution is $y(t) = e^{2t}$. The Euler formula is $y_{n+1} = (1 + 2h)y_n$. It is easy to see that $y_n = (1 + 2h)^n y_0 = (1 + 2h)^n$. Given $t > 0$, set $h = t/n$. Taking the limit, we find that $\lim_{n \rightarrow \infty} y_n = \lim_{n \rightarrow \infty} (1 + 2t/n)^n = e^{2t}$.

23. The exact solution is $y(t) = t/2 + e^{2t}$. The Euler formula is $y_{n+1} = (1 + 2h)y_n + h/2 - ht_n$. Since $y_0 = 1$, $y_1 = (1 + 2h) + h/2 = (1 + 2h) + t_1/2$. It is easy to show by mathematical induction, that $y_n = (1 + 2h)^n + t_n/2$. For $t > 0$, set $h = t/n$ and thus $t_n = t$. Taking the limit, we find that $\lim_{n \rightarrow \infty} y_n = \lim_{n \rightarrow \infty} [(1 + 2t/n)^n + t/2] = e^{2t} + t/2$. Hence pointwise convergence is proved.

2.8

2. Let $z = y - 3$ and $\tau = t + 1$. It follows that $dz/d\tau = (dz/dt)(dt/d\tau) = dz/dt$. Furthermore, $dz/dt = dy/dt = 4 - y^3$. Hence $dz/d\tau = 4 - (z + 3)^3$. The new initial condition is $z(0) = 0$.

3.(a) The approximating functions are defined recursively by


$$\phi_{n+1}(t) = \int_0^t 3[\phi_n(s) + 1] ds.$$

Setting $\phi_0(t) = 0$, $\phi_1(t) = 3t$. Continuing, $\phi_2(t) = 9t^2/2 + 3t$, $\phi_3(t) = 9t^3/2 + 9t^2/2 + 3t$, $\phi_4(t) = 9t^4/4 + 9t^3/2 + 9t^2/2 + 3t$, Based upon these we conjecture that $\phi_n(t) = \sum_{k=1}^n 3^k t^k / k!$ and use mathematical induction to verify this form for $\phi_n(t)$. First, let $n = 1$, then $\phi_n(t) = 3t$, so it is certainly true for $n = 1$. Then, using Eq.(7) again we have

$$\phi_{n+1}(t) = \int_0^t 3[\phi_n(s) + 1] ds = \int_0^t 3 \left[\sum_{k=1}^n \frac{3^k}{k!} s^k + 1 \right] ds = \sum_{k=1}^{n+1} \frac{3^k}{k!} t^k,$$


and we have verified our conjecture.

(b)

(c) Recall from calculus that $e^{at} = 1 + \sum_{k=1}^{\infty} a^k t^k / k!$. Thus

$$\phi(t) = \sum_{k=1}^{\infty} \frac{3^k}{k!} t^k = e^{3t} - 1.$$

(d)




From the plot it appears that ϕ_4 is a good estimate for $|t| < 1/2$.

5.(a) The approximating functions are defined recursively by

$$\phi_{n+1}(t) = \int_0^t [-\phi_n(s)/3 + s] ds.$$


Setting $\phi_0(t) = 0$, $\phi_1(t) = t^2/2$. Continuing, $\phi_2(t) = t^2/2 - t^3/18$, $\phi_3(t) = t^2/2 - t^3/18 + t^4/216$, $\phi_4(t) = t^2/2 - t^3/18 + t^4/216 - t^5/3240$, ... Based upon these we conjecture that $\phi_n(t) = \sum_{k=1}^n 9(-1/3)^{k+1} t^{k+1} / (k+1)!$ and use mathematical induction to verify this form for $\phi_n(t)$.

(b)

(c) Recall from calculus that $e^{at} = 1 + \sum_{k=1}^{\infty} a^k t^k / k!$. Thus

$$\phi(t) = \sum_{k=1}^{\infty} 9 \frac{(-1/3)^{k+1}}{k+1!} t^{k+1} = 9e^{-t/3} + 3t - 9.$$

(d)




From the plot it appears that ϕ_4 is a good estimate for $|t| < 2$.

6.(a) The approximating functions are defined recursively by


$$\phi_{n+1}(t) = \int_0^t [\phi_n(s) + 1 - s] ds.$$

Setting $\phi_0(t) = 0$, $\phi_1(t) = t - t^2/2$, $\phi_2(t) = t - t^3/6$, $\phi_3(t) = t - t^4/24$, $\phi_4(t) = t - t^5/120$, Based upon these we conjecture that $\phi_n(t) = t - t^{n+1}/(n+1)!$ and use mathematical induction to verify this form for $\phi_n(t)$.

(b)

(c) Clearly $\phi(t) = t$.

(d)



From the plot it appears that ϕ_4 is a good estimate for $|t| < 1$.


8.(a) The approximating functions are defined recursively by

$$\phi_{n+1}(t) = \int_0^t [s^2 \phi_n(s) - s] ds.$$

Set $\phi_0(t) = 0$. The iterates are given by $\phi_1(t) = -t^2/2$, $\phi_2(t) = -t^2/2 - t^5/10$, $\phi_3(t) = -t^2/2 - t^5/10 - t^8/80$, $\phi_4(t) = -t^2/2 - t^5/10 - t^8/80 - t^{11}/880, \dots$ Upon inspection, it becomes apparent that

$$\begin{aligned} \phi_n(t) &= -t^2 \left[\frac{1}{2} + \frac{t^3}{2 \cdot 5} + \frac{t^6}{2 \cdot 5 \cdot 8} + \dots + \frac{(t^3)^{n-1}}{2 \cdot 5 \cdot 8 \dots [2 + 3(n-1)]} \right] = \\ &= -t^2 \sum_{k=1}^n \frac{(t^3)^{k-1}}{2 \cdot 5 \cdot 8 \dots [2 + 3(k-1)]}. \end{aligned}$$

(b)



(c) Using the identity $\phi_n(t) = \phi_1(t) + [\phi_2(t) - \phi_1(t)] + [\phi_3(t) - \phi_2(t)] + \dots + [\phi_n(t) - \phi_{n-1}(t)]$, consider the series $\phi_1(t) + \sum_{k=1}^{\infty} [\phi_{k+1}(t) - \phi_k(t)]$. Fix any t value now. We use the Ratio Test to prove the convergence of this series:

$$\left| \frac{\phi_{k+1}(t) - \phi_k(t)}{\phi_k(t) - \phi_{k-1}(t)} \right| = \left| \frac{\frac{(-t^2)(t^3)^k}{2 \cdot 5 \cdots (2+3k)}}{\frac{(-t^2)(t^3)^{k-1}}{2 \cdot 5 \cdots (2+3(k-1))}} \right| = \frac{|t|^3}{2+3k}.$$


The limit of this quantity is 0 for any fixed t as $k \rightarrow \infty$, and we obtain that $\phi_n(t)$ is convergent for any t .

9.(a) The approximating functions are defined recursively by

$$\phi_{n+1}(t) = \int_0^t [2s^2 + \phi_n^2(s)] ds.$$

Set $\phi_0(t) = 0$. The first three iterates are given by $\phi_1(t) = 2t^3/3$, $\phi_2(t) = 2t^3/3 + 4t^7/63$, $\phi_3(t) = 2t^3/3 + 4t^7/63 + 16t^{11}/2079 + 16t^{15}/59535$.

(b)



The iterates appear to be converging.

12.(a) The approximating functions are defined recursively by


$$\phi_{n+1}(t) = \int_0^t \left[\frac{3s^2 + 4s + 2}{2(\phi_n(s) - 1)} \right] ds.$$

Note that $1/(2y - 2) = -(1/2) \sum_{k=0}^6 y^k + O(y^7)$. For computational purposes, use the geometric series sum to replace the above iteration formula by

$$\phi_{n+1}(t) = -\frac{1}{2} \int_0^t \left[(3s^2 + 4s + 2) \sum_{k=0}^6 \phi_n^k(s) \right] ds.$$

Set $\phi_0(t) = 0$. The first four approximations are given by $\phi_1(t) = -t - t^2 - t^3/2$, $\phi_2(t) = -t - t^2/2 + t^3/6 + t^4/4 - t^5/5 - t^6/24 + \dots$, $\phi_3(t) = -t - t^2/2 + t^4/12 - 3t^5/20 + 4t^6/45 + \dots$, $\phi_4(t) = -t - t^2/2 + t^4/8 - 7t^5/60 + t^6/15 + \dots$

(b)



The approximations appear to be converging to the exact solution, which can be found by separating the variables: $\phi(t) = 1 - \sqrt{1 + 2t + 2t^2 + t^3}$.

14.(a) $\phi_n(0) = 0$, for every $n \geq 1$. Let $a \in (0, 1]$. Then $\phi_n(a) = 2na e^{-na^2} = 2na/e^{na^2}$. Using l'Hospital's rule, $\lim_{z \rightarrow \infty} 2az/e^{az^2} = \lim_{z \rightarrow \infty} 1/ze^{az^2} = 0$. Hence $\lim_{n \rightarrow \infty} \phi_n(a) = 0$.

(b) $\int_0^1 2nx e^{-nx^2} dx = -e^{-nx^2} \Big|_0^1 = 1 - e^{-n}$. Therefore,

$$\lim_{n \rightarrow \infty} \int_0^1 \phi_n(x) dx \neq \int_0^1 \lim_{n \rightarrow \infty} \phi_n(x) dx.$$

15. Let t be fixed, such that $(t, y_1), (t, y_2) \in D$. Without loss of generality, assume that $y_1 < y_2$. Since f is differentiable with respect to y , the mean value theorem asserts that there exists $\xi \in (y_1, y_2)$ such that $f(t, y_1) - f(t, y_2) = f_y(t, \xi)(y_1 - y_2)$. This means that $|f(t, y_1) - f(t, y_2)| = |f_y(t, \xi)| |y_1 - y_2|$. Since, by assumption, $\partial f / \partial y$ is continuous in D , f_y attains a maximum K on any closed and bounded subset of D . Hence $|f(t, y_1) - f(t, y_2)| \leq K |y_1 - y_2|$.

16. For a sufficiently small interval of t , $\phi_{n-1}(t), \phi_n(t) \in D$. Since f satisfies a Lipschitz condition, $|f(t, \phi_n(t)) - f(t, \phi_{n-1}(t))| \leq K |\phi_n(t) - \phi_{n-1}(t)|$. Here $K = \max |f_y|$.

17.(a) $\phi_1(t) = \int_0^t f(s, 0)ds$. Hence $|\phi_1(t)| \leq \int_0^{|t|} |f(s, 0)| ds \leq \int_0^{|t|} M ds = M|t|$, in which M is the maximum value of $|f(t, y)|$ on D .

(b) By definition, $\phi_2(t) - \phi_1(t) = \int_0^t [f(s, \phi_1(s)) - f(s, 0)] ds$. Taking the absolute value of both sides, $|\phi_2(t) - \phi_1(t)| \leq \int_0^{|t|} |[f(s, \phi_1(s)) - f(s, 0)]| ds$. Based on the results in Problems 16 and 17,

$$|\phi_2(t) - \phi_1(t)| \leq \int_0^{|t|} K |\phi_1(s) - 0| ds \leq KM \int_0^{|t|} |s| ds.$$

Evaluating the last integral, we obtain that $|\phi_2(t) - \phi_1(t)| \leq MK|t|^2/2$.

(c) Suppose that

$$|\phi_i(t) - \phi_{i-1}(t)| \leq \frac{MK^{i-1}|t|^i}{i!}$$

for some $i \geq 1$. By definition,

$$\phi_{i+1}(t) - \phi_i(t) = \int_0^t [f(s, \phi_i(s)) - f(s, \phi_{i-1}(s))] ds.$$

It follows that

$$\begin{aligned} |\phi_{i+1}(t) - \phi_i(t)| &\leq \int_0^{|t|} |f(s, \phi_i(s)) - f(s, \phi_{i-1}(s))| ds \\ &\leq \int_0^{|t|} K |\phi_i(s) - \phi_{i-1}(s)| ds \leq \int_0^{|t|} K \frac{MK^{i-1}|s|^i}{i!} ds = \\ &= \frac{MK^i|t|^{i+1}}{(i+1)!} \leq \frac{MK^i h^{i+1}}{(i+1)!}. \end{aligned}$$

Hence, by mathematical induction, the assertion is true.

18.(a) Use the triangle inequality, $|a + b| \leq |a| + |b|$.

(b) For $|t| \leq h$, $|\phi_1(t)| \leq Mh$, and $|\phi_n(t) - \phi_{n-1}(t)| \leq MK^{n-1}h^n/(n!)$. Hence

$$|\phi_n(t)| \leq M \sum_{i=1}^n \frac{K^{i-1}h^i}{i!} = \frac{M}{K} \sum_{i=1}^n \frac{(Kh)^i}{i!}.$$

(c) The sequence of partial sums in (b) converges to $M(e^{Kh} - 1)/K$. By the comparison test, the sums in (a) also converge. Since individual terms of a convergent series must tend to zero, $|\phi_n(t) - \phi_{n-1}(t)| \rightarrow 0$, and it follows that the sequence $|\phi_n(t)|$ is convergent.

19.(a) Let $\phi(t) = \int_0^t f(s, \phi(s))ds$ and $\psi(t) = \int_0^t f(s, \psi(s))ds$. Then by linearity of the integral, $\phi(t) - \psi(t) = \int_0^t [f(s, \phi(s)) - f(s, \psi(s))] ds$.

(b) It follows that $|\phi(t) - \psi(t)| \leq \int_0^{|t|} |f(s, \phi(s)) - f(s, \psi(s))| ds$.

(c) We know that f satisfies a Lipschitz condition, $|f(t, y_1) - f(t, y_2)| \leq K |y_1 - y_2|$, based on $|\partial f / \partial y| \leq K$ in D . Therefore,

$$|\phi(t) - \psi(t)| \leq \int_0^t |f(s, \phi(s)) - f(s, \psi(s))| ds \leq \int_0^t K |\phi(s) - \psi(s)| ds.$$

2.9

1. Writing the equation for each $n \geq 0$, $y_1 = -0.8 y_0$, $y_2 = -0.8 y_1 = (-0.8)^2 y_0$, $y_3 = -0.8 y_2 = (-0.8)^3 y_0$ and so on, it is apparent that $y_n = (-0.8)^n y_0$. The terms constitute an alternating series, which converge to zero, regardless of y_0 .

3. Write the equation for each $n \geq 0$, $y_1 = \sqrt{2} y_0$, $y_2 = \sqrt{3/2} y_1$, $y_3 = \sqrt{4/3} y_2$, ... Upon substitution, we find that $y_2 = \sqrt{(2 \cdot 3)/2} y_0 = \sqrt{3} y_0$, $y_3 = \sqrt{(4 \cdot 3 \cdot 2)/(3 \cdot 2)} y_0 = \sqrt{4} y_0$, ... It can be proved by mathematical induction, that $y_n = \sqrt{n+1} y_0$. This sequence is divergent, except for $y_0 = 0$.

4. Writing the equation for each $n \geq 0$, $y_1 = -y_0$, $y_2 = y_1$, $y_3 = -y_2$, $y_4 = y_3$, and so on. It can be shown that

$$y_n = \begin{cases} y_0, & \text{for } n = 4k \text{ or } n = 4k - 1 \\ -y_0, & \text{for } n = 4k - 2 \text{ or } n = 4k - 3 \end{cases}$$

The sequence is convergent only for $y_0 = 0$.

6. Writing the equation for each $n \geq 0$,

$$\begin{aligned} y_1 &= -0.5 y_0 + 4 \\ y_2 &= -0.5 y_1 + 4 = -0.5(-0.5 y_0 + 4) + 4 = (-0.5)^2 y_0 + 4 + (-0.5)4 \\ y_3 &= -0.5 y_2 + 4 = -0.5(-0.5 y_1 + 4) + 4 = (-0.5)^3 y_0 + 4 [1 + (-0.5) + (-0.5)^2] \\ &\vdots \\ y_n &= (-0.5)^n y_0 + (8/3) [1 - (-0.5)^n] \end{aligned}$$

which follows from Eq.(13) and (14). The sequence is convergent for all y_0 , and in fact $y_n \rightarrow 8/3$.

8. Let y_n be the balance at the end of the n th month. Then $y_{n+1} = (1 + r/12)y_n + 25$. We have $y_n = \rho^n [y_0 - 25/(1 - \rho)] + 25/(1 - \rho)$, in which $\rho = (1 + r/12)$. Here r is the annual interest rate, given as 7%. Thus $y_{48} = (1.00583333)^{48} [1000 + 12 \cdot 25/r] - 12 \cdot 25/r = \$2,702.28$.

9. Let y_n be the balance due at the end of the n th month. The appropriate difference equation is $y_{n+1} = (1 + r/12) y_n - P$. Here r is the annual interest rate and P is the monthly payment. The solution, in terms of the amount borrowed, is given by $y_n = \rho^n [y_0 + P/(1 - \rho)] - P/(1 - \rho)$, in which $\rho = (1 + r/12)$ and $y_0 = 9,000$. To figure out the monthly payment P , we require that $y_{36} = 0$. That is,

$\rho^{36}[y_0 + P/(1 - \rho)] = P/(1 - \rho)$. After the specified amounts are substituted, we find that $P = \$282.03$.

11. Let y_n be the balance due at the end of the n th month. The appropriate difference equation is $y_{n+1} = (1 + r/12)y_n - P$, in which $r = .06$ and P is the monthly payment. The initial value of the mortgage is $y_0 = \$100,000$. Then the balance due at the end of the n -th month is $y_n = \rho^n[y_0 + P/(1 - \rho)] - P/(1 - \rho)$, where $\rho = (1 + r/12)$. In terms of the specified values, $y_n = (1.005)^n[10^5 - 12P/r] + 12P/r$. Setting $n = 30 \cdot 12 = 360$, and $y_{360} = 0$, we find that $P = \$599.55$. For the monthly payment corresponding to a 20 year mortgage, set $n = 240$ and $y_{240} = 0$ to find that $P = \$716.43$. The total amount paid during the term of the loan is $360 \times 599.55 = \$215,838.00$ for the 30-year loan and is $240 \times 716.43 = \$171,943.20$ for the 20-year loan.


12. Let y_n be the balance due at the end of the n th month, with y_0 the initial value of the mortgage. The appropriate difference equation is $y_{n+1} = (1 + r/12)y_n - P$, in which $r = 0.08$ and $P = \$1000$ is the maximum monthly payment. Given that the life of the mortgage is 20 years, we require that $y_{240} = 0$. The balance due at the end of the n -th month is $y_n = \rho^n[y_0 + P/(1 - \rho)] - P/(1 - \rho)$. In terms of the specified values for the parameters, the solution of $(1.00666667)^{240}[y_0 - 12 \cdot 1000/0.08] = -12 \cdot 1000/0.08$ is $y_0 = \$119,554.29$.

19.(a) $\delta_2 = (\rho_2 - \rho_1)/(\rho_3 - \rho_2) = (3.449 - 3)/(3.544 - 3.449) = 4.7263$.

(b) $\text{diff} = (|\delta - \delta_2|/\delta) \cdot 100 = (|4.6692 - 4.7363|/4.6692) \cdot 100 \approx 1.22\%$.

(c) Assuming $(\rho_3 - \rho_2)/(\rho_4 - \rho_3) = \delta$, $\rho_4 \approx 3.5643$

(d) A period 16 solution appears near $\rho \approx 3.565$.



(e) Note that $(\rho_{n+1} - \rho_n) = \delta_n^{-1}(\rho_n - \rho_{n-1})$. With the assumption that $\delta_n = \delta$, we have $(\rho_{n+1} - \rho_n) = \delta^{-1}(\rho_n - \rho_{n-1})$, which is of the form $y_{n+1} = \alpha y_n$, $n \geq 3$. It

follows that $(\rho_k - \rho_{k-1}) = \delta^{3-k}(\rho_3 - \rho_2)$ for $k \geq 4$. Then

$$\begin{aligned}\rho_n &= \rho_1 + (\rho_2 - \rho_1) + (\rho_3 - \rho_2) + (\rho_4 - \rho_3) + \dots + (\rho_n - \rho_{n-1}) \\ &= \rho_1 + (\rho_2 - \rho_1) + (\rho_3 - \rho_2) [1 + \delta^{-1} + \delta^{-2} + \dots + \delta^{3-n}] \\ &= \rho_1 + (\rho_2 - \rho_1) + (\rho_3 - \rho_2) \left[\frac{1 - \delta^{4-n}}{1 - \delta^{-1}} \right].\end{aligned}$$

Hence $\lim_{n \rightarrow \infty} \rho_n = \rho_2 + (\rho_3 - \rho_2) \left[\frac{\delta}{\delta - 1} \right]$. Substitution of the appropriate values yields

$$\lim_{n \rightarrow \infty} \rho_n = 3.5699$$

PROBLEMS

1. The equation is *linear*. It can be written in the form $y' + 2y/x = x^3$, and the integrating factor is $\mu(x) = e^{\int (2/x) dx} = e^{2 \ln x} = x^2$. Multiplication by $\mu(x)$ yields $x^2 y' + 2yx = (yx^2)' = x^5$. Integration with respect to x and division by x^2 gives that $y = x^4/6 + c/x^2$.

5. The equation is *exact*. Algebraic manipulations give the symmetric form of the equation, $(2xy + y^2 + 3)dx + (x^2 + 2xy)dy = 0$. We can check that $M_y = 2x + 2y = N_x$, so the equation is really exact. Integrating M with respect to x gives that $\psi(x, y) = x^2 y + xy^2 + 3x + g(y)$, then $\psi_y = x^2 + 2xy + g'(y) = x^2 + 2xy$, so we get that $g'(y) = 0$, so we obtain that $g(y) = 0$ is acceptable. Therefore the solution is defined implicitly as $x^2 y + xy^2 + 3x = c$.

6. The equation is *linear*. It can be written in the form $y' + (1 + (1/x))y = 1/x$ and the integrating factor is $\mu(x) = e^{\int 1 + (1/x) dx} = e^{x + \ln x} = xe^x$. Multiplication by $\mu(x)$ yields $xe^x y' + (xe^x + e^x)y = (xe^x y)' = e^x$. Integration with respect to x and division by xe^x shows that the general solution of the equation is $y = 1/x + c/(xe^x)$. The initial condition implies that $0 = 1 + c/e$, which means that $c = -e$ and the solution is $y = 1/x - e/(xe^x) = x^{-1}(1 - e^{1-x})$.

7. The equation is *separable*. Separation of variables gives the differential equation $y(2 + 3y)dy = (5x^4 + 1)dx$, and then after integration we obtain that the solution is $x^5 + x - y^2 - y^3 = c$.

8. The equation is *linear*. It can be written in the form $y' + 2y/x = \cos x/x^2$ and the integrating factor is $\mu(x) = e^{\int (2/x) dx} = e^{2 \ln x} = x^2$. Multiplication by $\mu(x)$ gives $x^2 y' + 2xy = (x^2 y)' = \cos x$, and after integration with respect to x and division by x^2 we obtain the general solution $y = (c + \sin x)/x^2$. The initial condition implies that $c = 4 - \sin 2$ and the solution becomes $y = (4 - \sin 2 + \sin x)/x^2$.

11. The equation is *exact*. It is easy to check that $M_y = 1 = N_x$. Integrating M with respect to x gives that $\psi(x, y) = x^4/4 + xy + g(y)$, then $\psi_y = x + g'(y) =$

$x + e^y$, which means that $g'(y) = e^y$, so we obtain that $g(y) = e^y$. Therefore the solution is defined implicitly as $x^4/4 + xy + e^y = c$.

13. The equation is *separable*. Factoring the right hand side leads to the equation $y' = (1 + y^2)(1 + 3x^2)$. We separate the variables to obtain $dy/(1 + y^2) = (1 + 3x^2)dx$, then integration gives us $\arctan y = x + x^3 + c$. The solution is $y = \tan(x + x^3 + c)$.

14. The equation is *exact*. We can check that $M_y = 1 = N_x$. Integrating M with respect to x gives that $\psi(x, y) = x^2/2 + xy + g(y)$, then $\psi_y = x + g'(y) = x + 2y$, which means that $g'(y) = 2y$, so we obtain that $g(y) = y^2$. Therefore the general solution is defined implicitly as $x^2/2 + xy + y^2 = c$. The initial condition gives us $c = 10$, so the solution is $x^2 + 2xy + 2y^2 = 20$.

15. The equation is *separable*. Separation of variables leads us to the equation

$$\frac{dy}{y} = \frac{1 - e^x}{1 + e^x} dx.$$

Note that $1 + e^x - 2e^x = 1 - e^x$. We obtain that

$$\ln |y| = \int \frac{1 - e^x}{1 + e^x} dx = \int 1 - \frac{2e^x}{1 + e^x} dx = x - 2 \ln(1 + e^x) + \tilde{c}.$$

This means that $y = ce^x(1 + e^x)^{-2}$, which also can be written as $y = c/\cosh^2(x/2)$ after some algebraic manipulations.

16. The equation is *exact*. The symmetric form is $(-e^{-x} \cos y + e^{2y} \cos x)dx + (-e^{-x} \sin y + 2e^{2y} \sin x)dy = 0$. We can check that $M_y = e^{-x} \sin y + 2e^{2y} \cos x = N_x$. Integrating M with respect to x gives that $\psi(x, y) = e^{-x} \cos y + e^{2y} \sin x + g(y)$, then $\psi_y = -e^{-x} \sin y + 2e^{2y} \sin x + g'(y) = -e^{-x} \sin y + 2e^{2y} \sin x$, so we get that $g'(y) = 0$, so we obtain that $g(y) = 0$ is acceptable. Therefore the solution is defined implicitly as $e^{-x} \cos y + e^{2y} \sin x = c$.

17. The equation is *linear*. The integrating factor is $\mu(x) = e^{-\int 3 dx} = e^{-3x}$, which turns the equation into $e^{-3x}y' - 3e^{-3x}y = (e^{-3x}y)' = e^x$. We integrate with respect to x to obtain $e^{-3x}y = e^x + c$, and the solution is $y = ce^{3x} + e^{4x}$ after multiplication by e^{3x} .

18. The equation is *linear*. The integrating factor is $\mu(x) = e^{\int 2 dx} = e^{2x}$, which gives us $e^{2x}y' + 2e^{2x}y = (e^{2x}y)' = e^{-x^2}$. The antiderivative of the function on the right hand side can not be expressed in a closed form using elementary functions, so we have to express the solution using integrals. Let us integrate both sides of this equation from 0 to x . We obtain that the left hand side turns into

$$\int_0^x (e^{2s}y(s))' ds = e^{2x}y(x) - e^0y(0) = e^{2x}y - 4.$$

The right hand side gives us $\int_0^x e^{-s^2} ds$. So we found that

$$y = e^{-2x} \int_0^x e^{-s^2} ds + 4e^{-2x}.$$

19. The equation is *exact*. Algebraic manipulations give us the symmetric form $(y^3 + 2y - 4x^3)dx + (2x + 3xy^2)dy = 0$. We can check that $M_y = 3y^2 + 2 = N_x$. Integrating M with respect to x gives that $\psi(x, y) = xy^3 + 2xy - x^4 + g(y)$, then $\psi_y = 3xy^2 + 2x + g'(y) = 2x + 3xy^2$, which means that $g'(y) = 0$, so we obtain that $g(y) = 0$ is acceptable. Therefore the solution is $xy^3 + 2xy - x^4 = c$.

20. The equation is *separable*, because $y' = e^{x+y} = e^x e^y$. Separation of variables yields the equation $e^{-y} dy = e^x dx$, which turns into $-e^{-y} = e^x + c$ after integration and we obtain the implicitly defined solution $e^x + e^{-y} = c$.

22. The equation is *separable*. Separation of variables turns the equation into $(y^2 + 2)dy = (x^2 - 1)dx$, which, after integration, gives $y^3/3 + 2y = x^3/3 - x + c$. The initial condition yields $c = 5/3$, and the solution is $y^3 + 6y - x^3 + 3x = 5$.

23. The equation is *linear*. Division by t gives $y' + (1 + (1/t))y = e^{2t}/t$, so the integrating factor is $\mu(t) = e^{\int(1+(1/t))dt} = e^{t+\ln t} = te^t$. The equation turns into $te^t y' + (te^t + e^t)y = (te^t y)' = e^{3t}$. Integration therefore leads to $te^t y = e^{3t}/3 + c$ and the solution is $y = e^{2t}/(3t) + ce^{-t}/t$.

24. The equation is *exact*. We can check that $M_y = 2\cos y \sin x \cos x = N_x$. Integrating M with respect to x gives that $\psi(x, y) = \sin y \sin^2 x + g(y)$, then $\psi_y = \cos y \sin^2 x + g'(y) = \cos y \sin^2 x$, which means that $g'(y) = 0$, so we obtain that $g(y) = 0$ is acceptable. Therefore the solution is defined implicitly as $\sin y \sin^2 x = c$.

25. The equation is *exact*. We can check that

$$M_y = -\frac{2x}{y^2} - \frac{x^2 - y^2}{(x^2 + y^2)^2} = N_x.$$

Integrating M with respect to x gives that $\psi(x, y) = x^2/y + \arctan(y/x) + g(y)$, then $\psi_y = -x^2/y^2 + x/(x^2 + y^2) + g'(y) = x/(x^2 + y^2) - x^2/y^2$, which means that $g'(y) = 0$, so we obtain that $g(y) = 0$ is acceptable. Therefore the solution is defined implicitly as $x^2/y + \arctan(y/x) = c$.

28. The equation can be made *exact* by choosing an appropriate integrating factor. We can check that $(M_y - N_x)/N = (2 - 1)/x = 1/x$ depends only on x , so $\mu(x) = e^{\int(1/x)dx} = e^{\ln x} = x$ is an integrating factor. After multiplication, the equation becomes $(2yx + 3x^2)dx + x^2 dy = 0$. This equation is exact now, because $M_y = 2x = N_x$. Integrating M with respect to x gives that $\psi(x, y) = yx^2 + x^3 + g(y)$, then $\psi_y = x^2 + g'(y) = x^2$, which means that $g'(y) = 0$, so we obtain that $g(y) = 0$ is acceptable. Therefore the solution is defined implicitly as $x^3 + x^2 y = c$.

29. The equation is *homogeneous*. (See Section 2.2, Problem 30) We can see that

$$y' = \frac{x+y}{x-y} = \frac{1+(y/x)}{1-(y/x)}.$$

We substitute $u = y/x$, which means also that $y = ux$ and then $y' = u'x + u =$

$(1-u)/(1+u)$, which implies that

$$u'x = \frac{1-u}{1+u} - u = \frac{1-2u-u^2}{1+u},$$

a separable equation. Separating the variables yields

$$\frac{1+u}{1-2u-u^2} du = \frac{dx}{x},$$

and then integration gives $-\ln(1-2u-u^2)/2 = \ln|x| + c$. Substituting $u = y/x$ back into this expression and using that

$$-\ln(1-2(y/x)-(y/x)^2)/2 - \ln|x| = -\ln(|x|\sqrt{1-2y/x-(y/x)^2}) = -\ln(\sqrt{x^2-2yx-y^2})$$

we obtain that the solution is $x^2 - 2yx - y^2 = c$.

30. The equation is *homogeneous*. (See Section 2.2, Problem 30) Algebraic manipulations show that it can be written in the form

$$y' = \frac{3y^2 + 2xy}{2xy + x^2} = \frac{3(y/x)^2 + 2(y/x)}{2(y/x) + 1}.$$

Substituting $u = y/x$ gives that $y = ux$ and then

$$y' = u'x + u = \frac{3u^2 + 2u}{2u + 1},$$

which implies that

$$u'x = \frac{3u^2 + 2u}{2u + 1} - u = \frac{u^2 + u}{2u + 1},$$

a separable equation. We obtain that $(2u+1)du/(u^2+u) = dx/x$, which in turn means that $\ln(u^2+u) = \ln|x| + \tilde{c}$. Therefore, $u^2+u = cx$ and then substituting $u = y/x$ gives us the solution $(y^2/x^3) + (y/x^2) = c$.

31. The equation can be made *exact* by choosing an appropriate integrating factor. We can check that $(M_y - N_x)/M = -(3x^2 + y)/(y(3x^2 + y)) = -1/y$ depends only on y , so $\mu(y) = e^{\int (1/y) dy} = e^{\ln y} = y$ is an integrating factor. After multiplication, the equation becomes $(3x^2y^2 + y^3)dx + (2x^3y + 3xy^2)dy = 0$. This equation is exact now, because $M_y = 6x^2y + 3y^2 = N_x$. Integrating M with respect to x gives that $\psi(x, y) = x^3y^2 + y^3x + g(y)$, then $\psi_y = 2x^3y + 3y^2x + g'(y) = 2x^3y + 3xy^2$, which means that $g'(y) = 0$, so we obtain that $g(y) = 0$ is acceptable. Therefore the general solution is defined implicitly as $x^3y^2 + xy^3 = c$. The initial condition gives us $4 - 8 = c = -4$, and the solution is $x^3y^2 + xy^3 = -4$.

33. Let y_1 be a solution, i.e. $y_1' = q_1 + q_2y_1 + q_3y_1^2$. Now let $y = y_1 + (1/v)$ also be a solution. Differentiating this expression with respect to t and using that y is also a solution we obtain $y' = y_1' - (1/v^2)v' = q_1 + q_2y + q_3y^2 = q_1 + q_2(y_1 + (1/v)) + q_3(y_1 + (1/v))^2$. Now using that y_1 was also a solution we get that $-(1/v^2)v' = q_2(1/v) + 2q_3(y_1/v) + q_3(1/v^2)$, which, after some simple algebraic manipulations turns into $v' = -(q_2 + 2q_3y_1)v - q_3$.

35.(a) The equation is $y' = (1 - y)(x + by) = x + (b - x)y - by^2$. We set $y = 1 + (1/v)$ and differentiate: $y' = -v^{-2}v' = x + (b - x)(1 + (1/v)) - b(1 + (1/v))^2$, which, after simplification, turns into $v' = (b + x)v + b$.

(b) When $x = at$, the equation is $v' - (b + at)v = b$, so the integrating factor is $\mu(t) = e^{-bt - at^2/2}$. This turns the equation into $(v\mu(t))' = b\mu(t)$, so $v\mu(t) = \int b\mu(t)dt$, and then $v = (b \int \mu(t)dt)/\mu(t)$.

36. Substitute $v = y'$, then $v' = y''$. The equation turns into $t^2v' + 2tv = (t^2v)' = 2$, which yields $t^2v = 2t + c_1$, so $y' = v = (2/t) + (c_1/t^2)$. Integrating this expression gives us the solution $y = 2 \ln t - (c_1/t) + c_2$.

37. Set $v = y'$, then $v' = y''$. The equation with this substitution is $tv' + v = (tv)' = 4$, which gives $tv = 4t + c_1$, so $y' = v = 4 + (c_1/t)$. Integrating this expression yields the solution $y = 4t + c_1 \ln t + c_2$.

38. Set $v = y'$, so $v' = y''$. The equation is $v' + 2tv^2 = 0$, which is a separable equation. Separating the variables we obtain $dv/v^2 = -2tdt$, so $-1/v = -t^2 + c$, and then $y' = v = 1/(t^2 + c_1)$. Now depending on the value of c_1 , we have the following possibilities: when $c_1 = 0$, then $y = -1/t + c_2$, when $0 < c_1 = k^2$, then $y = (1/k) \arctan(t/k) + c_2$, and when $0 > c_1 = -k^2$ then

$$y = (1/2k) \ln |(t - k)/(t + k)| + c_2.$$

We also divided by $v = y'$ when we separated the variables, and $v = 0$ (which is $y = c$) is also a solution.

39. Substitute $v = y'$ and $v' = y''$. The equation is $2t^2v' + v^3 = 2tv$. This is a *Bernoulli* equation (See Section 2.4, Problem 27), so the substitution $z = v^{-2}$ yields $z' = -2v^{-3}v'$, and the equation turns into $2t^2v'v^3 + 1 = 2t/v^2$, i.e. into $-2t^2z'/2 + 1 = 2tz$, which in turn simplifies to $t^2z' + 2tz = (t^2z)' = 1$. Integration yields $t^2z = t + c$, which means that $z = (1/t) + (c/t^2)$. Now $y' = v = \pm\sqrt{1/z} = \pm t/\sqrt{t + c_1}$ and another integration gives

$$y = \pm \frac{2}{3}(t - 2c_1)\sqrt{t + c_1} + c_2.$$

The substitution also loses the solution $v = 0$, i.e. $y = c$.

40. Set $v = y'$, then $v' = y''$. The equation reads $v' + v = 2e^{-t}$, which is a linear equation with integrating factor $\mu(t) = e^t$. This turns the equation into $e^tv' + e^tv = (e^tv)' = 2$, which means that $e^tv = 2t + c$ and then $y' = v = 2te^{-t} + ce^{-t}$. Another integration yields the solution $y = -2te^{-t} + c_1e^{-t} + c_2$.

41. Let $v = y'$ and $v' = y''$. The equation is $t^2v' = v^2$, which is a separable equation. Separating the variables we obtain $dv/v^2 = dt/t^2$, which gives us $-1/v = -(1/t) + c_1$, and then $y' = v = t/(1 + c_1t)$. Now when $c_1 = 0$, then $y = t^2/2 + c_2$, and when $c_1 \neq 0$, then $y = t/c_1 - (\ln |1 + c_1t|)/c_1^2 + c_2$. Also, at the separation we divided by $v = 0$, which also gives us the solution $y = c$.

43. Set $y' = v(y)$. Then $y'' = v'(y)(dy/dt) = v'(y)v(y)$. We obtain the equation $v'v + y = 0$, where the differentiation is with respect to y . This is a separable equation which simplifies to $v dv = -y dy$. We obtain that $v^2/2 = -y^2/2 + c$, so $y' = v(y) = \pm\sqrt{c - y^2}$. We separate the variables again to get $dy/\sqrt{c - y^2} = \pm dt$, so $\arcsin(y/\sqrt{c}) = t + d$, which means that $y = \sqrt{c} \sin(\pm t + d) = c_1 \sin(t + c_2)$.

44. Set $y' = v(y)$. Then $y'' = v'(y)(dy/dt) = v'(y)v(y)$. We obtain the equation $v'v + yv^3 = 0$, where the differentiation is with respect to y . Separation of variables turns this into $dv/v^2 = -y dy$, which gives us $y' = v = 2/(c_1 + y^2)$. This implies that $(c_1 + y^2)dy = 2dt$ and then the solution is defined implicitly as $c_1 y + y^3/3 = 2t + c_2$. Also, $y = c$ is a solution which we lost when divided by $y' = v = 0$.

46. Set $y' = v(y)$. Then $y'' = v'(y)(dy/dt) = v'(y)v(y)$. We obtain the equation $yv'v - v^3 = 0$, where the differentiation is with respect to y . This separable equation gives us $dv/v^2 = dy/y$, which means that $-1/v = \ln|y| + c$, and then $y' = v = 1/(c - \ln|y|)$. We separate variables again to obtain $(c - \ln|y|)dy = dt$, and then integration yields the implicitly defined solution $cy - (y \ln|y| - y) = t + d$. Also, $y = c$ is a solution which we lost when we divided by $v = 0$.

49. Set $y' = v(y)$. Then $y'' = v'(y)(dy/dt) = v'(y)v(y)$. We obtain the equation $v'v - 3y^2 = 0$, where the differentiation is with respect to y . Separation of variables gives $v dv = 3y^2 dy$, and after integration this turns into $v^2/2 = y^3 + c$. The initial conditions imply that $c = 0$ here, so $(y')^2 = v^2 = 2y^3$. This implies that $y' = \sqrt{2}y^{3/2}$ (the sign is determined by the initial conditions again), and this separable equation now turns into $y^{-3/2} dy = \sqrt{2} dt$. Integration yields $-2y^{-1/2} = \sqrt{2}t + d$, and the initial conditions at this point give that $d = -\sqrt{2}$. Algebraic manipulations find that $y = 2(1 - t)^{-2}$.

50. Set $v = y'$, then $v' = y''$. The equation with this substitution turns into the equation $(1 + t^2)v' + 2tv = ((1 + t^2)v)' = -3t^{-2}$. Integrating this we get that $(1 + t^2)v = 3t^{-1} + c$, and $c = -5$ from the initial conditions. This means that $y' = v = 3/(t(1 + t^2)) - 5/(1 + t^2)$. The partial fraction decomposition of the first expression shows that $y' = 3/t - 3t/(1 + t^2) - 5/(1 + t^2)$ and then another integration here gives us that $y = 3 \ln t - (3/2) \ln(1 + t^2) - 5 \arctan t + d$. The initial conditions identify $d = 2 + (3/2) \ln 2 + 5\pi/4$, and we obtained the solution.