

Sediment model manuscript

Christopher M. Tasich^{a,1}, Jonathan Gilligan^{a,2}, and Steven L. Goodbred^{a,3}

^aVanderbilt University

This manuscript was compiled on November 19, 2020

Please provide an abstract of no more than 250 words in a single paragraph. Abstracts should explain to the general reader the major contributions of the article. References in the abstract must be cited in full within the abstract itself and cited in the text.

aggradation | sea level rise | tidal river management | polders | tidal amplification

Sea level rise threatens the densely populated and ecologically significant low-lying coastal region of Bengal. Mitigating the effects of sea level rise (SLR) in this region will be especially difficult considering current widespread land management practices and the ongoing geopolitical tension between India and Bangladesh.

Coastal Bengal is situated within the delta formed by the confluence of the Ganges and Brahmaputra rivers and straddles the border between West Bengal, India to the east and Bangladesh to the west. The region is home to ~35 million people and the ecologically critical Sundarbans mangrove forest. Since the 1960s, the region has seen a vast transformation through the reduction in natural mangrove habitat and the widespread construction of earthen embankments. These embankments surround large swaths of clearcut land, known as polders, that accommodate the swelling population of the region.

While, preventing regular inundation by spring high tides, the creation of these embankments have had the unintended consequence of starving the interior of the polders of fresh sediment. Without this sediment, polder interiors have compacted resulting in significant elevation offset (1.0 m to 1.5 m) relative to the natural mangrove forest⁽¹⁾. Polder elevations often sit precariously below local mean high water (MHW) levels leading to persistent waterlogging. Furthermore, many of these embankments are in disrepair and susceptible to breaching especially by storm surge as was the case with Cyclone Sidr (2007), Cyclone Aila (2009), and recently with Cyclone Amphan (2020).

Tidal river management (TRM) has been proposed as a possible augmentation of current land management practices to alleviate some of the issues caused by poldering. Under TRM, embankments are seasonally breached to allow tidal water inundation and sediment aggradation. Many low-lying areas may recover to an acceptable elevation within 5 to 10 years, though, some may take longer. There are a host of local socioeconomic, political, and governance consideration that may influence the success of TRM. Here, we neglect those considerations and focus on the general feasibility of TRM in regards to SLR and the sediment supply of the GB system. Future studies will focus on the social dynamics surround TRM.

Coastal Bengal is often seen as one of the most at-risk regions for SLR due to climate change. Infographics often depict large swaths of the Bengal coastline flooded under different SLR scenarios. However, this overly simplifies the threats to the region and neglects the significant sediment

contribution of the GB system in maintaining the natural elevation.

Estimates for increases in Relative Mean Sea Level (RMSL) in the GB delta range from 2.8 mm yr⁻¹ to 8.8 mm yr⁻¹. However, RMSL neglects the widening of the tidal range in the polder region. On average, local high water levels in the polder region are increasing at a rate of 15.9 mm yr⁻¹⁽²⁾. While SLR is of paramount importance, tidal range amplification is the more imminent threat to the region. This is especially important considering the Bangladeshi government's recent reinvestment in poldering with a \$400 million loan from the World Bank for the Coastal Embankment Improvement Project - Phase I (CEIP-I).

As for the natural mangrove system, it is unclear how changing water levels will affect elevation. Some studies have shown that the region is incredibly resilient to increasing water levels due, in large part, to the abundant sediment supply of the GB system. This sediment is delivered to the platform periodically during spring high tide which helps maintain an equilibrium elevation approximately equivalent to mean higher high water (MHHW). But, this large volume of sediment delivered to coastal Bengal is not guaranteed.

Water has long been the focus of the geopolitical disputes between India and Bangladesh. However, the reduction in waterflow across the border portends a significant decrease in sediment flux. Estimates suggest sediment flux may be reduced by 39 % to 75 % for the Ganges and 9 % to 25 % for the Brahmaputra resulting in a change in aggradation from 3.6 mm yr⁻¹ to 2.5 mm yr⁻¹⁽³⁾.

The combination of increasing water levels and decreasing sediment supply may further intensify an already dire situation. Here, we use a zero-dimensional mass balance model of sediment aggradation to understand the impact that increasing water levels and decreasing sediment flux will have on the regions equilibrium elevation and consequently its resilience to climate change. We consider both the resilience of the natural mangrove system and the ability of the polder system

Significance Statement

Authors must submit a 120-word maximum statement about the significance of their research paper written at a level understandable to an undergraduate educated scientist outside their field of speciality. The primary goal of the significance statement is to explain the relevance of the work in broad context to a broad readership. The significance statement appears in the paper itself and is required for all research papers.

Author contributions: C.M.T., J.G., and S.L.G. designed research; C.M.T. and J.G. performed research; C.M.T. analyzed data; C.M.T. wrote the paper; and C.M.T., J.G., and S.L.G. provided discussion and input on paper

The authors declare no conflict of interest.

¹To whom correspondence should be addressed. E-mail: chris.tasichvanderbilt.edu

84 to recover to a more resilient elevation through TRM.

85 **Numerical model.** We modeled tidal platform elevation (ζ)
86 using a zero-dimensional mass balance model using the basic
87 formulation provided by Krone (4) and further refined by Allen
88 (5), French (6), and Temmerman et al. (7, 8). The rate of
89 tidal platform elevation change is described as

$$\frac{d\zeta}{dt} = \frac{dS_M}{dt} + \frac{dS_O}{dt} + \frac{dP}{dt} + \frac{dM}{dt}, \quad [1]$$

91 where S_M is mineral sedimentation, S_O is organic matter
92 sedimentation, P is compaction, and M is tectonic subsidence.
93 Each term of equation 1 can be further expanded.

94 We approximate S_M as

$$S_M(t) = \int \frac{w_s C(t)}{\rho_b} dt, \quad [2]$$

96 where w_s is the characteristic settling velocity of a given grain
97 size, $C(t)$ is the depth-averaged and time-varying sediment
98 concentration in the water column, and ρ_b is the dry bulk
99 density of the sediment. We assume there is no resuspension
100 of mineral sediment which is consistent with Krone's (4) initial
101 formulation.

102 w_s is calculated for a given grain size by using Stokes' law
103 to determine the terminal velocity of a sphere falling through
104 a fluid given by

$$w_s = \frac{2}{9} \frac{\rho_p - \rho_f}{\mu} g R^2 \quad [3]$$

106 where ρ_p is the mass density of the particle or grain, ρ_f is
107 the mass density of the fluid, μ is the dynamic viscosity of the
108 fluid, g is the acceleration due to gravity, and R is the radius
109 of the grain. This is approximation for w_s is consistent with
110 previous similar studies (5, 7, 8). We assume basic properties
111 of water for ρ_f (1000 kg m^{-3}) and μ ($1 \times 10^{-3} \text{ kg m}^{-1} \text{ s}^{-1}$) for
112 simplicity. Salinity does vary seasonally which will change
113 these values, but had little affect on the model output.

114 We capture the temporal variation of $C(t)$ through the
115 mass balance given as

$$\frac{d[h(t) - \zeta(t)]C(t)}{dt} = -w_s C(t) + C_{in} \frac{dh}{dt}, \quad [4]$$

117 where h is the height of the water column and C_{in} is the
118 incoming suspended sediment concentration of the adjacent
119 water column. We consider ζ to be a function of time and
120 update it at every timestep which differs from previous studies
121 (4–9) which only update ζ after every tidal cycle. The physical
122 interpretation of equation 4 is that the first term is the mass
123 flux above an area on the tidal platform, the second term is
124 the mass flux extracted from the water column, and the third
125 term is the mass flux from the adjacent water column. Further
126 derivation of equation 4 results in

$$\frac{dC}{dt} [h(t) - \zeta(t)] = -w_s C(t) + [C_{in} - C(t)] \frac{dh}{dt} + C(t) \frac{d\zeta}{dt} \quad [5]$$

128 From equation 5, we can approximate the solution for
129 concentration numerically.

Model inputs. We obtained model inputs from field measure-
ments. We observed tidal height, grain size, suspended sedi-
ment concentration (SSC), and dry bulk density around Polder
32 over multiple field seasons from 2011 to 2016 (1, 10, 11).

For h , we extracted one year of contiguous tidal data from
from a pressure sensor deployed within the tidal channel near
Polder 32. We used the oce package in R (3.6.3) to create
an idealized tidal curve from our data (12). The tidal curve
was then shifted down so that mean higher high water would
be 0.3 m above the Sundarban platform and 1.8 m above the
polder surface (ζ). We replicated this tidal curve for each
subsequent year for the length of the model run. Field obser-
vations confirm these benchmark elevation (1, 11, 13). In order
to simulate sea level rise, the subsequent year tidal curves were
increased at a linear rate of 2 mm yr^{-1} which is consistent with
field observations.

For C_{in} , we use observed values of SSC from Hale et al.
(10) that are characteristic of the tidal channels in the region.
Similar to Temmerman et al. (7, 8), we scaled the observed
tidal channel SSC by a factor as the flood waters are expected
to have a lower SSC than the tidal channel due to lower flow
velocities. For our preliminary study, we use a k-factor of
0.7. In future model iterations, we will better explore this
relationship and determine an appropriate k-factor.

For ρ , we used values derived from conversations with
Steven Goodbred and Carol Wilson.

Materials and Methods

Please describe your materials and methods here. This can be more
than one paragraph, and may contain subsections and equations as
required.

Subsection for Method. Example text for subsection.

ACKNOWLEDGMENTS. Please include your acknowledgments
here, set in a single paragraph. Please do not include any acknowl-
edgments in the Supporting Information, or anywhere else in the
manuscript.

1. LW Auerbach, et al., Flood risk of natural and embanked landscapes on the Ganges–Brahmaputra tidal delta plain. *Nat. Clim. Chang.* **5**, 153–157 (2015).
2. J Pethick, JD Orford, Rapid rise in effective sea-level in southwest Bangladesh: Its causes and contemporary rates. *Glob. Planet. Chang.* **111**, 237–245 (2013).
3. SA Higgins, I Overeem, KG Rogers, EA Kalina, River linking in India: Downstream impacts on water discharge and suspended sediment transport to deltas. *Elem Sci Anth* **6**, 20 (2018).
4. R Krone, A Method for Simulating Marsh Elevations in *Coastal Sediments*. (American Society of Civil Engineers, New Orleans, Louisiana), pp. 316–323 (1987).
5. JRL Allen, Salt-marsh growth and stratification: A numerical model with special reference to the Severn Estuary, southwest Britain. *Mar. Geol.* **95**, 77–96 (1990).
6. JR French, Numerical simulation of vertical marsh growth and adjustment to accelerated sea-level rise, North Norfolk, U.K. *Earth Surf. Process. Landforms* **18**, 63–81 (1993).
7. S Temmerman, G Govers, P Meire, S Wartel, Modelling long-term tidal marsh growth under changing tidal conditions and suspended sediment concentrations, Scheldt estuary, Belgium. *Mar. Geol.* **193**, 151–169 (2003).
8. S Temmerman, G Govers, S Wartel, P Meire, Modelling estuarine variations in tidal marsh sedimentation: Response to changing sea level and suspended sediment concentrations. *Mar. Geol.* **212**, 1–19 (2004).
9. J French, Tidal marsh sedimentation and resilience to environmental change: Exploratory modelling of tidal, sea-level and sediment supply forcing in predominantly allocthonous systems. *Mar. Geol.* **235**, 119–136 (2006).
10. R Hale, R Bain, S Goodbred Jr., J Best, Observations and scaling of tidal mass transport across the lower Ganges–Brahmaputra delta plain: Implications for delta management and sustainability. *Earth Surf. Dyn.* **7**, 231–245 (2019).
11. RP Hale, CA Wilson, EJ Bomer, Seasonal Variability of Forces Controlling Sedimentation in the Sundarbans National Forest, Bangladesh. *Front. Earth Sci.* **7**, 211 (2019).
12. D Kelley, C Richards, Oce: Analysis of Oceanographic Data (2020).
13. EJ Bomer, CA Wilson, RP Hale, ANM Hossain, FMA Rahman, Surface elevation and sedimentation dynamics in the Ganges–Brahmaputra tidal delta plain, Bangladesh: Evidence for mangrove adaptation to human-induced tidal amplification. *CATENA* **187**, 104312 (2020).