

Crosby Lab Code

by

Christelinda Laureijs

Scripts for analyzing evoked currents and action potentials in R

June 20, 2024

Contents

| | |
|-----------------------------------|---|
| Evoked Currents (eEPSCs) Analysis | 2 |
| Raw plots | 2 |
| Pruned individual plots | 2 |
| Pruned summary plots | 2 |
| Statistical analysis | 2 |
| Levene's test | 2 |
| Univariate Shapiro | 2 |
| Box's M | 2 |
| Multivariate normality | 2 |
| T-tests | 2 |
| Paired Pulse Ratio analysis | 3 |
| Shapiro | 3 |
| Action Potential (AP) Analysis | 4 |
| AP Frequency t-test | 4 |
| Shapiro test | 4 |
| Wilcoxon Test | 4 |
| Summary plot | 4 |
| Individual plots | 4 |

Evoked Currents (eEPSCs) Analysis

Raw plots

Pruned individual plots

Pruned summary plots

Statistical analysis

Levene's test

Univariate Shapiro

Box's M

Multivariate normality

T-tests

Paired Pulse Ratio analysis

Shapiro

Action Potential (AP) Analysis

AP Frequency t-test

Shapiro test

Wilcoxon Test

Summary plot

Individual plots




Figure 1: Insulin decreases the excitability of DMH neurons. Data were recorded before and then after 25 minutes of insulin exposure. *A*) Insulin significantly decreases action potential frequency (mean \pm SE; n is the number of cells). *B*) Representative traces from a current injection of 40 pA (top: baseline, bottom: insulin). *C*) Insulin significantly decreases action potential amplitudes *C*) and thresholds *D*), while significantly increasing half-widths *E*). Insulin does not significantly affect latency to fire *F*), afterhyperpolarization amplitude *G*) or after-hyperpolarization time *H*). Overlay on Figures C-H: mean \pm SE. $n = 15$ for baseline and $n = 11$ for insulin because 4 cells did not fire any action potentials after insulin exposure. Wilcoxon signed-rank test, * = $p < 0.05$, ** = $p < 0.01$, *** = $p < 0.001$.