

Crosby Lab Code

by

Christelinda Laureijs

Scripts for analyzing evoked currents and action potentials in R

June 20, 2024

Contents

Action Potential (AP) Analysis	I
AP Frequency t-test	I
Shapiro test	I
Wilcoxon Test	I
Summary plot	I
Individual plots	I

Action Potential (AP) Analysis

AP Frequency t-test

Shapiro test

Wilcoxon Test

Summary plot

Individual plots




Figure 1: Insulin decreases the excitability of DMH neurons. Data were recorded before and then after 25 minutes of insulin exposure. *A*) Insulin significantly decreases action potential frequency (mean \pm SE; n is the number of cells). *B*) Representative traces from a current injection of 40 pA (top: baseline, bottom: insulin). *C*) Insulin significantly decreases action potential amplitudes *C*) and thresholds *D*), while significantly increasing half-widths *E*). Insulin does not significantly affect latency to fire *F*), after-hyperpolarization amplitude *G*) or after-hyperpolarization time *H*). Overlay on Figures C-H: mean \pm SE. $n = 15$ for baseline and $n = 11$ for insulin because 4 cells did not fire any action potentials after insulin exposure. Wilcoxon signed-rank test, * = $p < 0.05$, ** = $p < 0.01$, *** = $p < 0.001$.