

Abstract Algebra 2024–I

Homework 2

Christian Chávez

February 19, 2024

1. Determine which of the following binary operations are associative.
 - (a) the operation \star on \mathbb{Z} defined by $a \star b = a - b$
 - (b) the operation \star on \mathbb{R} defined by $a \star b = a + b + ab$
 - (c) the operation \star on \mathbb{Q} defined by $a \star b = \frac{a+b}{5}$
 - (d) the operation \star on $\mathbb{Z} \times \mathbb{Z}$ defined by $(a, b) \star (c, d) = (ad + bc, bd)$
 - (e) the operation \star on $\mathbb{Q} \setminus \{0\}$ defined by $a \star b = \frac{a}{b}$
2. Prove that addition of residue classes in $\mathbb{Z}/n\mathbb{Z}$ is associative. (Assume it is well defined.)
3. Determine which of the following sets are groups under addition:
 - (a) the set of rational numbers (including $0 = 0/1$) in lowest terms whose denominators are odd
 - (b) the set of rational numbers (including $0 = 0/1$) in lowest terms whose denominators are even
 - (c) the set of rational numbers of absolute value < 1
 - (d) the set of rational numbers of absolute value ≥ 1 together with 0
 - (e) the set of rational numbers with denominators equal to 1 or 2
 - (f) the set of rational numbers with denominators equal to 1, 2 or 3
4. Let $G = \{z \in \mathbb{C} \mid z^n = 1 \text{ for some } n \in \mathbb{Z}^+\}$.
 - (a) Prove that G is a group under multiplication (called the group of *roots of unity* in \mathbb{C}).
 - (b) Prove that G is not a group under addition.
5. Let $G = \{a + b\sqrt{2} \in \mathbb{R} \mid a, b \in \mathbb{Q}\}$.
 - (a) Prove that G is a group under addition.

- (b) Prove that the nonzero elements of G are a group under multiplication. (“Rationalize the denominators” to find multiplicative inverses.)
6. Find the orders of each element of the additive group $\mathbb{Z}/12\mathbb{Z}$.
7. Find the orders of the following elements of the multiplicative group $(\mathbb{Z}/12\mathbb{Z})^\times$:
- $$\overline{1}, \overline{-1}, \overline{5}, \overline{7}, \overline{-7}, \overline{13}.$$
8. Find the orders of the following elements of the additive group $\mathbb{Z}/36\mathbb{Z}$:
- $$\overline{1}, \overline{2}, \overline{6}, \overline{9}, \overline{10}, \overline{12}, \overline{-1}, \overline{-10}, \overline{-18}.$$
9. Let x be an element of G . Prove that $x^2 = 1$ if and only if $|x|$ is either 1 or 2.
10. Let x be an element of G . Prove that if $|x| = n$ for some positive integer n then $x^{-1} = x^{n-1}$.
11. Let x and y be elements of G . Prove that $xy = yx$ if and only if $y^{-1}xy = x$ if and only if $x^{-1}y^{-1}xy = 1$.
12. Let $x \in G$ and let $a, b \in \mathbb{Z}^+$.
- Prove that $x^{a+b} = x^a x^b$ and $(x^a)^b = x^{ab}$.
 - Prove that $(x^a)^{-1} = x^{-a}$.
 - Establish part (a) for arbitrary integers a and b (positive, negative or zero).
13. For x an element in G show that x and x^{-1} have the same order.
14. If x and g are elements of the group G , prove that $|x| = |g^{-1}xg|$. Deduce that $|ab| = |ba|$ for all $a, b \in G$.
15. Prove that if $x^2 = 1$ for all $x \in G$, then G is abelian.
16. Assume H is a nonempty subset of (G, \star) which is closed under the binary operation on G and is closed under inverses, i.e., for all h and k elements of H it holds $hk, h^{-1} \in H$. Prove that H is a group under the operation \star restricted to H (such a subset H is called a subgroup of G).
17. Prove that if x is an element of the group G then $\{x^n \mid n \in \mathbb{Z}\}$ is a subgroup (cf. the preceding exercise) of G (called the cyclic subgroup of G generated by x).
18. Compute the order of each of the elements in (a) D_6 , (b) D_8 , and (c) D_{10} .
19. Let σ be the permutation

$$1 \mapsto 3 \quad 2 \mapsto 4 \quad 3 \mapsto 5 \quad 4 \mapsto 2 \quad 5 \mapsto 1$$

and let τ be the permutation

$$1 \mapsto 5 \quad 2 \mapsto 3 \quad 3 \mapsto 2 \quad 4 \mapsto 4 \quad 5 \mapsto 1.$$

Find the cycle decompositions of each of the following permutations: $\sigma, \tau, \sigma^2, \sigma\tau, \tau\sigma$, and $\tau^2\sigma$.

20. Compute the order of each of the elements in the following in (a) S_3 and (b) S_4 .
21. Find the order of $(1\ 12\ 8\ 10\ 4)(2\ 13)(5\ 11\ 7)(6\ 9)$.
22. Write out the cycle decomposition of each element of order 4 in S_4 .
23. (a) Let σ be the 12-cycle $(1\ 2\ 3\ 4\ 5\ 6\ 7\ 8\ 9\ 10\ 11\ 12)$. For which positive integers i is σ^i also a 12-cycle?
(b) Let τ be the 8-cycle $(1\ 2\ 3\ 4\ 5\ 6\ 7\ 8)$. For which positive integers i is τ^i also an 8-cycle?
(c) Let w be the 14-cycle $(1\ 2\ 3\ 4\ 5\ 6\ 7\ 8\ 9\ 10\ 11\ 12\ 13\ 14)$. For which positive integers i is w^i also a 14-cycle?
24. Prove that if σ is the m -cycle $(a_1a_2\ldots a_m)$, then for all $i \in \{1, 2, \ldots, m\}$, it holds $\sigma^i(a_k) = a_{k+i}$, where $k+i$ is replaced by its least residue mod m when $k+i > m$. Deduce that $|\sigma| = m$.
25. Let σ be the m -cycle $(1\ 2\ 3\ \cdots\ m)$. Show that σ^i is also an m -cycle if and only if i is relatively prime to m .
26. Let p be a prime. Show that an element has order p in S_n if and only if its cycle decomposition is a product of commuting p -cycles. Show by an explicit example that this need not be the case if p is not prime.
27. Prove that the order of an element in S_n equals the least common multiple of the lengths of the cycles in its cycle decomposition. (Hint: use problem 24.)
28. Prove that $|GL_2(\mathbb{F}_2)| = 6$.
29. Write out all the elements of $GL_2(\mathbb{F}_2)$ and compute the order of each element.
30. 3. Show that $GL_2(\mathbb{F}_2)$ is non-abelian.
31. 4. Show that if n is not prime then $\mathbb{Z}/n\mathbb{Z}$ is not a field.
32. 5. Show that $GL_n(F)$ is a finite group if and only if F has a finite number of elements.
33. 6. If $|F| = q$ is finite prove that $|GL_n(F)| < q^{n^2}$.

34. 8. Show that $GL_n(F)$ is non-abelian for any $n \geq 2$ and any F .

The next exercise introduces the Heisenberg group over the field F and develops some of its basic properties. When $F = \mathbb{R}$ this group plays an important role in quantum mechanics and signal theory by giving a group theoretic interpretation (due to H. Weyl) of Heisenberg's Uncertainty Principle. Note also that the Heisenberg group may be defined more generally for example, with entries in \mathbb{Z} .

35. Let $H(F) = \left\{ \begin{pmatrix} 1 & a & b \\ 0 & 1 & c \\ 0 & 0 & 1 \end{pmatrix} \mid a, b, c \in F \right\}$ — called the Heisenberg group over F . Let $X = \begin{pmatrix} 1 & a & b \\ 0 & 1 & c \\ 0 & 0 & 1 \end{pmatrix}$ and $Y = \begin{pmatrix} 1 & d & e \\ 0 & 1 & f \\ 0 & 0 & 1 \end{pmatrix}$ be elements of $H(F)$.

Compute the matrix product XY and deduce that $H(F)$ is closed under matrix multiplication. Exhibit explicit matrices such that $XY \neq YX$ (so that $H(F)$ is always non-abelian).

36. Let G and H be groups. Let $\varphi : G \rightarrow H$ be a homomorphism.

(a) Prove that $\varphi(x^n) = \varphi(x)^n$ for all $n \in \mathbb{Z}^+$.

(b) Do part (a) for $n = -1$ and deduce that $\varphi(x^n) = \varphi(x)^n$ for all $n \in \mathbb{Z}$.

37. Let G and H be groups. If $\varphi : G \rightarrow H$ is an isomorphism, prove that $|\varphi(x)| = |x|$ for all $x \in G$. Deduce that any two isomorphic groups have the same number of elements of order n for each $n \in \mathbb{Z}^+$. Is the result true if φ is only assumed to be a homomorphism?

38. Let G and H be groups. If $\varphi : G \rightarrow H$ is an isomorphism, prove that G is abelian if and only if H is abelian. If $\varphi : G \rightarrow H$ is a homomorphism, what additional conditions on φ (if any) are sufficient to ensure that if G is abelian, then so is H ?

39. Prove that D_{24} and S_4 are not isomorphic.

40. Let A and B be groups. Prove that $A \times B \cong B \times A$.

41. Let G and H be groups and let $\varphi : G \rightarrow H$ be a homomorphism. Prove that the image of φ is a subgroup of H . Prove that, if φ is injective, then $G \cong \varphi(G)$.

42. Let G and H be groups and let $\varphi : G \rightarrow H$ be a homomorphism. Define the kernel of φ to be $\{g \in G \mid \varphi(g) = 1_H\}$ (so the kernel is the set of elements in G which map to the identity of H , i.e., is the *fiber* over the identity of H). Prove that the kernel of φ is a subgroup of G . Prove that φ is injective if and only if the kernel of φ is the identity subgroup of G .

43. Define a map $\pi : \mathbb{R}^2 \rightarrow \mathbb{R}$ by $\pi((x, y)) = x$. Prove that π is a homomorphism and find the kernel of π .
44. Let G be any group. Prove that the map from G to itself defined by $g \mapsto g^{-1}$ is a homomorphism if and only if G is abelian.
45. Let G be any group. Prove that the map from G to itself defined by $g \mapsto g^2$ is a homomorphism if and only if G is abelian.
46. Let $G = \{z \in \mathbb{C} \mid z^n = 1 \text{ for some } n \in \mathbb{Z}^+\}$. Prove that for any fixed integer $k > 1$ the map from G to itself defined by $z \mapsto z^k$ is a surjective homomorphism but is not an isomorphism.
47. Let G be a group and let $\text{Aut}(G)$ be the set of all isomorphisms from G onto G . Prove that $\text{Aut}(G)$ is a group under function composition (called the *automorphism group* of G and the elements of $\text{Aut}(G)$ are called automorphisms of G).
48. In each of (a) – (e) below prove that the specified subset is *not* a subgroup of the given group:
- (a) the set of 2-cycles in S_n for $n \geq 3$,
 - (b) the set of reflections in D_{2n} for $n \geq 3$,
 - (c) for n a composite integer > 1 and G a group containing an element of order n , the set $\{x \in G : |x| = n\} \cup \{1\}$,
 - (d) the set of (positive and negative) odd integers in \mathbb{Z} together with 0, and
 - (e) the set of real numbers whose square is a rational number (under addition).
49. Show that the following subsets of the dihedral group D_8 are actually subgroups: (a) $\{1, r^2, s, sr^2\}$, (b) $\{1, r^2, sr, sr^3\}$.
50. Give an explicit example of a group G and an infinite subset H of G that is closed under the group operation but is not a subgroup of G .
51. Prove that G cannot have a subgroup H with $|H| = n - 1$, where $n = |G| > 2$.
52. Let G be an abelian group. Prove that $\{g \in G : |g| < \infty\}$ is a subgroup of G (called the *torsion subgroup* of G). Give an explicit example where this set is not a subgroup when G is non-abelian.
53. Fix some $n \in \mathbb{Z}$ with $n > 1$. Find the torsion subgroup (cf. the previous exercise) of $\mathbb{Z} \times (\mathbb{Z}/n\mathbb{Z})$. Show that the set of elements of infinite order together with the identity is *not* a subgroup of this direct product.
54. Let H and K be subgroups of G . Prove that $H \cup K$ is a subgroup if and only if either $H \subseteq K$ or $K \subseteq H$.

55. Let $G = GL_n(F)$, where F is any field. Define

$$SL_n(F) = \{A \in GL_n(F) \mid \det(A) = 1\}$$

(called the *special linear group*). Prove that $SL_n(F) \leq GL_n(F)$.

56. (a) Prove that if H and K are subgroups of G then so is their intersection $H \cap K$.

(b) Prove that the intersection of an arbitrary nonempty collection of subgroups of G is again a subgroup of G (do not assume the collection is countable).

57. Let A and B be groups. Prove that the following sets are subgroups of the direct product $A \times B$:

(a) $\{(a, 1) \mid a \in A\}$,

(b) $\{(1, b) \mid b \in B\}$, and

(c) $\{(a, a) \mid a \in A\}$, where we assume $A = B$.

58. Let $H_1 \leq H_2 \leq \cdots$ be an ascending chain of subgroups of G . Prove that $\cup_{i=1}^{\infty} H_i$ is a subgroup of G .

59. Let $n \in \mathbb{Z}^+$ and let F be a field. Prove that the set $\{(a_{ij}) \in GL_n(F) \mid a_{ij} = 0 \text{ for all } i > j\}$ is a subgroup of $GL_n(F)$ (called the *group of upper triangular matrices*).

60. Prove that $C_G(A) = \{g \in G \mid g^{-1}ag = a \text{ for all } a \in A\}$.

61. Prove that $C_G(Z(G)) = G$ and deduce that $N_G(Z(G)) = G$.

62. In each of parts (a) to (c) show that for the specified group G and subgroup A of G , $C_G(A) = A$ and $N_G(A) = G$.

(a) $G = S_3$ and $A = \{1, (123), (132)\}$

(b) $G = D_8$ and $A = \{1, s, r^2, sr^2\}$

(c) $G = D_{10}$ and $A = \{1, r, r^2, r^3, r^4\}$

63. Let H be a subgroup of the group G .

(a) Show that $H \leq N_G(H)$. Give an example to show that this is not necessarily true if H is not a subgroup.

(b) Show that $H \leq C_G(H)$ if and only if H is abelian.

64. Prove that $Z(G) \leq N_G(A)$ for any subset A of G .