

The interleaving distance between two interval modules

Christian Chavez

I found nowhere the explicit computation of the interleaving distance between interval modules, though it is well known that it equals the bottleneck distance.

1 Preliminaries

Let $\mathbf{k}$ be an arbitrary field, fixed throughout. We denote by $\mathbb{R}$ either the real line or the category induced by the poset $(\mathbb{R}, \leq)$. All functors are covariant.

1.1 The category of persistence modules

A **persistence module** is a functor $\mathbb{R} \rightarrow \text{Vect}(\mathbf{k})$.

Example 1 (Important). Let $I \subseteq \mathbb{R}$ be an interval. Define $\mathbf{k}^I: \mathbb{R} \rightarrow \text{Vect}(\mathbf{k})$ by

$$t \mapsto \begin{cases} \mathbf{k} & \text{if } t \in I \\ \mathbf{0} & \text{else} \end{cases} \quad \text{and} \quad r \rightarrow s \mapsto \begin{cases} \mathbf{k} \xrightarrow{\text{Id}} \mathbf{k} & \text{if } r, s \in I \\ \mathbf{0} & \text{else} \end{cases}$$

on objects and morphisms, respectively. Here $\mathbf{0}$ denotes both the zero vector space and the zero map, respectively. We call $\mathbf{k}^I$ the **interval module** over I .

Remark. Interval modules are the building blocks of persistence modules. In fact, one objective of persistence theory is to understand persistence modules by decomposing them into interval modules. However, this is not always possible. A situation when such a decomposition is guaranteed is when we work with finite dimensional vector spaces, that is, if we work over $\text{vect}(\mathbf{k})$ instead of $\text{Vect}(\mathbf{k})$.

The concise definition of persistence module can be expanded to aide comprehension. Let $V: \mathbb{R} \rightarrow \text{Vect}(\mathbf{k})$ be a persistence module.

- For each $t \in \mathbb{R}$ and each arrow $r \rightarrow s$ denote $V(t) = V_t$ and $V(r \rightarrow s) = v_s^r$.
- Since V is a functor, we have the so-called **composition law**:

$$v_t^s \circ v_s^r = v_t^r \quad \text{whenever } r \leq s \leq t$$

$$\text{and } V(t \rightarrow t) = v_t^t = \text{Id}_{V_t}.$$

Thus, a persistence module V is a family of vector spaces $(V_t)_{t \in \mathbb{R}}$ together with a family $(v_t^s: V_s \rightarrow V_t \mid s \leq t)_{s,t \in \mathbb{R}}$ of $\mathbf{k}$ -linear maps that satisfy the composition law.

We got a set theoretical definition from the categorical one above. In fact, the two are equivalent. We will use the first, but prefer the notation of the second.

Since persistence modules are functors, we can consider the natural transformations between them, and since those can be composed (using vertical composition) we obtain a category.

Definition 1. The **category of persistence modules** is $\mathbf{X} = \text{Fun}(\mathbb{R}, \text{Vect}(\mathbf{k}))$.

Yes, this category is fun. Concretely, a morphism of persistence modules $\alpha: U \rightarrow V$ is a family of linear maps $(\alpha_r: U_r \rightarrow V_r)_{r \in \mathbb{R}}$ such that the diagram

$$\begin{array}{ccc} U_s & \xrightarrow{v_t^s} & U_t \\ \alpha_s \downarrow & & \downarrow \alpha_t \\ V_s & \xrightarrow{v_t^s} & V_t \end{array} \quad (1)$$

commutes for all $s \leq t$, i.e., $v_t^s \circ \alpha_s = \alpha_t \circ v_t^s$.

Explain composition with a diagram.

Example 2. If U and V are any persistence modules, then there is a zero morphism $U \rightarrow V$, given by $(U_r \xrightarrow{0} V_r)_{r \in \mathbb{R}}$, and denoted by $\mathbf{0}$.

Example 3. Consider two interval modules $\mathbf{k}^{[0,1]}$ and $\mathbf{k}^{[0,2]}$. For each $r \in \mathbb{R}$, define

$$\alpha_r = \mathbf{k}^{[0,1]}(x) \rightarrow \mathbf{k}^{[0,2]}(x)$$

So $\alpha = (\alpha_r)_{r \in \mathbb{R}}$ is the inclusion $\mathbf{k}^{[0,1]} \hookrightarrow \mathbf{k}^{[0,2]}$. However, α is not a morphism of persistence modules. For example, for the arrow $1 \rightarrow 2$ we have

$$\mathbf{k}^{[0,1]}(1 \rightarrow 2) = \mathbf{k} \xrightarrow{0} \mathbf{0}$$

and

$$\mathbf{k}^{[0,2]}(1 \rightarrow 2) = \mathbf{k} \xrightarrow{\text{Id}} \mathbf{k}$$

but $\alpha_1 = \mathbf{k} \xrightarrow{\text{Id}} \mathbf{k}$ and $\alpha_2 = \mathbf{0} \xrightarrow{0} \mathbf{k}$. Therefore, the diagram

$$\begin{array}{ccc} \mathbf{k}^{[0,1]} & \xrightarrow{1 \rightarrow 2} & \mathbf{k}^{[0,2]} \\ \alpha_1 \downarrow & & \downarrow \alpha_2 \\ \mathbf{k} & \xrightarrow{1 \rightarrow 2} & \mathbf{k} \end{array}$$

does not commute. The same situation happens if we'd taken $r \rightarrow s$ with $r \in [0, 1] \cap [0, 2] = [0, 1]$ and $s \in [0, 2] \setminus [0, 1] = (1, 2]$.

1.2 The shift functor

Let U be a persistence module and fix any $t \in \mathbb{R}$. Define a new persistence module $U[t]$ by

$$U[t](r) = U_{r+t} \quad \text{and} \quad U[t](r \rightarrow s) = u_{r+t}^{s+t}.$$

We say that $U[t]$ is a **shifted module**. Note that

$$U[0] = U \quad \text{and} \quad U[t][t'] = U[t + t']$$

for any $t, t' \in \mathbb{R}$. Similarly, for any morphism of persistence modules $\alpha: U \rightarrow V$, we can define a **shifted module morphism** $\alpha[t] = (\alpha_{r+t})_{r \in \mathbb{R}}$.

Suppose we've made two shifts of U , say $U[s]$ and $U[t]$, where $s \leq t$. We might ask ourselves: *what is the relationship between these shifted modules?* In other words, we wonder if there is a canonical natural transformation $\alpha: U[s] \rightarrow U[t]$. To find out, consider any $p \rightarrow q$ and apply both $U[s]$ and $U[t]$ to this arrow. We get

$$U[s](p \rightarrow q) = U_{p+s} \xrightarrow{u_{q+s}^{p+s}} U_{q+s}$$

and

$$U[t](p \rightarrow q) = U_{p+t} \xrightarrow{u_{q+t}^{p+t}} U_{q+t}.$$

Since $\alpha = (\alpha_r)_{r \in \mathbb{R}}$ is determined by its components, we need to find the vertical arrows that make the square

![[diagram3]](/assets/2025-10-29-images-interleaving-distance-between-interval-modules/diagram3)

commute. Set $\alpha_p = u_{p+t}^{p+s}$ and $\alpha_q = u_{q+t}^{q+s}$. Then the composition law give us

$$u_{q+t}^{p+t} \circ u_{p+t}^{p+s} = u_{q+t}^{p+s} = u_{q+t}^{q+s} \circ u_{q+s}^{p+s},$$

which means the square commutes. So, yes there is a canonical way to map $U[s]$ to $U[t]$. Denote $\alpha = U[s \rightarrow t]$ and define

$$U[s] \xrightarrow{U[s \rightarrow t]} U[t] = (u_{r+t}^{r+s})_{r \in \mathbb{R}}.$$

Remark 1.

- For $t \geq 0$, we denote $\mathbb{K}_U^t = U[0 \rightarrow t]$.
- Observe that

$$\mathbb{K}_U^0 = U[0 \rightarrow 0] = (u_r^r)_{r \in \mathbb{R}} = (\text{Id}_{U_r})_{r \in \mathbb{R}} = \mathbb{K}_U$$

is an identity natural transformation. So, for $t \geq 0$, we might think of $\mathbb{K}_U^t$ as a generalized identity.

- By definition, we get $U[t \rightarrow t] = \mathbb{K}_{U[t]}$ for any $t \in \mathbb{R}$. Moreover, if $r \leq s \leq t$, for any $p \in \mathbb{R}$ we have

$$\begin{aligned} (U[s \rightarrow t] \circ U[r \rightarrow s])_p &= U[s \rightarrow t]_p \circ U[r \rightarrow s]_p \\ &= u_{p+t}^{p+s} \circ u_{p+s}^{p+r} \\ &= u_{p+t}^{p+r} \\ &= U[r \rightarrow t]_p \end{aligned}$$

thanks to the composition law. Thus

$$U[s \rightarrow t] \circ U[r \rightarrow s] = U[r \rightarrow t].$$

The last remark leads us to the following definition.

Definition 2. The **shift functor** $T = [\cdot]: \mathbb{R} \rightarrow \text{End}(X)$ is defined as follows

- (i) Objects: for each $t \in \mathbb{R}$ we have a functor $T(t): X \rightarrow X$ denoted T_t , which is defined by $T_t(U) = U[t]$ on objects $U \in X$, and by $T_t(\alpha) = \alpha[t] = (\alpha_{r+t})_{r \in \mathbb{R}}$ on morphisms $\alpha = (\alpha_r)_{r \in \mathbb{R}}$. (Equivalently, we can define $T_t(\alpha) = \alpha \star [t]$, using a whiskering)
- (ii) Morphisms: for each $r \rightarrow s$, define a natural transformation $T(r \rightarrow s): T_r \rightarrow T_s$ by

$$T(r \rightarrow s) = (U[r \rightarrow s])_{U \in X}.$$

Example 4. Let $\mathcal{I} = \mathbf{k}^I$ be an interval module and fix $t \in \mathbb{R}$. Then, for any $r \in \mathbb{R}$,

$$\mathbf{k}^I(r+t) = \begin{cases} \mathbf{k} & \text{if } r+t \in I \\ \mathbf{0} & \text{else} \end{cases} = \begin{cases} \mathbf{k} & \text{if } r \in I-t \\ \mathbf{0} & \text{else} \end{cases} = \mathbf{k}^{I-t}(r)$$

and similarly, for any $r \leq s$ we obtain

$$\mathbf{k}^I(r+t \rightarrow s+t) = \mathbf{k}^{I-t}(r \rightarrow s).$$

Thus $\mathbf{k}^I[t] = \mathbf{k}^{I-t}$. We will denote $\mathcal{I}[t] = \mathcal{I} - t$.

1.3 The interleaving distance between persistence modules

Let U and V be persistence modules. A **t -interleaving** of U and V is a pair of natural transformations $(\phi: U \rightarrow V[t], \psi: V \rightarrow U[t])$ that make the diagrams

![[diagram4]](/assets/2025-10-29-images-interleaving-distance-between-interval-modules/diagram4)

commute. If such a pair exists, we say that U and V are **t -interleaved**.

Example 5. Two persistence modules are isomorphic if and only if they are 0-interleaved. Thus, we can think of interleavings as generalized isomorphisms. Let us see why...

Definition 3. The **interleaving distance** between two persistence modules U and V is

$$d_T(U, V) = \inf \{t \geq 0 \mid \text{there exists a } t\text{-interleaving of } U \text{ and } V\}.$$

in case there is at least one interleaving of U and V . If no such interleaving exists, we set $d_T(U, V) = \infty$.

The subscript T remind us that d_T depends on the shift functor. On the other hand, d_T is not an actual metric, but a *pseudometric*, since $d_T(U, V) = 0$ does not imply $U = V$.

Example 6. An example that proves the last sentence.


Let us prove d_T satisfies the triangle inequality.

2 The computation

We'll do the computation of $d_T(\mathbf{k}^I, \mathbf{k}^J)$ for the case of closed intervals. Let $I = [a, b]$ and $J = [c, d]$. Denote by $\mathcal{I} = \mathbf{k}^I$ and $\mathcal{J} = \mathbf{k}^J$ the associated interval modules.

Our strategy is to find some appropriate bounds for $d_T(U, V)$. Before going into the search for the pair of morphisms that would make the triangles above commute, we need to know how are the morphisms between arbitrary interval modules, because then we can characterize the morphisms between shifted interval modules.

Let $\alpha: \mathcal{I} \rightarrow \mathcal{J}$ be any persistence module morphism. For each $s \in \mathbb{R}$ we have a map of vector spaces $\alpha_s: \mathcal{I}_s \rightarrow \mathcal{J}_s$. The only available options for $\mathcal{I}_s$ and $\mathcal{J}_s$ are either $\mathbf{k}$ or $\mathbf{0}$.




Note that, if $s \notin I \cap J$, then either $\mathcal{I}_s = \mathbf{0}$ or $\mathcal{J}_s = \mathbf{0}$, and in any case α_s is the zero map. Thus, we always get trivial morphisms outside of the intersection. Inside the intersection, when $s \in I \cap J$, we have a linear map $\alpha_s: \mathbf{k} \rightarrow \mathbf{k}$, but since $\mathbf{k}$ is a field, α_s is nothing more than multiplication by some scalar $\lambda \in \mathbf{k}$, so $\alpha_s: x \mapsto \lambda x$. Such λ could be zero, though.

As a first result, we have the following.

Fact 1. If $I \cap J = \emptyset$, then $\text{Hom}(\mathbf{k}^I, \mathbf{k}^J) = \mathbf{0}$.

With this trivial case out of the way, assume $I \cap J \neq \emptyset$, and let us consider two cases: $a < c$ or $c \leq a$.

Case 1. Suppose $a < c$. Recall that our goal is to characterize the morphisms $\mathbf{k}^I \rightarrow \mathbf{k}^J$. Since we already know what happens when $t \notin I \cap J$, let us focus on $\alpha_t: \mathbf{k}_t^I \rightarrow \mathbf{k}_t^J$ with $t \in I \cap J$, that is with $c \leq t \leq b$.



Take the arrow $a \rightarrow t$ and apply both $\mathbf{k}^I$ and $\mathbf{k}^J$. We have $\mathbf{k}^I(a \rightarrow t) = \mathbf{k} \xrightarrow{\text{Id}} \mathbf{k}$ and $\mathbf{k}^J(a \rightarrow t) = \mathbf{0} \xrightarrow{0} \mathbf{k}$. Note that $\alpha_a: \mathbf{k} \rightarrow \mathbf{0}$ is the zero map and $\alpha_t: \mathbf{k} \rightarrow \mathbf{k}$ is some linear map. By (1), naturality implies

$$\alpha_t = \alpha_t \circ \mathbf{k}^I(a \rightarrow t) = \mathbf{k}^J(a \rightarrow t) \circ \alpha_a = 0.$$

Thus α_t is zero even on $I \cap J$. The conclusion is that $\text{Hom}(\mathbf{k}^I, \mathbf{k}^J) \cong \mathbf{0}$ if I was *born* before J .

Case 2. Suppose $c \leq a$, meaning that J was *born* before I . Now we consider two more cases.

Case 2.1. Suppose $b < d$. This means that J *died* after I . We also get $\text{Hom}(\mathbf{k}^I, \mathbf{k}^J) \cong \mathbf{0}$.

Case 2.2. Suppose $d \leq b$. Here we prove there is an isomorphism $\text{Hom}(\mathbf{k}^I, \mathbf{k}^J) \cong \mathbf{k}$.

Now we come to the following conclusion.

Fact 2. If $c \leq a \leq d \leq b$, then $\text{Hom}(\mathbf{k}^I, \mathbf{k}^J) \cong \mathbf{k}$. Given $\lambda \in \mathbf{k}$, define $\alpha^\lambda: \mathbf{k}^I \rightarrow \mathbf{k}^J$ by

$$\alpha_r^\lambda = \begin{cases} \mathbf{k} \xrightarrow{\mu_\lambda} \mathbf{k} & \text{if } r \in I \cap J \\ \mathbf{k}_r^I \xrightarrow{0} \mathbf{k}_r^J & \text{else} \end{cases}$$

where $\mu_\lambda: r \mapsto \lambda r$. The map

$$\Psi: \mathbf{k} \rightarrow \text{Hom}(\mathbf{k}^I, \mathbf{k}^J) : \lambda \mapsto \alpha^\lambda$$

is injective, and surjective, and linear (proof...)

Both Fact 1 and Fact 2 will help us determine $d_T(\mathbf{k}^I, \mathbf{k}^J)$ for any closed intervals I and J .

2.1 Bound above


We start looking for values of $t \geq 0$ for which a t -interleaving of $\mathcal{I} = \mathbf{k}^I$ and $\mathcal{J} = \mathbf{k}^J$ exists, that is, for which we can find maps $\phi: \mathcal{I} \rightarrow \mathcal{J}[t]$ and $\psi: \mathcal{J} \rightarrow \mathcal{I}[t]$ that make the diagrams

![[diagram5]](/assets/2025-10-29-images-interleaving-distance-between-interval-modules/diagram5.png)

commute. By using the fact that a shift of an interval module is the interval module of the shifted interval (Example 4), we can redraw these diagram as

![[diagram6]](/assets/2025-10-29-images-interleaving-distance-between-interval-modules/diagram6.png)

The first observation is that if the top arrows are both 0, then we can put $\phi = 0$ and $\psi = 0$ because then the triangles commute trivially. But, how do we get the top arrows to be zero? Here we can use Fact 1: if $[a, b] \cap [a - 2t, b - 2t] = \emptyset$, then $\text{Hom}(\mathbf{k}^{[a, b]}, \mathbf{k}^{[a - 2t, b - 2t]}) = \mathbf{0}$. In order to get an empty intersection, we just need to make a sufficiently large shift. Since $t \geq 0$, the shift moves to the left, so we need $b - 2t < a$, that is $t > \frac{b-a}{2}$.



By the same reason we arrive at $t > \frac{d-c}{2}$. Thus the top arrows are both zero if $t > \frac{b-a}{2}$ and $t > \frac{d-c}{2}$, which is equivalent to

$$t > \frac{1}{2} \max \{b - a, d - c\}.$$

If we denote the right term by M , we get that an $(M + \varepsilon)$ -interleaving of $\mathcal{I}$ and $\mathcal{J}$ exists for every $\varepsilon > 0$. Thus $d_T(\mathbf{k}^I, \mathbf{k}^J) \leq M + \varepsilon$ for every $\varepsilon > 0$, and taking the infimum over ε give us

$$d_T(\mathbf{k}^{[a, b]}, \mathbf{k}^{[c, d]}) \leq \frac{1}{2} \max \{b - a, d - c\}.$$

Notice that this bound does not take into account the relationship (or position) between I and J , so our next step is to find a bound that does consider this aspect.

Now we use Fact 2. Framed in our context, it tell us that

$$\begin{aligned} c - t \leq a \leq d - t \leq b &\implies \text{Hom}(\mathbf{k}^{[a, b]}, \mathbf{k}^{[c-t, d-t]}) \cong \mathbf{k} \\ a - t \leq c \leq b - t \leq d &\implies \text{Hom}(\mathbf{k}^{[c, d]}, \mathbf{k}^{[a-t, b-t]}) \cong \mathbf{k}. \end{aligned} \tag{2}$$

Now we start looking for values of $t \geq 0$ for which the two sets of inequalities are satisfied, because then we can choose appropriate nonzero ϕ and ψ such that (ϕ, ψ) is a t -interleaving.

The conditions $c - t \leq a$ and $a - t \leq c$ are equivalent to $t \geq |c - a|$, and the conditions $d - t \leq b$ and $b - t \leq d$ are equivalent to $t \geq |d - b|$. So, let $t \geq \max\{|c - a|, |d - b|\}$.

With respect to the middle inequalities, we consider two cases.

Case A. Suppose $d - t < a$ and $b - t < c$. Using the fact that $t \geq |d - b| \geq b - d$, we have

$$b - 2t = b - t - t \leq b - (b - d) - t \leq d - t < a$$

whence $t > \frac{b-a}{2}$. Similarly, if $b - t < c$, we get $t > \frac{d-c}{2}$. Thus $t \geq \max\{\frac{b-a}{2}, \frac{d-c}{2}\}$, which lead us to the situation above where a t -interleaving trivially exists, but the two hom-sets are zero.

Case B. Suppose then that $a \leq d - t$ or $c \leq b - t$. Using the assumption that $t \geq \max\{|c - a|, |d - b|\}$, we see that either inequality implies the other. Thus we have $a \leq d - t$ and $c \leq b - t$. Therefore both sets of inequalities in (2) above are satisfied, whence

$$\text{Hom}(\mathbf{k}^{[a,b]}, \mathbf{k}^{[c-t, d-t]}) \cong \mathbf{k} \quad \text{and} \quad \text{Hom}(\mathbf{k}^{[c,d]}, \mathbf{k}^{[a-t, b-t]}) \cong \mathbf{k}.$$

Using Fact 2, we see that this implies that for $1 \in \mathbf{k}$, there exists $\phi: \mathbf{k}^{[a,b]} \rightarrow \mathbf{k}^{[c-t, d-t]}$ and $\psi: \mathbf{k}^{[c,d]} \rightarrow \mathbf{k}^{[a-t, b-t]}$ defined by

$$\phi_r = \begin{cases} \mathbf{k} \xrightarrow{\text{Id}} \mathbf{k} & \text{if } r \in [a, d - t] \\ \mathbf{0} & \text{else} \end{cases}$$


and

$$\psi_r = \begin{cases} \mathbf{k} \xrightarrow{\text{Id}} \mathbf{k} & \text{if } r \in [c, b - t] \\ \mathbf{0} & \text{else} \end{cases}$$

The final, and most important, step is to prove that (ϕ, ψ) is a t -interleaving of $\mathbf{k}^I$ and $\mathbf{k}^J$. This amounts to prove that the triangles

!

commute for every $r \in \mathbb{R}$. Let's consider the first triangle. We consider three cases.



If $r \in (-\infty, a)$, then

$$\begin{aligned}\psi_{r+t} \circ \phi_r &= \psi_{r+t} \circ (\mathbf{0} \xrightarrow{0} \mathbf{k}^*) \\ &= \mathbf{0} \xrightarrow{0} \mathbf{k}^* \\ &= \mathbf{k}^{[a,b]}(r \rightarrow r + 2t)\end{aligned}$$

no matter the value of ψ_{r+t} . Here $\mathbf{k}^*$ denotes either $\mathbf{0}$ or $\mathbf{k}$.

If $r \in [a, b - 2t]$, then

$$\begin{aligned}\psi_{r+t} \circ \phi_r &= (\mathbf{k} \xrightarrow{\text{Id}} \mathbf{k}) \circ (\mathbf{k} \xrightarrow{\text{Id}} \mathbf{k}) \\ &= \mathbf{k} \xrightarrow{\text{Id}} \mathbf{k} \\ &= \mathbf{k}^{[a,b]}(r \rightarrow r + 2t).\end{aligned}$$

Finally, if $r \in (b - 2t, \infty)$,

$$\begin{aligned}\psi_{r+t} \circ \phi_r &= (\mathbf{k}^* \xrightarrow{0} \mathbf{0}) \circ \phi_r \\ &= \mathbf{k}^* \xrightarrow{0} \mathbf{0} \\ &= \mathbf{k}^{[a,b]}(r \rightarrow r + 2t).\end{aligned}$$

where $\mathbf{k}^*$ denotes either $\mathbf{0}$ or $\mathbf{k}$. We have proved that the left triangle commutes for every $r \in \mathbb{R}$. In an entirely analogous manner we prove the right triangle commutes. Therefore (ϕ, ψ) is a t -interleaving of $\mathbf{k}^I$ and $\mathbf{k}^J$.

We have proved that, in either case A or B, $\mathbf{k}^I$ and $\mathbf{k}^J$ are t -interleaved if $t \geq \max\{|c - a|, |d - b|\}$. Therefore

$$d_T(\mathbf{k}^{[a,b]}, \mathbf{k}^{[c,d]}) \leq \max\{|c - a|, |d - b|\}.$$

Up to now we've found two upper bounds for $d_T(\mathbf{k}^{[a,b]}, \mathbf{k}^{[c,d]})$, so we know that the distance is smaller than both of them. We summarize what we've found in the following

Fact 3. For any closed intervals $I = [a, b]$ and $J = [c, d]$,

$$d_T(\mathbf{k}^{[a,b]}, \mathbf{k}^{[c,d]}) \leq \min \left\{ \frac{1}{2} \max \{b - a, d - c\}, \max \{|c - a|, |d - b|\} \right\}. \quad (3)$$

2.2 Bound below

Let's prove that (3) is actually an equality. Denote that minimum by m . Our goal is to prove that if a t -interleaving exists, then necessarily $t \geq m$, for then we can take the infimum over all those $t \geq 0$ and get that $d_T(\mathbf{k}^{[a,b]}, \mathbf{k}^{[c,d]}) \geq m$.

Let $t \geq 0$. We will actually prove that if $t < m$, then no interleaving can exist.