

UNIVERSITÀ DEGLI STUDI DI TORINO

Corso di Laurea Magistrale in Matematica



UNIVERSITÀ  
DI TORINO

Tesi di Laurea Magistrale

## The ideally exact categories

A non-pointed generalization  
of the notion of semiabelian category

**Relatore**

professor Alan Cigoli

**Candidato**

Christian D'Ascoli

Anno Accademico 2024-2025

# Table of Contents

|                                                                     |           |
|---------------------------------------------------------------------|-----------|
| <b>Introduction</b>                                                 | <b>1</b>  |
| <b>1 Preliminaries</b>                                              | <b>2</b>  |
| <b>2 Pullback functor</b>                                           | <b>5</b>  |
| 2.1 Connected limits . . . . .                                      | 5         |
| 2.2 The pullback functor and its adjunction . . . . .               | 7         |
| 2.3 Nullary and essentially nullary monads . . . . .                | 15        |
| <b>3 Ideally exact categories</b>                                   | <b>17</b> |
| 3.1 Preliminary results on the Eilenberg-Moore categories . . . . . | 17        |
| 3.2 The notion of ideally exact category . . . . .                  | 28        |
| <b>Appendix</b>                                                     | <b>32</b> |



# Introduction

# Chapter 1

## Preliminaries

In order to understand the notions introduced in this thesis is important to recall some definitions and constructions. We also take this opportunity to introduce and fix some notations.

First we define some categories that will be used throughout the thesis. Let  $\mathcal{A}$  be a category, we defines the following categories.

- The **slice category**: given an object  $B$  in a category  $\mathcal{A}$ , we call  $(\mathcal{A} \downarrow B)$  the category of objects over  $E$ , we denote an object by  $(A, \alpha : A \rightarrow B)$ . A morphism between two objects  $(A, \alpha)$  and  $(A', \alpha')$  is a morphism  $f : A \rightarrow A'$  such that  $\alpha' f = \alpha$ .

$$\begin{array}{ccc} A & & A \xrightarrow{f} A' \\ \downarrow \alpha & & \downarrow \alpha \swarrow \alpha' \\ B & & B \end{array}$$

- The **coslice category**: given an object  $B$  in a category  $\mathcal{A}$ , we call  $(B \downarrow \mathcal{A})$  the category of objects under  $B$ , we denote an object by  $(\alpha : B \rightarrow A, A)$ . A morphism between two objects  $(\alpha, A)$  and  $(\alpha', A')$  is a morphism  $f : A \rightarrow A'$  such that  $f\alpha = \alpha'$ .

$$\begin{array}{ccc} B & & B \\ \downarrow \alpha & & \downarrow \alpha \searrow \alpha' \\ A & & A \xrightarrow{f} A' \end{array}$$

- The **category of points**: given an object  $B$  in a category  $\mathcal{A}$ , we call  $\text{Pt}(B)$  the category of points of  $B$ , we denote an object by  $(A, B, p, s)$  which is a diagram of the type  $A \begin{smallmatrix} \xrightarrow{p} \\ \xleftarrow{s} \end{smallmatrix} B$  with the condition  $ps = 1_B$ . Morphisms between two objects  $(A, B, p, s)$  and  $(A', B', p', s')$  are diagrams of the type

$$\begin{array}{ccc} A & \xrightarrow{f} & A' \\ \swarrow p & & \swarrow p' \\ & B & \\ \searrow s & & \searrow s' \end{array}$$

with the conditions  $p'f = p$  and  $fs = s'$

We recall the following definitions.

**Definition 1.0.1.** A finitely complete category  $\mathcal{A}$  is said to be **regular** if


- Every morphism in  $\mathcal{A}$  admits an (regular epi, mono) factorization;
- Regular epis in  $\mathcal{A}$  are pullback stable.

**Definition 1.0.2.** A category  $\mathcal{A}$  is said to be **exact** if

- It is regular;
- Any equivalence relation is the kernel pair of some arrow.

In order to define the next notion, we need the following construction. Suppose  $\mathcal{A}$  has pullbacks and consider a morphism  $f : B' \rightarrow B$  in the same category. We define the **change-of-base functor** of  $f$ .

$$f^* : \text{Pt}(B) \rightarrow \text{Pt}(B')$$



where  $\langle 1_{B'}, sf \rangle$  is the section of  $\text{proj}_1$ . It holds that  $(\text{proj}_1)\langle 1_{B'}, sf \rangle = 1_{B'}$ . The action on a morphism  $\alpha : A \rightarrow A'$  is given by



**Definition 1.0.3.** A finitely complete category  $\mathcal{A}$  is said to be **protomodular** if for every morphism  $f : A \rightarrow B$  in  $\mathcal{A}$ , the change-of-base functor  $f^* : \text{Pt}(B) \rightarrow \text{Pt}(A)$  is conservative.

We conclude recalling the following.

**Definition 1.0.4.** A category  $\mathcal{A}$  is said to be **semiabelian** if


- It is pointed;
- It is exact;
- It is protomodular;
- It has binary coproducts.

# Chapter 2

## Pullback functor

### 2.1 Connected limits


**Definition 2.1.1.** A **connected limit** is a limit over a connected category. A category is connected if it is inhabited and any two objects can be connected by a zigzag of morphisms




where each symbol  $\leftrightarrow$  is a morphism  $y_i \rightarrow y_{i+1}$  or  $y_{i+1} \rightarrow y_i$ .

**Example 2.1.2.** Some examples

- The pullback is a connected limit.




- The equalizer is a connected limit.



**Remark 2.1.3.** The connected limits in the category  $(\mathcal{A} \downarrow E)$  can be computed in  $\mathcal{A}$ . In fact we can provide to the object  $L$  of the limit cone  $(L, (q_i)_{i \in \mathcal{I}})$  in  $\mathcal{A}$  a morphism to form an object in  $(\mathcal{A} \downarrow E)$ . In the zigzag below we chose  $(\varphi_1)(q_1)$ ,

but we could use whatever morphism of the type  $(\varphi_i)(q_i)$ . The connection guarantees that all the morphisms  $q_1 \dots q_n$  involved in the limit in  $\mathcal{A}$  are also morphisms in  $(\mathcal{A} \downarrow E)$ , ie  $(\varphi_i)(q_i) = (\varphi_1)(q_1) \forall i$ .



By the previous result, the following holds.

**Proposition 2.1.4.** *Let  $E$  be an object in  $\mathcal{A}$ . Then the forgetful functor  $U : (\mathcal{A} \downarrow E) \rightarrow \mathcal{A}$  creates all connected limits that  $\mathcal{A}$  has.*

Note that this result is not true for all the limits. This is extensively shown in the following example.

**Example 2.1.5.** *The product is not a connected limit, in fact as we can observe the following category is not connected*

1                      2

We try to compute the product directly in  $(\mathcal{A} \downarrow E)$ . Let  $(D, \delta)$  and  $(D', \delta')$  be two objects in  $(\mathcal{A} \downarrow E)$ , the product is given by the following pullback

$$\begin{array}{ccccc}
 & D \times_B D' & \longrightarrow & D' & \\
 & \swarrow & & \searrow \delta' & \\
 D & \xrightarrow{\delta} & E & & \\
 & \searrow \delta & \downarrow 1_E & \swarrow \delta' & \\
 & & E & & 
 \end{array}$$

which doesn't correspond to the product  $D \times D'$  in  $\mathcal{A}$ .

## 2.2 The pullback functor and its adjunction

Given a category  $\mathcal{A}$  with pullbacks and a morphism  $p : E \rightarrow B$  in  $\mathcal{A}$ , one can consider functors pullback and shrink

$$\begin{array}{ccc}
 & \xrightarrow{p^*} & \\
 (\mathcal{A} \downarrow B) & & (\mathcal{A} \downarrow E) \\
 & \xleftarrow{p_!} &
 \end{array}$$

defined as follows:

- $p^* : (\mathcal{A} \downarrow B) \rightarrow (\mathcal{A} \downarrow E)$  is the **pullback functor**

$$\begin{array}{ccc}
 \begin{array}{c} A \\ \downarrow \alpha \\ B \end{array} & \xrightarrow{p^*} & \begin{array}{ccc} E \times_B A & \xrightarrow{\text{proj}_2} & A \\ \downarrow \text{proj}_1 & & \downarrow \alpha \\ E & \xrightarrow{p} & B \end{array} \\
 \\ 
 \begin{array}{ccc} A & \xrightarrow{f} & A' \\ \downarrow \alpha & \nearrow \alpha' & \\ B & & \end{array} & \xrightarrow{p^*} & \begin{array}{c} \begin{array}{ccccc} E \times_B A & \xrightarrow{\text{proj}_2} & A & \xrightarrow{f} & A' & \xleftarrow{\text{proj}_2'} & E \times_B A' \\ & \searrow \alpha & & \swarrow \alpha' & & \\ & & B & & & \\ \text{proj}_1 \swarrow & & \uparrow p & & \searrow \text{proj}_1' & \\ & E & & & & \end{array} \\ \text{dashed arc: } p^*(f) = \tilde{f} \end{array}
 \end{array}$$

- $p_! : (\mathcal{A} \downarrow E) \rightarrow (\mathcal{A} \downarrow B)$  is the **shrink functor**

$$\begin{array}{ccc}
 \begin{array}{c} D \\ \downarrow \delta \\ E \end{array} & \xrightarrow{p_!} & \begin{array}{ccc} D & & \\ \delta \downarrow & \searrow p\delta & \\ E & \xrightarrow{p} & B \end{array} \\
 \\ 
 \begin{array}{ccc} D & \xrightarrow{g} & D' \\ \delta \downarrow & \nearrow \delta' & \\ E & & \end{array} & \xrightarrow{p_!} & \begin{array}{ccc} D & \xrightarrow{g} & D' \\ & \searrow p\delta & \downarrow p\delta' \\ E & \xrightarrow{p} & B \end{array}
 \end{array}$$

**Proposition 2.2.1.** *Let  $p : E \rightarrow B$  be a morphism in  $\mathcal{A}$  with pullbacks, then  $p_!$  preserves connected limits.*

*Proof.* By Remark 2.1.3 connected limits are computed in  $\mathcal{A}$  and  $p_!$  doesn't affect objects and morphism in  $\mathcal{A}$ .  $\square$

**Proposition 2.2.2.** *Let  $p : E \rightarrow B$  be a morphism in  $\mathcal{A}$  with pullbacks, then*

$$(\mathcal{A} \downarrow B) \begin{array}{c} \xrightarrow{p^*} \\ \xleftarrow{p_!} \end{array} (\mathcal{A} \downarrow E) \quad \text{with } p_! \dashv p^*$$

*is an adjunction. The unit is given by*

$$\eta : \text{Id}_{(\mathcal{A} \downarrow E)} \rightarrow p^* p_!$$

$$\eta_{(D, \delta)} = \langle \delta, 1_D \rangle : (D, \delta) \rightarrow (E \times_B D, \text{proj}_1)$$

*and the counit is given by*

$$\epsilon : p_! p^* \rightarrow \text{Id}_{(\mathcal{A} \downarrow B)}$$

$$\epsilon_{(A, \alpha)} = \text{proj}_2 : (E \times_B A, p(\text{proj}_1)) \rightarrow (A, \alpha)$$

*Proof.* We want to prove this is adjunction. We start proving that the unit is a natural transformation. Let  $g : (D, \delta) \rightarrow (D', \delta')$  be a morphism in  $(\mathcal{A} \downarrow E)$ , we want to prove that the following diagram commutes:

$$\begin{array}{ccc} D & \xrightarrow{\eta_{(D, \delta)}} & E \times_B D \\ \downarrow g & & \downarrow p^* p_!(g) = \tilde{g} \\ D' & \xrightarrow{\eta_{(D', \delta')}} & E \times_B D' \end{array}$$

In order to do that we need to expand the action of the functor  $p^* p_!$  on the morphism  $g$ . We have:

$$\begin{array}{ccc} D & \xrightarrow{g} & D' \\ \downarrow \delta & \searrow \delta' & \\ E & & \end{array} \quad \xrightarrow{p_!} \quad \begin{array}{ccc} D & \xrightarrow{g} & D' \\ & \searrow p\delta & \downarrow p\delta' \\ & & B \end{array}$$

$$\begin{array}{c}
 \xrightarrow{p^*} \\
 \begin{array}{ccccc}
 & & \tilde{g} & & \\
 & \swarrow & \text{---} & \searrow & \\
 E \times_B D & \xrightarrow{\text{proj}_2} & D & \xrightarrow{g} & D' & \xleftarrow{\text{proj}'_2} & E \times_B D' \\
 & \nwarrow \eta_{(D,\delta)} & \searrow p\delta & \swarrow p\delta' & \nwarrow \eta_{(D',\delta')} & & \\
 & & B & & & & \\
 & \nwarrow \text{proj}_1 & \uparrow p & \swarrow \text{proj}'_1 & & & \\
 & & E & & & & 
 \end{array}
 \end{array}$$

Consider the following commutative diagram.

$$\begin{array}{ccccc}
 D & \xrightarrow{\eta_{(D,\delta)}} & E \times_B D & & \\
 \downarrow g & \searrow \eta_{(D',\delta')} & \downarrow p^*p_!(g)=\tilde{g} & & \\
 D' & \xrightarrow{\eta_{(D',\delta')}} & E \times_B D' & \xrightarrow{\text{proj}'_2} & D' \\
 & & \downarrow \text{proj}'_1 & & \downarrow p\delta' \\
 & & E & \xrightarrow{p} & B
 \end{array}$$

Since  $(\text{proj}'_1, \text{proj}'_2)$  are jointly monic, by the following compositions

- $(\text{proj}'_2)(\tilde{g})(\eta_{(D,\delta)}) = (g)(\text{proj}_2)(\eta_{(D,\delta)}) = (g)(1_D) = g$
- $(\text{proj}'_2)(\eta_{(D',\delta')})(g) = (1_{D'})(g) = g$
- $(\text{proj}'_1)(\tilde{g})(\eta_{(D,\delta)}) = (\text{proj}_1)(\eta_{(D,\delta)}) = \delta$
- $(\text{proj}'_1)(\eta_{(D',\delta')})(g) = (\delta')(g) = \delta$

we conclude that  $(\tilde{g})(\eta_{(D,\delta)}) = (\eta_{(D',\delta')})(g)$ . Thus the naturality condition of  $\eta$  is satisfied.

Now we want to prove that the counit is a natural transformation. Consider  $f : (A, \alpha) \rightarrow (A', \alpha')$  a morphism in  $(\mathcal{A} \downarrow B)$ , we want to prove that the following diagram commutes:

$$\begin{array}{ccc}
 E \times_B A & \xrightarrow{\epsilon_{(A,\alpha)}} & A \\
 p_!p^*(f)=\tilde{f} \downarrow & & \downarrow f \\
 E \times_B A' & \xrightarrow{\epsilon_{(A',\alpha')}} & A'
 \end{array}$$

As similarly done above, we need to expand the action of the functor  $p_!p^*$  on the morphism  $f$ , we have:

$$\begin{array}{ccc}
 \begin{array}{c} A \xrightarrow{f} A' \\ \downarrow \alpha \quad \swarrow \alpha' \\ B \end{array} & \xrightarrow{p^*} & \begin{array}{c} \begin{array}{ccccc} & & \tilde{f} & & \\ & & \curvearrowright & & \\ E \times_B A & \xrightarrow{\text{proj}_2} & A & \xrightarrow{f} & A' & \xleftarrow{\text{proj}_2'} & E \times_B A' \\ & \searrow \alpha & \swarrow \alpha' & & \\ & B & & & \\ \text{proj}_1 \swarrow & & \searrow \text{proj}_1' & & \\ & E & & & \end{array} \end{array} \\
 \\
 \xrightarrow{p!} & & \begin{array}{c} \begin{array}{ccccc} & & \tilde{f} & & \\ & & \curvearrowright & & \\ E \times_B A & \xrightarrow{\text{proj}_2} & A & \xrightarrow{f} & A' & \xleftarrow{\text{proj}_2'} & E \times_B A' \\ & \searrow \alpha & \swarrow \alpha' & & \\ & B & & & \\ p(\text{proj}_1) \swarrow & & \searrow p(\text{proj}_1') & & \\ & E & & & \end{array} \end{array}
 \end{array}$$

Since  $\epsilon_{(A,\alpha)} = \text{proj}_{2,E \times_B A}$ , the composition  $(f)(\epsilon_{(A,\alpha)}) = (\epsilon_{(A',\alpha')})(\tilde{f})$  holds by the definition of  $\tilde{f}$ . Thus also the naturality of  $\epsilon$  is satisfied.

Now we can check that the unit and counit fulfil the first triangle identities. Let  $(A, \alpha)$  be an object in  $(\mathcal{A} \downarrow B)$ .

$$((p^* \epsilon) \circ (\eta p^*))_{(A,\alpha)} = (p^* \epsilon)_{(A,\alpha)} (\eta p^*)_{(A,\alpha)} = p^* (\text{proj}_{2,E \times_B A}) (\eta_{(E \times_B A, \text{proj}_1)}) = \circledast$$

Considering first how  $p^*$  acts on the morphism  $\text{proj}_{1,E \times_B A}$

$$\begin{array}{ccc}
 \begin{array}{ccc} E \times_B A & \xrightarrow{\text{proj}_{2,E \times_B A}} & A \\ & \searrow \alpha(\text{proj}_{2,E \times_B A}) & \swarrow \alpha \\ & B & \end{array} & \xrightarrow{p^*} & \begin{array}{c} \begin{array}{ccccc} & & \text{proj}_{2,\tilde{E} \times_B A} & & \\ & & \curvearrowright & & \\ E \times_B (E \times_B A) & \xrightarrow{\text{proj}_{2,E \times_B (E \times_B A)}} & E \times_B A & \xrightarrow{\text{proj}_{2,E \times_B A}} & A & \xleftarrow{\text{proj}_{2,E \times_B A}} & E \times_B A \\ & \searrow (\alpha)(\text{proj}_{2,E \times_B A}) & \swarrow \alpha & & \\ & B & & & \\ \text{proj}_{1,E \times_B (E \times_B A)} \swarrow & & \searrow \text{proj}_{1,E \times_B A} & & \\ & E & & & \end{array} \end{array}
 \end{array}$$



and the second composition

$$\begin{aligned}
 & (\text{proj}_{1,E \times_B A})(\tilde{\text{proj}}_{2,E \times_B A})\langle \text{proj}_{1,E \times_B A}, 1_{E \times_B A} \rangle = \\
 & = (\text{proj}_{1,E \times_B (E \times_B A)})(\text{proj}_{1,E \times_B A}, 1_{E \times_B A}) \\
 & = \text{proj}_{1,E \times_B A} \\
 & = (\text{proj}_{1,E \times_B A})(1_{(E \times_B A, \text{proj}_1)})
 \end{aligned}$$

Hence

$$\circledast = (\tilde{\text{proj}}_{2,E \times_B A})\langle \text{proj}_{1,E \times_B A}, 1_{E \times_B A} \rangle = 1_{(E \times_B A, \text{proj}_1)}$$

Finally we can check the second triangle identities. Let  $(D, \delta)$  be an object in  $(\mathcal{A} \downarrow E)$ , we have:

$$\begin{aligned}
 ((\epsilon p_!) \circ (p_! \eta))_{(D, \delta)} &= ((\epsilon p_!)_{(D, \delta)}(p_! \eta))_{(D, \delta)} = (\epsilon_{p_!(D, \delta)})p_!\langle \delta, 1_D \rangle = \\
 &= (\epsilon_{(D, p\delta)})p_!(\langle \delta, 1_D \rangle) = \text{proj}_{2,E \times_B D}\langle \delta, 1_D \rangle = 1_{(D, p\delta)}
 \end{aligned}$$

$$\begin{array}{ccc}
 E \times_B D & \xrightarrow{\text{proj}_{2,E \times_B D}} & D \\
 \downarrow \text{proj}_{1,E \times_B D} & & \downarrow p\delta \\
 E & \xrightarrow{p} & B
 \end{array}$$

So we can conclude that the adjunction holds.  $\square$

**Remark 2.2.3.**  $p^*$  is a right adjoint functor, so it preserves limits.  $p_!$  is a left adjoint functor, so it preserves colimits.

**Property 2.2.4.** The diagram constructed by  $p^*$  acting on a morphism  $f : (A, \alpha) \rightarrow (A', \alpha')$  is a pullback in  $(\mathcal{A} \downarrow B)$ .

$$\begin{array}{ccc}
 E \times_B A & \xrightarrow{p^*(f)=\tilde{f}} & E \times_B A' \\
 \downarrow \text{proj}_2 & & \downarrow \text{proj}'_2 \\
 A & \xrightarrow{f} & A'
 \end{array}$$

*Proof.* We prove it in  $\mathcal{A}$ , then by Remark 2.1.3 it holds in  $\mathcal{A} \downarrow B$ . Since the right and the total diagrams below are pullback, then the left diagram is a

pullback.

$$\begin{array}{ccccc}
 & & \text{proj}_1 & & \\
 & \nearrow & & \searrow & \\
 E \times_B A & \xrightarrow{p^*(f)=\tilde{f}} & E \times_B A' & \xrightarrow{\text{proj}'_1} & E \\
 \downarrow \text{proj}_2 & & \downarrow \text{proj}'_2 & & \downarrow p \\
 A & \xrightarrow{f} & A' & \xrightarrow{\alpha'} & B \\
 & \searrow & & \nearrow & \\
 & & \alpha & & 
 \end{array}$$

□

**Property 2.2.5.** *The unit  $\eta$  and the counit  $\epsilon$  are cartesian.*

*Proof.* First we prove that  $\eta$  is cartesian. Let  $g : (D, \delta) \rightarrow (D', \delta')$  be a morphism in  $(\mathcal{A} \downarrow E)$ . Applying *Property 2.2.4* to  $p_!(g)$ , we obtain that the following diagram is a pullback.

$$\begin{array}{ccc}
 E \times_B D & \xrightarrow{\text{proj}_2} & D \\
 \downarrow p^*p_!(g)=\tilde{g} & & \downarrow g \\
 E \times_B D' & \xrightarrow{\text{proj}'_2} & D'
 \end{array}$$

Considering the following commutative diagram, since the right diagram is a pullback, as we just proved, and the total diagram is a pullback, we can conclude that the left diagram is a pullback, as stated.

$$\begin{array}{ccccc}
 & & 1_D & & \\
 & \nearrow & & \searrow & \\
 D & \xrightarrow{\eta_{(D,\delta)}} & E \times_B D & \xrightarrow{\text{proj}_2} & D \\
 \downarrow g & & \downarrow p^*p_!(g)=\tilde{g} & & \downarrow g \\
 D' & \xrightarrow{\eta_{(D',\delta')}} & E \times_B D' & \xrightarrow{\text{proj}'_2} & D' \\
 & \searrow & & \nearrow & \\
 & & 1_{D'} & & 
 \end{array}$$

Now we want to prove that  $\epsilon$  is cartesian. Let  $f : (A, \alpha) \rightarrow (A', \alpha')$  be a morphism in  $(\mathcal{A} \downarrow B)$ . By *Proposition 2.2.1* applied at the pullback in *Property 2.2.4* for  $f$  we obtain that the following diagram is a pullback.

$$\begin{array}{ccc}
 E \times_B A & \xrightarrow{\epsilon_{(A,\alpha)}=\text{proj}_2} & A \\
 \downarrow p_!p^*(f)=\tilde{f} & & \downarrow f \\
 E \times_B A' & \xrightarrow{\epsilon_{(A',\alpha')}=\text{proj}'_2} & A'
 \end{array}$$

□

**Proposition 2.2.6.** *Let  $p : E \rightarrow B$  be a morphism in  $\mathcal{A}$  with pullbacks. Then  $p^*$  preserves coequalizer of kernel pairs whose coequalizers are pullback stable regular epimorphisms.*

*Proof.* Consider the following diagram:

$$\begin{array}{ccccc}
 p^*(\text{Eq}(q)) & \xrightarrow[p_1]{\tilde{p}_2} & p^*(A) = A \times_B E & \xrightarrow{\tilde{q}} & p^*(\text{Coeq}(p_1, p_2)) \\
 \downarrow & & \downarrow & & \downarrow \\
 \text{Eq}(q) & \xrightarrow[p_1]{p_2} & A & \xrightarrow{q} & \text{Coeq}(p_1, p_2)
 \end{array}$$

Let  $q$  be the coequalizer of the kernel pair  $\text{Eq}(q)$ . We can affirm by *Property 2.2.4* applied to  $q$  that the diagram in the bottom-right corner is a pullback. Since, by hypothesis,  $q$  is a pullback stable regular epimorphism, we have that  $\tilde{q}$  is a regular epimorphism.

We now aim to show that  $\tilde{q}$  is also the coequalizer of  $(\tilde{p}_1, \tilde{p}_2)$ . Since  $\tilde{q}$ ,  $\tilde{p}_1$  and  $\tilde{p}_2$  are obtained applying  $p^*$  to the kernel pair  $\text{Eq}(q)$ , and  $p^*$  preserves limits, it follows that  $p^*(\text{Eq}(q))$  is the kernel pair of  $\tilde{q}$ . We already know that  $\tilde{q}$  is a regular epi. Thus, by *Proposition A.0.7*,  $\tilde{q}$  is also the coequalizer of  $(\tilde{p}_1, \tilde{p}_2)$ . □

Later on it will be useful to study monads. Thus, we state the following proposition.

**Property 2.2.7.** *Let  $p : E \rightarrow B$  be a morphism in  $\mathcal{A}$  with pullbacks. The associated monad  $(p^*p_!, \eta, p^*\epsilon p_!)$  is cartesian.*

*Proof.* We need to prove that the category  $(\mathcal{A} \downarrow E)$  has pullback, the monad preserves pullbacks, and both the multiplication  $p^*\epsilon p_!$  and the unit  $\eta$  are cartesian.

The category  $(\mathcal{A} \downarrow E)$  has pullbacks, by *Proposition 2.1.4*, since  $\mathcal{A}$  has. The monad  $p^*p_!$  preserves pullbacks, in fact  $p^*$  is a right adjoint so it preserves limits, therefore preserves pullbacks, and  $p_!$  preserves pullbacks by *Proposition 2.2.1*. We already proved that the unit  $\eta$  is cartesian, so it remains to prove that the multiplication  $p^*\epsilon p_!$  is cartesian. Since  $\epsilon$  is cartesian and  $p^*$  preserves pullbacks as shown above, we can conclude that the multiplication is cartesian. □

## 2.3 Nullary and essentially nullary monads

**Definition 2.3.1.** A monad  $(T, \eta, \mu)$  on a category  $\mathcal{X}$  with finite coproducts is said to be **nullary** if, for each object  $X$  in  $\mathcal{X}$ , the morphism  $[T(!_X), \eta_X]$  is an iso, where  $!_X$  is the unique morphism from the initial object of  $\mathcal{X}$  to  $X$ . A monad is said to be **essentially nullary** if, for each object  $X$  in  $\mathcal{X}$ , the morphism  $[T(!_X), \eta_X]$  is a strong epi.

Now we claim a lemma which will be useful to prove some theorems.

**Lemma 2.3.2.** If a category  $\mathcal{X}$  is protomodular, then for each commutative diagram of the form

$$\begin{array}{ccc} V & \xrightarrow{u} & X \\ q \downarrow \uparrow t & & p \downarrow \uparrow s \\ W & \xrightarrow{v} & Y \end{array}$$

where  $p, q$  are split epimorphisms with respective sections  $s, t$  and the downward directed square is a pullback, it holds that the pair  $(u, s)$  is a strongly epimorphic pair.

*Proof.* Assume  $\mathcal{X}$  is protomodular. We want to prove that if  $u$  and  $s$  factorize through a mono  $m$  as  $u = mf$  and  $s = ms'$ , then  $m$  is an iso. By a standard result, proved in *Proposition A.0.5*, this implies that  $(u, s)$  is a strongly epimorphic pair.

Define  $p' = pm$ , we have that  $p's' = 1_Y$ , in fact  $p's' = pms' = ps = 1_Y$ . Pulling back with the change-of-base functor  $v^*$  the mono  $m$  we obtain the following diagram. In this operation we also generate the morphisms  $q', t'$  and  $w$ . In the same way as in *Property 2.2.4* we have that the diagram between  $m$  and  $v^*(m)$  is a pullback.

Now we want to prove that  $v^*(m)$  is an iso. Consider the following diagram.

We know that the external diagram commutes by the hypothesis factorization.

$$\begin{array}{ccccc}
 V & & & & \\
 & \searrow \langle f, 1_V \rangle & & \nearrow 1_V & \\
 & & W & \xrightarrow{v^*(m)} & V \\
 & \searrow f & \downarrow w & & \downarrow u \\
 & & X' & \xrightarrow{m} & X
 \end{array}$$

Since the square is a pullback, it's obvious that  $v^*(m)\langle f, 1_V \rangle = 1_V$ . It remains to prove that  $\langle f, 1_V \rangle v^*(m) = 1_W$ .

Notice that  $mf v^*(m) = uv^*(m) = mw$ , so  $fv^*(m) = w$  since  $m$  is a mono. Now consider the following compositions, and use what we just noticed.

- $v^*(m)(\langle f, 1_V \rangle v^*(m)) = (v^*(m)\langle f, 1_V \rangle)v^*(m) = 1_V v^*(m) = v^*(m)1_W$
- $w(\langle f, 1_V \rangle v^*(m)) = (w\langle f, 1_V \rangle)v^*(m) = fv^*(m) = w = w1_W$

Since the pair  $(v^*(m), w)$  is jointly monic we have that  $\langle f, 1_V \rangle v^*(m) = 1_W$ . Consdiering this composition and the one above, it follows that  $v^*(m)$  is an iso.

By the propomodularity of  $\mathcal{X}$ , since  $v^*(m)$  is an iso, we have that  $m$  is an iso, as claimed.  $\square$

**Theorem 2.3.3.** *If  $\mathcal{X}$  is a pointed protomodular category with finite coproducts and  $(T, \eta, \mu)$  is a monad on  $\mathcal{X}$  with cartesian  $\eta$ , then the monad is essentially nullary.*

*Proof.* For each object  $X$  in  $\mathcal{X}$  we can consider the following diagram:

$$\begin{array}{ccc}
 X & \xrightarrow{\eta_X} & T(X) \\
 !^X \downarrow \uparrow !_X & & T(!^X) \downarrow \uparrow T(!_X) \\
 0 & \xrightarrow{\eta_0} & T(0)
 \end{array}$$

By hypothesis  $\eta$  is cartesian, so the downward directed square is a pullback. Since  $\mathcal{X}$  is protomodular then by *Lemma 2.3.2*, the pair  $(\eta_X, T(!_X))$  is strongly epimorphic. By a standard result, proved in *Proposition A.0.6*, this yields that  $[\eta_X, T(!_X)]$  is a strong epi. Thus the monad is essentially nullary.  $\square$

## Chapter 3

# Ideally exact categories

### 3.1 Preliminary results on the Eilenberg-Moore categories

We present here some technical results on the Eilenberg-Moore categories. These will play a key role in the next section, where we define and characterize the ideally exact categories through monadic functors.

**Proposition 3.1.1.** *Let  $T : \mathcal{X} \rightarrow \mathcal{X}$  be a monad preserving regular epimorphisms and kernel pairs. If  $\mathcal{X}$  is exact, then the Eilenberg-Moore category  $\mathcal{X}^T$  is exact.*

*Proof.* Assume  $\mathcal{X}$  is exact. With the given hypotheses on the monad  $T$ , we aim to prove that  $\mathcal{X}^T$  is exact. We need to establish that it is finitely complete, that each equivalent relation is the kernel pair of some arrow, that regular epis are pullback stable and that each morphism admits a (regular epi, mono) factorization.

First we show that  $\mathcal{X}^T$  is finitely complete. We recall that the forgetful functor  $U^T : \mathcal{X}^T \rightarrow \mathcal{X}$  creates all the limits that  $\mathcal{X}$  has. Since  $\mathcal{X}$  is exact, it has all finite limits. Therefore,  $\mathcal{X}^T$  has all finite limits.

We proceed to prove that each equivalent relation is the kernel pair of some arrow. Consider an equivalence relation  $R$  in  $\mathcal{X}^T$  with  $\xi_R$  and  $\xi_A$  giving the structure of algebras. Then, under the forgetful functor  $U^T$ ,  $R$  is an equivalence relation in  $\mathcal{X}$ .  $\mathcal{X}$  is exact, so  $R$  is the kernel pair of some arrow. In particular, by *Proposition A.0.9*, we can suppose that  $R$  is the kernel pair of its coequalizer. Let call it  $f : A \rightarrow B$ . Since  $T$  preserves kernel pairs, we have that  $T(R)$  is the kernel pair of  $T(f)$ . Moreover  $T$  preserves regular epis, so  $T(f)$  is a regular epi. By *Proposition A.0.7*, we can assume that  $T(f)$  is the coequalizer of its kernel pair  $(T(R), r_1, r_2)$ . Since  $f\xi_A T(r_1) = fr_1\xi_R = fr_2\xi_R = f\xi_A T(r_2)$ , by the universal property of the coequalizer, there exists a unique morphism  $\xi_B$  such that  $\xi_B T(f) = f\xi_A$ . The morphism  $\xi_B$  is the algebra structure of  $T(B)$ . Since  $U^T$  creates all the limits that  $\mathcal{X}$  has, we can conclude that  $R$  is

the kernel pair in  $\mathcal{X}^T$  of some arrow.

$$\begin{array}{ccccc}
 T(R) & \xrightleftharpoons[T(r_2)]{T(r_1)} & T(A) & \xrightarrow{T(f)} & T(B) \\
 \downarrow \xi_R & & \downarrow \xi_A & & \downarrow \xi_B \\
 R & \xrightleftharpoons[r_2]{r_1} & A & \xrightarrow{f} & B
 \end{array}$$

Consider now a morphism  $f : (A, \xi_A) \rightarrow (B, \xi_B)$  in  $\mathcal{X}^T$ , we aim to prove that it admits a (regular epi, mono) factorization. The morphism  $f$ , under the forgetful functor  $U^T$ , is a morphism in  $\mathcal{X}$ , which is exact and in particular regular. Thus  $f$  admits a (regular epi, mono) factorization in  $\mathcal{X}$ . Let it be  $f = me$  with  $m : Q \rightarrow B$  mono and  $e : A \rightarrow Q$  regular epi. Notice that  $A$  and  $B$  are already algebras, we want to provide  $Q$  and the domain of the parallel arrows with a structure of algebra, so that we can lift the factorization to  $\mathcal{X}^T$ .

$$\begin{array}{ccccc}
 D & \xrightleftharpoons{\quad} & T(A) & \xrightarrow{T(f)} & T(B) \\
 & & \downarrow \xi_A & & \downarrow \xi_B \\
 C & \xrightleftharpoons{\quad} & A & \xrightarrow{f} & B \\
 & & \searrow e & & \nearrow m \\
 & & & Q & 
 \end{array}$$

In order to do so, we first need to show that  $T$  preserves coequalizers in  $\mathcal{X}$  of kernel pairs arising from  $\mathcal{X}^T$ , once the algebra structure is forgotten, and considering only the pair of the whole kernel pair. Let  $(\text{Eq}, r_1, r_2)$  be a kernel pair in  $\mathcal{X}$ , arising from a kernel pair in  $\mathcal{X}^T$  of a morphism, and let  $p$  be the coequalizer in  $\mathcal{X}$  of  $(r_1, r_2)$ . By *Proposition A.0.8*, we can affirm that  $(r_1, r_2)$  is the kernel pair of  $p$ . Given that  $(\text{Eq}, r_1, r_2)$  is a kernel pair arising from  $\mathcal{X}^T$ , it follows that  $(T(\text{Eq}), T(r_1), T(r_2))$  is the kernel pair of some arrow. Since  $T$  preserves regular epi and kernel pair we can assert that  $T(p)$  is a regular epi and, by *Proposition A.0.7*, it is the coequalizer of the kernel pair  $(T(\text{Eq}), T(r_1), T(r_2))$ , as claimed.

$$\begin{array}{ccccc}
 T(\text{Eq}) & \xrightleftharpoons[T(r_2)]{T(r_1)} & T(A) & \xrightarrow{T(p)} & T(B) \\
 \downarrow \xi_{\text{Eq}} & & \downarrow \xi_A & & \downarrow \xi_B \\
 \text{Eq} & \xrightleftharpoons[r_2]{r_1} & A & \xrightarrow{p} & B
 \end{array}$$

Picking up where we left off, consider  $f$  and its kernel pair  $(\text{Eq}(f), r_1, r_2)$ . Since  $U^T$  creates all the limits that  $\mathcal{X}$  has, we can lift the kernel pair

$(\text{Eq}(f), r_1, r_2)$  to a kernel pair in  $\mathcal{X}^T$ . Let  $\xi_{\text{Eq}}$  be the structure of algebra of  $\text{Eq}$ .

We know that  $U^T$  creates all the colimits that  $\mathcal{X}$  has and  $T$  preserves. Hence,  $U^T$  creates all coequalizers that  $\mathcal{X}$  has of kernel pairs arising from  $\mathcal{X}^T$ , once the algebra structure is forgotten. Observe that  $e$  is such a coequalizer, in fact it is a regular epi and by *Proposition A.0.10*  $f$  and  $e$  share the same kernel pair, because of the factorization  $f = me$  with  $m$  mono, thus  $(\text{Eq}(f), r_1, r_2)$  is also the kernel pair of  $e$ . Hence, by *Proposition A.0.7*,  $e$  is the coequalizer of  $(\text{Eq}(f), r_1, r_2)$ . In particular by the creation of such colimit we proved that  $Q$  has the structure of algebra, called  $\xi_Q$ , and that  $e$  can be lifted to a coequalizer in  $\mathcal{X}^T$ .

$$\begin{array}{ccccc}
 T(\text{Eq}(f)) & \xrightleftharpoons[T(r_2)]{T(r_1)} & T(A) & \xrightarrow{T(f)} & T(B) \\
 \downarrow \xi_{\text{Eq}(f)} & & \downarrow \xi_A & \searrow T(e) & \nearrow T(m) \\
 & & & T(Q) & \\
 & & & \downarrow \xi_Q & \\
 \text{Eq}(f) & \xrightleftharpoons[r_2]{r_1} & A & \xrightarrow{f} & B \\
 & & \searrow e & & \nearrow m \\
 & & & Q & 
 \end{array}$$

In order to prove the factorization of  $f$  in  $\mathcal{X}^T$ , it remains to establish that  $m$  is a morphism in  $\mathcal{X}^T$  and that it is a mono. Notice that  $m$  is a mono in  $\mathcal{X}^T$ , in fact it is a mono in  $\mathcal{X}$  and furthermore it is merely a morphism  $\mathcal{X}$  that satisfies the algebra condition. We still need to establish that  $m$  is a morphism in  $\mathcal{X}^T$ .

$$\begin{aligned}
 f = mq &\implies f\xi_A = mq\xi_A \\
 &\implies \xi_B T(f) = m\xi_Q T(q) \\
 &\implies \xi_B T(m)T(q) = m\xi_Q T(q)
 \end{aligned}$$

We know that  $T(q)$  is a regular epi, thus it is a epi. So we can cancel it from both sides and we obtain the condition for  $m$  to be a morphism in  $\mathcal{X}^T$ :

$$\xi_B T(m) = m\xi_Q$$

We proved that  $m$  and  $p$  are respectively a mono and a regular epi in  $\mathcal{X}^T$ . This allows the lifting of the factorization  $f = me$  to  $\mathcal{X}^T$ .

It remains to prove the last condition of a regular category, that is the pullback stability of regular epi. Let  $q : A \rightarrow B$  be a regular epi in  $\mathcal{X}^T$ , ie a coequalizer of some arrows in  $\mathcal{X}^T$ . In particular,  $q$  is a regular epi in  $\mathcal{X}$ . Since  $\mathcal{X}$  is exact, it is pullback stable. Let  $\tilde{q}$  the pullback in  $\mathcal{X}^T$  of  $q$  along some morphism. As just noticed,  $\tilde{q}$  is a regular epi in  $\mathcal{X}$ . Let  $(\text{Eq}(\tilde{q}), r_1, r_2)$

be the kernel pair of  $\tilde{q}$ . By *Proposition A.0.7*,  $\tilde{q}$  is the coequalizer of such kernel pair. We want to lift  $\tilde{Q}$  to a regular epi in  $\mathcal{X}^T$ . By construction, the pullback considered is a pullback in  $\mathcal{X}^T$ , thus it comes with algebra structures of  $A \times_B C$  and  $C$ . Let  $\xi_{A \times_B C}$  and  $\xi_C$  be the algebra structures of  $A \times_B C$  and  $C$  respectively. Moreover,  $\tilde{q}$  is a morphism in  $\mathcal{X}^T$ . We know that the forgetful functor  $U^T$  creates all the limits that  $\mathcal{X}$  has, so we can lift the kernel pair  $(\text{Eq}(\tilde{q}), r_1, r_2)$  to a kernel pair in  $\mathcal{X}^T$ . Let  $\xi_{\text{Eq}(\tilde{q})}$  be the algebra structure of  $\text{Eq}(\tilde{q})$ . We know that  $U^T$  creates all the colimits that  $\mathcal{X}$  has and the monad  $T$  preserves, and we already proved that  $T$  preserves coequalizers of kernel pairs. Thus, since  $\tilde{q}$  is such a coequalizer, we can lift it to a coequalizer in  $\mathcal{X}^T$ .

$$\begin{array}{ccccccc}
 & T(K) & \rightrightarrows & T(A) & \xrightarrow{T(q)} & T(B) & \\
 & \downarrow & & \downarrow & & \downarrow & \\
 T(\text{Eq}(\tilde{q})) & \xrightarrow{T(r_1)} & T(A \times_B C) & \xrightarrow{T(\tilde{q})} & T(C) & & \\
 & \downarrow T(r_2) & \downarrow \xi_{A \times_B C} & \downarrow & \downarrow \xi_C & & \\
 & K & \rightrightarrows & A & \xrightarrow{q} & B & \\
 & \downarrow & & \downarrow & & \downarrow & \\
 \text{Eq}(\tilde{q}) & \xrightarrow{r_1} & A \times_B C & \xrightarrow{\tilde{q}} & C & & \\
 & \downarrow r_2 & & & & & 
 \end{array}$$

□

**Proposition 3.1.2.** *Let  $T : \mathcal{X} \rightarrow \mathcal{X}$  be a monad. If  $\mathcal{X}$  is protomodular, then the Eilenberg-Moore category  $\mathcal{X}^T$  is protomodular.*

*Proof.* Since the forgetful functor  $\mathcal{X}^T \rightarrow \mathcal{X}$  creates all the limits that  $\mathcal{X}$  has, we can conclude that  $\mathcal{X}^T$  is finitely complete.

Let now  $f : (X, \xi) \rightarrow (Y, \xi')$  be a morphism in  $\mathcal{X}^T$ . The following diagram of functors is commutative. This essentially follows from the fact that the forgetful functor  $\mathcal{X}^T \rightarrow \mathcal{X}$  creates all the limits that  $\mathcal{X}$  has, in particular the pullbacks, which are used to define the change-of-base functor  $f^*$ .

$$\begin{array}{ccc}
 \text{Pt}_{\mathcal{X}^T}(Y, \xi') & \xrightarrow{f^*} & \text{Pt}_{\mathcal{X}^T}(X, \xi) \\
 \downarrow U^T & & \downarrow U^T \\
 \text{Pt}_{\mathcal{X}}(Y) & \xrightarrow{f^*} & \text{Pt}_{\mathcal{X}}(X)
 \end{array}$$

Let  $\alpha$  be a morphism in  $\text{Pt}_{\mathcal{X}^T}(Y, \xi')$ . Suppose that  $f^*(\alpha)$  is iso, we want to prove that  $\alpha$  is iso. Since  $f^*(\alpha)$  is iso, we have that  $U^T(f^*(\alpha))$  is iso. By the commutativity of the diagram, we have that  $U^T(f^*(\alpha)) = f^*(U^T(\alpha))$ . Thus  $f^*(U^T(\alpha))$  is an iso. Since  $\mathcal{X}$  is protomodular,  $U^T(\alpha)$  is an iso. Finally,  $U^T$  reflects isomorphisms, so  $\alpha$  is an iso. □

We now proceed with results that will allow us to construct coproducts in the Eilenberg–Moore category, assuming they exist in the base category. The following will be rather intricate and technical; readers may choose to skip directly to the final outcome *Proposition 3.1.6*.

For the sake of clarity, we first present the following technical definition.

**Definition 3.1.3.** Let  $U : \mathcal{X} \rightarrow \mathcal{A}$  be a functor, let  $X$  be an object in  $\mathcal{X}$  and let  $\{(f_i, g_i)\}_{i \in I}$  be a family of pair of parallel arrows in  $\mathcal{A}$ , where  $f_i, g_i : A_i \rightarrow U(X)$ . A morphism  $q : X \rightarrow Q$  in  $\mathcal{X}$  is said to be a  *$U$ -coequalizer* of the pair  $\{(f_i, g_i)\}_{i \in I}$  if

- $U(q)f_i = U(q)g_i \ \forall i \in I$  in  $\mathcal{A}$ ;
- If  $h : X \rightarrow Y$  is a morphism in  $\mathcal{X}$  such that  $U(h)f_i = U(h)g_i \ \forall i \in I$  in  $\mathcal{A}$ , then there exists a unique morphism  $t : Q \rightarrow Y$  in  $\mathcal{X}$  such that  $tq = h$ .

$$\begin{array}{ccc}
 A & \xrightarrow[f_i]{g_i} & U(X) \xrightarrow{U(q)} U(Q) \\
 & & \searrow U(h) \quad \downarrow U(t) \\
 & & U(Y)
 \end{array}
 \qquad
 \begin{array}{ccc}
 X & \xrightarrow{q} & Q \\
 & \searrow h & \downarrow t \\
 & & Y
 \end{array}$$

Before continuing, recall the universal property of the unit in an adjunction. Consider a functor  $U : \mathcal{A} \rightarrow \mathcal{B}$  right adjoint to a functor  $F$ . Then for each object  $A$  in  $\mathcal{A}$ ,  $B$  in  $\mathcal{B}$ , and  $f : B \rightarrow U(A)$  in  $\mathcal{B}$  there exists a unique morphism  $g : F(B) \rightarrow A$  in  $\mathcal{A}$  such that  $f = U(g)\eta_B$ . We denote  $g = \bar{f}$ .

$$\begin{array}{ccc}
 B & \xrightarrow{\eta_B} & UF(B) \\
 & \searrow f & \downarrow U(g) \\
 & & U(A)
 \end{array}
 \qquad
 \begin{array}{ccc}
 & & F(B) \\
 & & \downarrow g \\
 & & A
 \end{array}$$

Also recall that a monad  $(T, \mu, \eta)$  induces a functor  $U^T$  and a functor  $F^T$  defined as follows:

$$\begin{array}{ccc}
 \mathcal{X}^T & & (X, \xi) \xrightarrow{U^T} X \\
 \uparrow F^T \quad \downarrow U^T & & \\
 \mathcal{X} & & X \xrightarrow{F^T} (T(X), \mu_X)
 \end{array}$$

**Lemma 3.1.4.** Let  $U : \mathcal{X} \rightarrow \mathcal{A}$  be a functor right adjoint to a functor  $F : \mathcal{A} \rightarrow \mathcal{X}$ , let  $X$  be an object in  $\mathcal{X}$  and let  $\{(f_i, g_i)\}_{i \in I}$  be a family of parallel arrows in  $\mathcal{X}$ , where  $f_i, g_i : A_i \rightarrow U(X)$ . Consider the family of unique morphisms induced by the universal property of the unit  $\{(\bar{f}_i, \bar{g}_i)\}_{i \in I}$ , such that  $\bar{f}_i = U(f_i)\eta_{A_i}$  and  $\bar{g}_i = U(g_i)\eta_{A_i}$ . If  $q : X \rightarrow Q$  is a coequalizer of the family  $\{(\bar{f}_i, \bar{g}_i)\}_{i \in I}$  in  $\mathcal{X}$ , then  $q$  is a  $U^T$ -coequalizer of the family  $\{(f_i, g_i)\}_{i \in I}$  in  $\mathcal{A}$ .

$$\begin{array}{ccc}
 A_i & \xrightarrow{\eta_{A_i}} & UF(A_i) \\
 & \searrow f_i & \downarrow U(\bar{g}_i) \quad \downarrow U(\bar{f}_i) \\
 & \searrow g_i & U(X)
 \end{array}
 \qquad
 \begin{array}{ccc}
 F(A_i) & \xrightarrow[\bar{g}_i]{\bar{f}_i} & X \xrightarrow{q} Q \\
 A_i \xrightarrow[\bar{g}_i]{\bar{f}_i} & U(X) & \xrightarrow{U(q)} U(Q)
 \end{array}$$

*Proof.* Assume  $q : X \rightarrow Q$  is a coequalizer of the family  $\{(\bar{f}_i, \bar{g}_i)\}_{i \in I}$  in  $\mathcal{X}$ , we want to prove that  $q$  is a  $U^T$ -coequalizer of the family  $\{(f_i, g_i)\}_{i \in I}$  in  $\mathcal{A}$ . First we show, for each  $i$ , that  $U(q)f_i = U(q)g_i$ . Assume that  $q\bar{f}_i = q\bar{g}_i$ .

$$\begin{aligned}
 q\bar{f}_i = q\bar{g}_i &\implies U(q)U(\bar{f}_i) = U(q)U(\bar{g}_i) \\
 &\implies U(q)U(\bar{f}_i)\eta_{A_i} = U(q)U(\bar{g}_i)\eta_{A_i} \\
 &\implies U(q)f_i = U(q)g_i
 \end{aligned}$$

We aim to prove now the universal property of the  $U$ -coequalizer. Let  $h : X \rightarrow Y$  be a morphism such that  $U(h)f_i = U(h)g_i$ . By the universal property of the unit applied to  $U(h)f_i = U(h)g_i : A_i \rightarrow U(Y)$  there exist a unique morphism  $k : F(A_i) \rightarrow Y$  such that  $U(k)\eta_{A_i} = U(h)f_i = U(h)g_i$ . Both  $h\bar{f}_i$  and  $h\bar{g}_i$  satisfies such condition; in fact  $U(h\bar{f}_i)\eta_{A_i} = U(h)U(\bar{f}_i)\eta_{A_i} = U(h)f_i$  and similarly for  $h\bar{g}_i$ . Thus by the uniqueness  $h\bar{f}_i = k = h\bar{g}_i$ .

$$\begin{array}{ccc}
 A_i & \xrightarrow{\eta_{A_i}} & UF(A_i) \\
 & \searrow f_i & \downarrow U(\bar{g}_i) \quad \downarrow U(\bar{f}_i) \\
 & \searrow g_i & U(X) \\
 & & \downarrow U(h) \\
 & & U(Y)
 \end{array}
 \qquad
 \begin{array}{ccc}
 F(A_i) & & \\
 \bar{f}_i \downarrow \quad \bar{g}_i \downarrow & & \\
 X & & \\
 \downarrow h & & \\
 Y & &
 \end{array}$$

$U(h)f_i = U(h)g_i$  (curved arrow from  $A_i$  to  $U(Y)$ )  
 $U(k)$  (curved arrow from  $UF(A_i)$  to  $U(Y)$ )  
 $k$  (curved arrow from  $F(A_i)$  to  $Y$ )

Then, by the universal property of the coequalizer  $q$  in  $\mathcal{X}^T$ , there exists a unique morphism  $t : Q \rightarrow Y$  such that  $tq = h$ .  $\square$

**Proposition 3.1.5.** *Let  $\mathcal{X}$  be a category, let  $(T, \mu, \eta)$  be a monad on  $\mathcal{X}$  and let  $\{(A_i, \xi_i)\}_{i \in I}$  be a family of objects in  $\mathcal{X}^T$ . Assume that the coproducts  $\sum_i A_i$  and  $\sum_i T(A_i)$  exist in  $\mathcal{X}$  with respective canonical injections  $\{\iota_i\}_{i \in I}$  and  $\{\iota'_i\}_{i \in I}$ . If*

the following parallel arrows in  $\mathcal{X}^T$  admits a coequalizer  $q : T(\sum_i) \rightarrow (Q, \xi_Q)$

$$\begin{array}{ccc}
 & (T^2(\sum_i A_i), \mu) & \\
 T[\dots, T(\iota_i), \dots] \nearrow & & \searrow \mu_{\sum_i A_i} \\
 (T(\sum_i T(A_i)), \mu) & \xrightarrow{T(\sum_i \xi_i)} & (T(\sum_i A_i), \mu)
 \end{array} \quad (3.1)$$

then both the following diagrams of parallel arrow in  $\mathcal{X}$  have  $p$  as  $U^T$ -coequalizer.

$$\begin{array}{ccc}
 \sum_i T(A_i) & \xrightarrow{[\dots, T(\iota_i), \dots]} & T(\sum_i A_i) \\
 \searrow \sum_i \xi_i & & \nearrow \eta_{\sum_i A_i} \\
 & \sum_i A_i &
 \end{array} \quad (3.2)$$

$$\begin{array}{ccc}
 T(A_i) & \xrightarrow{T(\iota_i)} & T(\sum_i A_i) \\
 \searrow \xi_i & & \nearrow \eta_{\sum_i A_i} \\
 & A_i \xrightarrow{\iota_i} \sum_i A_i &
 \end{array} \quad (3.3)$$

*Proof.* Consider the diagram (3.1) of parallel arrows in  $\mathcal{X}^T$ , and  $p$  its coequalizer, here below explicitly.

$$\begin{array}{ccccc}
 & & T^3(\sum_i A_i) & & \\
 & T^2[\dots, T(\iota_i), \dots] \nearrow & \downarrow & \searrow T(\mu_{\sum_i A_i}) & \\
 T^2(\sum_i T(A_i)) & \xrightarrow{T(\sum_i \xi_i)} & T^2(\sum_i A_i) & \xrightarrow{\mu_{\sum_i A_i}} & T^2(\sum_i A_i) \\
 \downarrow \mu_{\sum_i T(A_i)} & & \downarrow \mu_{T(\sum_i A_i)} & & \downarrow \mu_{\sum_i A_i} \\
 & T[\dots, T(\iota_i), \dots] \nearrow & T^2(\sum_i A_i) & \searrow \mu_{\sum_i A_i} & \\
 T(\sum_i T(A_i)) & \xrightarrow{T(\sum_i \xi_i)} & T(\sum_i A_i) & & 
 \end{array}$$

Note that the top and bottom arrows of the diagram (3.1) correspond, by the universal property of the unit of the adjunction, respectively to the top and bottom arrows of the diagram (3.2). In order to prove so, by the uniqueness of the morphism induced by the universal property of the unit, it suffices to show that the image of the diagram (3.1) through  $U^T$ , composed with the

unit, is the diagram (3.2). Such composition results to the following diagram.

$$\begin{array}{ccccc}
 & T(\sum_i T(A_i)) & \xrightarrow{T[\dots, T(\iota_i), \dots]} & T^2(\sum_i A_i) & \\
 \eta_{\sum_i T(A_i)} \nearrow & & & & \searrow \mu_{\sum_i A_i} \\
 \sum_i T(A_i) & & (\spadesuit) & & T(\sum_i A_i) \\
 \eta_{\sum_i T(A_i)} \searrow & & & & \nearrow T(\sum_i \xi_i) \\
 & T(\sum_i T(A_i)) & & & 
 \end{array}$$

Consider now the top arrow in the diagram  $(\spadesuit)$ . Using the naturality of  $\eta$ , we have that

$$\begin{array}{ccc}
 \sum_i T(A_i) & \xrightarrow{\eta_{\sum_i T(A_i)}} & T(\sum_i T(A_i)) \\
 \downarrow [\dots, T(\iota_i), \dots] & & \downarrow T[\dots, T(\iota_i), \dots] \\
 T(\sum_i A_i) & \xrightarrow{\eta_{T(\sum_i A_i)}} & T^2(\sum_i A_i)
 \end{array}$$

$$\begin{aligned}
 \mu_{\sum_i A_i} T[\dots, T(\iota_i), \dots] \eta_{\sum_i T(A_i)} &= \mu_{\sum_i A_i} \eta_{T(\sum_i A_i)} [\dots, T(\iota_i), \dots] \\
 &= 1_{T(\sum_i A_i)} [\dots, T(\iota_i), \dots] \\
 &= [\dots, T(\iota_i), \dots]
 \end{aligned}$$

which is the top arrow in (3.2). Then, for the bottom arrow in  $(\spadesuit)$ , by the naturality of  $\eta$ , we have that

$$\begin{array}{ccc}
 \sum_i T(A_i) & \xrightarrow{\eta_{\sum_i T(A_i)}} & T(\sum_i T(A_i)) \\
 \downarrow \xi_A + \xi_B & & \downarrow T(\xi_A + \xi_B) \\
 \sum_i A_i & \xrightarrow{\eta_{\sum_i A_i}} & T(\sum_i A_i)
 \end{array}$$

□

We are now ready to prove the result that will allow us to construct coproducts in the Eilenberg–Moore category, assuming they exist in the base category. This is possible if we assume the existence of reflexive coequalizers in the Eilenberg–Moore category.

**Proposition 3.1.6.** *Let  $T : \mathcal{X} \rightarrow \mathcal{X}$  be a monad. If  $\mathcal{X}$  has finite coproducts and the Eilenberg–Moore category  $\mathcal{X}^T$  has reflexive coequalizers, then  $\mathcal{X}^T$  has*

*finite coproducts. Moreover  $\mathcal{X}^T$  is finitely cocomplete.*

*Proof.* Let  $(T, \mu, \eta)$  be the given monad, and suppose that  $\mathcal{X}$  has finite coproducts. Let  $(A, \xi_A)$  and  $(B, \xi_B)$  be two objects in  $\mathcal{X}^T$ . We aim to prove that  $(A, \xi_A)$  and  $(B, \xi_B)$  have a coproduct in  $\mathcal{X}^T$ . All the coproducts in the diagram below are taken in  $\mathcal{X}$ , and therefore they exist. Let  $\iota_1$  and  $\iota_2$  be the canonical injections in  $\mathcal{X}$  of  $A$  and  $B$  into  $A + B$ , respectively. Our goal is to prove that the diagram below is composed by a reflexive pair in  $\mathcal{X}^T$ . Then, by assumption, it admits a coequalizer in  $\mathcal{X}^T$ , denoted by  $q$ .

$$\begin{array}{ccccc}
 & & T^3(A+B) & & \\
 & \nearrow T^2[T(\iota_1), T(\iota_2)] & \downarrow & \searrow T(\mu_{A+B}) & \\
 T^2(T(A) + T(B)) & \xleftarrow{T^2(\eta_A + \eta_B)} & T^2(A+B) & \xrightarrow{T(q)} & T(Q) \\
 & \xrightarrow{T^2[\xi_A, \xi_B]} & & & \\
 \downarrow \mu_{T(A)+T(B)} & & \downarrow \mu_{T(A+B)} & & \downarrow \mu_{A+B} \\
 & \nearrow T[T(\iota_1), T(\iota_2)] & T^2(A+B) & \searrow \mu_{A+B} & \\
 T(T(A) + T(B)) & \xleftarrow{T(\eta_A + \eta_B)} & T(A+B) & \xrightarrow{q} & Q \\
 & \xrightarrow{T[\xi_A, \xi_B]} & & & \\
 & & \downarrow \xi_Q & & 
 \end{array}$$

The parallel arrows in  $\mathcal{X}^T$  to be coequalized are

- $T[\xi_A, \xi_B]$
- $\mu_{A+B}T[T(\iota_1), T(\iota_2)]$

We proceed to show that they form a reflexive pair by exhibiting a common section. Consider  $T(\eta_A + \eta_B)$ ; it is a common section, as shown below. It follows the first composition

$$T[\xi_A, \xi_B]T(\eta_A + \eta_B) = T([\xi_A, \xi_B](\eta_A + \eta_B)) = T(1_{A+B}) = 1_{T(A+B)}$$

and the second composition, using the naturality of  $\eta$ .

$$\begin{aligned}
 \mu_{A+B}T[T(\iota_1), T(\iota_2)]T(\eta_A + \eta_B) &= \mu_{A+B}T([T(\iota_1), T(\iota_2)](\eta_A + \eta_B)) \\
 &= \mu_{A+B}T[T(\iota_1)\eta_A, T(\iota_2)\eta_B] \\
 &= \mu_{A+B}T[\eta_{A+B}\iota_1, \eta_{A+B}\iota_2] \\
 &= \mu_{A+B}T(\eta_{A+B}[\iota_1, \iota_2]) \\
 &= \mu_{A+B}T(\eta_{A+B}1_{A+B}) \\
 &= \mu_{A+B}T(\eta_{A+B}) = 1_{T(A+B)}
 \end{aligned}$$

Thus it is a reflexive pair.

Let  $q : T(A + B) \rightarrow Q$  be the coequalizer in  $\mathcal{X}$ , and  $\xi_Q$  be the algebra structure of  $Q$ . We aim to prove that  $(Q, \xi_Q)$  is the coproduct of  $(A, \xi_A)$  and  $(B, \xi_B)$  in  $\mathcal{X}^T$ . First we need to exhibit the canonical injections  $\iota_{Q,1}$  and  $\iota_{Q,2}$ . They are respectively

$$\begin{aligned} A &\xrightarrow{\iota_1} A + B \xrightarrow{\eta_{A+B}} T(A + B) \xrightarrow{q} Q \\ A &\xrightarrow{\iota_2} A + B \xrightarrow{\eta_{A+B}} T(A + B) \xrightarrow{q} Q \end{aligned}$$

We still need to prove that they are morphisms in  $\mathcal{X}^T$ . We prove the first one, the second one is similar. Note that  $\iota_1$  and  $\eta_{A+B}$  are not a morphism in  $\mathcal{X}^T$ , this is not a problem, since we are going to prove that the whole composition  $q\eta_{A+B}\iota_1$  is a morphism in  $\mathcal{X}^T$ . Consider the following diagram. All diagrams commute except the left and the external ones. We aim to prove that the left diagram commutes, so that the external one commutes as well.

$$\begin{array}{ccccccc} T(A) & \xrightarrow{T(\iota_1)} & T(A + B) & \xrightarrow{T(\eta_{A+B})} & T^2(A + B) & \xrightarrow{T(q)} & T(Q) \\ \downarrow \xi_A & & & \searrow & \downarrow \mu_{A+B} & & \downarrow \xi_Q \\ & & & & T(A + B) & \xrightarrow{q} & Q \\ A & \xrightarrow{\iota_1} & A + B & \xrightarrow{\eta_{A+B}} & T(A + B) & \xrightarrow{q} & Q \end{array}$$

□

We now want to study a sufficient condition for a category to admit reflexive coequalizers, so that we can apply the proposition above.

Recall that a pair  $(f, g)$  from  $A$  to  $B$  in a regular category induces a relation with the following construction. Consider the morphism induced by the universal property  $\langle f, g \rangle$ . Since the category is regular, we can take the factorization  $\langle f, g \rangle = \langle r_1, r_2 \rangle e$ , with  $\langle r_1, r_2 \rangle$  mono and  $e$  regular epi. Then  $\langle r_1, r_2 \rangle$  is the relation induced.

$$\begin{array}{ccc} A & \xrightarrow{\langle f, g \rangle} & B \times B \\ & \searrow e & \nearrow \langle r_1, r_2 \rangle \\ & R & \end{array}$$


**Proposition 3.1.7.** *The internal relation induced by a reflexive pair  $(f, g)$  in a regular category is reflexive.*

*Proof.* Consider the pair  $(f, g)$  from  $A$  to  $B$  and the section  $s$  with  $fs = 1_B$


and  $gs = 1_B$ .

$$A \begin{array}{c} \xrightarrow{f} \\ \xleftarrow{s} \\ \xrightarrow{g} \end{array} B$$

Let  $\langle f, g \rangle = \langle r_1, r_2 \rangle e$  be the (regular epi, mono) factorization. Moreover, by the universal property we have that  $\langle f, g \rangle s = \langle 1_B, 1_B \rangle$ .



Thus  $\langle r_1, r_2 \rangle es = \langle 1_B, 1_B \rangle$ . The morphism  $es$  makes the relation reflexive.



□

**Proposition 3.1.8.** *Let  $\mathcal{A}$  be an exact Mal'tsev category. Then  $\mathcal{A}$  has reflexive coequalizers.*

*Proof.* Assume  $\mathcal{A}$  is exact and Mal'tsev. We want to prove that  $\mathcal{A}$  has reflexive coequalizer. Consider a reflexive pair  $(f, g)$  from  $A$  to  $B$ . By *Proposition 3.1.7* the internal relation induced  $\langle r_1, r_2 \rangle$  is reflexive. Since  $\mathcal{A}$  is Malt'sev, then each internal reflexive relation is an equivalence relation. Hence  $\langle r_1, r_2 \rangle$  is an equivalence relation.  $\mathcal{A}$  is also exact, then, by *Proposition A.0.9* the equivalence relation  $(r_1, r_2)$  has a coequalizer  $k$ .

We want to prove that  $k$  is also the coequalizer of  $(f, g)$ . First we note that  $kf = kr_1e = kr_2e = kg$ . It remains to prove the universal property. Let  $h : B \rightarrow C$  be a morphism such that  $hf = hg$ . Then  $hr_1e = hr_2e$ . Since  $e$  is a regular epi, hence an epi, we have that  $hr_1 = hr_2$ . Thus there exists a unique

morphism  $t$  such that  $th = k$ .

$$\begin{array}{ccccc}
 A & & & & C \\
 \downarrow e & \searrow f & & \nearrow h & \downarrow \exists! t \\
 R & \xrightarrow[r_2]{r_1} & B & \xrightarrow[k]{} & \text{Coeq}(r_1, r_2)
 \end{array}$$

□

## 3.2 The notion of ideally exact category

We are now ready to prove the theorem, which will lead to the definition of the ideally exact categories.

**Theorem 3.2.1.** . *Let  $\mathcal{A}$  be a category. The following are equivalent:*

1.  $\mathcal{A}$  is exact and protomodular, has finite coproducts and the morphism  $! : 0 \rightarrow 1$  is a regular epimorphism.
2.  $\mathcal{A}$  is exact, has finite coproducts and there exists  $\mathcal{X}$  semiabelian category with a monadic functor  $U : \mathcal{A} \rightarrow \mathcal{X}$ .

$$\begin{array}{ccc}
 \mathcal{A} & \xrightarrow{\cong} & \mathcal{X}^{UF} \\
 \uparrow F & & \nearrow U \\
 \mathcal{X} & & 
 \end{array}$$

3. There exist  $\mathcal{X}$  semiabelian category with a monadic functor  $U : \mathcal{A} \rightarrow \mathcal{X}$  and  $F$  left adjoint to  $U$  such that the associated monad  $T = UF$  preserves regular epimorphisms and kernel pairs.

**Definition 3.2.2.** A category  $\mathcal{A}$  is said to be **ideally exact** if it satisfies one of the equivalent conditions of Theorem 3.2.1.

*Proof of Theorem 3.2.1.* (1  $\implies$  3) Suppose  $! : 0 \rightarrow 1$  is a regular epi. Since  $\mathcal{A}$  is exact, the pullback functor  $(!)^* : (\mathcal{A} \downarrow 1) \rightarrow (\mathcal{A} \downarrow 0)$  is monadic by (citazione). Note that  $(\mathcal{A} \downarrow 1) \cong \mathcal{A}$ . Now consider  $\mathcal{X} = (\mathcal{A} \downarrow 0)$ , we want to prove that  $\mathcal{X}$  is semiabelian.

$(\mathcal{A} \downarrow 0)$  is pointed, in fact if  $(A, \alpha)$  is an object in  $(\mathcal{A} \downarrow 0)$ , we have the

following initial and terminal objects, which are the same:

$$\begin{array}{ccc} 0 & \xrightarrow{!_A} & A \\ & \searrow & \downarrow \alpha \\ & & 0 \end{array} \qquad \begin{array}{ccc} A & \xrightarrow{\alpha} & 0 \\ \downarrow \alpha & \swarrow 1_0 & \\ 0 & & \end{array}$$

$\mathcal{A}$  has finite coproducts, so  $(\mathcal{A} \downarrow 0)$  has finite coproducts. In fact consider  $(A, \alpha)$ ,  $(B, \beta)$  in  $(\mathcal{A} \downarrow 0)$  and  $A + B$  in  $\mathcal{A}$ , with the canonical injections  $\iota_1 : A \rightarrow A + B$  and  $\iota_2 : B \rightarrow A + B$ . Let  $(C, \gamma)$  be an object in  $(\mathcal{A} \downarrow 0)$  with  $f : A \rightarrow C$  and  $g : B \rightarrow C$  morphisms in  $(\mathcal{A} \downarrow 0)$ . Providing to  $A + B$  the morphism  $[\alpha, \beta]$  we have that  $(A + B, [\alpha, \beta])$  is the coproduct of  $(A, \alpha)$  and  $(B, \beta)$  in  $(\mathcal{A} \downarrow 0)$ , in fact  $\gamma[f, g] = [\alpha, \beta]$ , so  $[f, g]$  is also a morphism in  $(\mathcal{A} \downarrow 0)$ .

$$\begin{array}{ccccc} A & \xrightarrow{\iota_1} & A + B & \xleftarrow{\iota_2} & B \\ & \searrow \alpha & \downarrow [\alpha, \beta] & \swarrow \beta & \\ & & 0 & & \\ & \searrow f & \downarrow \gamma & \swarrow g & \\ & & C & & \end{array}$$

Now we want to prove that  $(\mathcal{A} \downarrow 0)$  is exact, from the exactness of  $\mathcal{A}$ . First we prove it is finitely complete. It suffices to show that it has terminal a object and pullbacks. The terminal object is the one just shown above. It also has pullbacks, by *Proposition 2.1.4*, since pullbacks are connected limits and  $\mathcal{A}$  has them.

Consider now an equivalence relation  $(R, r_1, r_2)$  in  $(\mathcal{A} \downarrow 0)$ , endowed with the morphism  $\rho$  and  $\alpha$  as shown in the diagram below. We want to prove that such equivalence relation is the kernel pair of some arrow. Since  $\mathcal{A}$  is exact, it is a kernel pair in  $\mathcal{A}$ . By *Proposition A.0.9* we can suppose that it is the kernel pair of its coequalizer.

$$\begin{array}{ccccc} R & \xrightarrow{r_1} & A & \xrightarrow{q} & B \\ & \searrow r_2 & \downarrow \alpha & \swarrow \exists! \beta & \\ & \searrow \rho & 0 & & \end{array}$$

Since  $r_1$  and  $r_2$  are morphism in  $(\mathcal{A} \downarrow 0)$ , we have that  $\alpha r_1 = \rho = \alpha r_2$ . Thus, by the universal property of the coequalizer, there exists a unique morphism  $\beta : B \rightarrow 0$  such that  $\beta q = \alpha$ . Providing to  $B$  the morphism  $\beta$ , we have that  $(B, \beta)$  is the coequalizer of  $(R, r_1, r_2)$  in  $(\mathcal{A} \downarrow 0)$ .

We proceed by proving that  $(\mathcal{A} \downarrow 0)$  is regular. Both the conditions of the definition of regular category are easily extendable from  $\mathcal{A}$  to  $(\mathcal{A} \downarrow 0)$ . In fact since in  $\mathcal{A}$  each morphism admits a (regular epi, mono) factorization, then the same holds in  $(\mathcal{A} \downarrow 0)$ . Let  $f = me$  be such a factorization through  $Q$ , with  $f$  morphism in  $(\mathcal{A} \downarrow 0)$ . Endow  $Q$  with the morphism  $t$  induced by the

universal property of the coequalizer. Now we want to prove that  $e$  and  $m$  are morphism in  $(\mathcal{A} \downarrow 0)$ . Then they will be respectively regular epi and mono in  $(\mathcal{A} \downarrow 0)$ . Notice first that  $e$  is obviously a morphism in  $(\mathcal{A} \downarrow 0)$ , by the condition of  $t$ . Moreover, we have that  $te = \alpha = \beta f = \beta me$  and since  $e$  is a regular epi, hence an epi, it follows that  $m$  is a morphism in  $(\mathcal{A} \downarrow 0)$ .

$$\begin{array}{ccc}
 & Q & \\
 e \nearrow & \text{---} & \nwarrow m \\
 A & \xrightarrow{f} & B \\
 \searrow \alpha & \text{---} \exists! t & \swarrow \beta \\
 & 0 & 
 \end{array}$$

In order to prove that regular epis are pullback stable in  $(\mathcal{A} \downarrow 0)$  we can use the fact that they are in  $\mathcal{A}$ . In fact, we just endow the regular epi with the morphism induced by universal property of the coequalizer.

We must now establish that the  $(\mathcal{A} \downarrow 0)$  is protomodular. Since the construction of the change-of-base functor involves pullbacks, which are a connected limits, then by *Remark 2.1.3* the construction is the same in  $(\mathcal{A} \downarrow 0)$  and in  $\mathcal{A}$ . Thus  $(\mathcal{A} \downarrow 0)$  inherits protomodularity from  $\mathcal{A}$ , noting that the definition of iso in  $\mathcal{A}$  and  $(\mathcal{A} \downarrow 0)$  is the same. Hence, since  $\mathcal{X} = (\mathcal{A} \downarrow 0)$  is exact, protomodular, pointed and has binary coproducts, we can conclude that it is semiabelian.

It remains to prove that the monad  $(!)^*(!)$  preserves regular epimorphisms and kernel pairs. Consider a regular epi  $q : A \rightarrow B$  in  $(\mathcal{A} \downarrow 0)$ , it is the coequalizer of some parallel arrows. Since the coequalizer is a colimit, it is preserved by the left adjoint functor  $(!)$ . So it suffices to prove that, with  $\mathcal{A}$  regular,  $(!)^*$  preserves regular epis. Consider the following diagram:

$$\begin{array}{ccc}
 0 \times_1 A & \xrightarrow{(!)^*(q)} & 0 \times_1 B \\
 \downarrow & & \downarrow \\
 A & \xrightarrow{q} & B \\
 & \searrow & \swarrow \\
 & 1 & 
 \end{array}$$

By *Property 2.2.4* the square is a pullback. Since  $\mathcal{A}$  is regular, and hence so is  $(\mathcal{A} \downarrow 0)$ , it follows that  $(!)^*$  is a regular epi due to the stability of regular epis under pullbacks.

Finally, we want to establish that the monad  $(!)^*(!)$  preserves kernel pairs. This is obvious:  $(!)$  preserves connected limits, such as kernel pairs, by *Proposition 2.2.1*, while  $(!)^*$  preserves kernel pairs since it is a right adjoint. Thus  $(!)^*(!)$  preserves kernel pairs.

(3  $\implies$  2) Suppose there exists  $\mathcal{X}$  semiabelian category and a monadic functor  $U : \mathcal{A} \rightarrow \mathcal{X}$  satisfying the stated conditions. Assume that the associated monad  $T$  preserves regular epimorphisms and kernel pairs.

First we need to verify that  $\mathcal{A}$  is exact. Since  $U$  is monadic, we have that  $\mathcal{A} \cong \mathcal{X}^T$ . Then it suffices to show that  $\mathcal{X}^T$  is exact. We know that  $\mathcal{X}$  is semiabelian, so it is exact. Then, by *Proposition 3.1.1*,  $\mathcal{X}^T$  is exact.

In order to prove that  $\mathcal{A}$  has finite coproducts, it is first necessary to establish that  $\mathcal{X}^T$  is protomodular. This follows directly from *Proposition 3.1.2* as  $\mathcal{X}$  is protomodular. Since any protomodular category is Malt'sev, then  $\mathcal{X}^T$  is Malt'sev. We already prove that  $\mathcal{X}^T$  is also exact, then by *Proposition 3.1.8*,  $\mathcal{X}^T$  has reflexive coequalizers. Since  $\mathcal{X}$  has finite coproducts and  $\mathcal{X}^T$  has reflexive coequalizers, then by *Proposition 3.1.6*,  $\mathcal{X}^T$  has finite coproducts. Finally, since  $\mathcal{A} \cong \mathcal{X}^T$ , it follows that  $\mathcal{A}$  has finite coproducts. □

# Appendix

**Definition A.0.1.** Let  $\gamma$  be a natural transformation between two functors  $F, G : \mathcal{C} \rightarrow \mathcal{D}$ , we say that  $\gamma$  is **cartesian** if for every morphism  $f : A \rightarrow B$  in  $\mathcal{C}$ , the following diagram is a pullback:

$$\begin{array}{ccc} F(A) & \xrightarrow{\gamma_A} & G(A) \\ \downarrow F(f) & & \downarrow G(f) \\ F(B) & \xrightarrow{\gamma_B} & G(B) \end{array}$$

ie the naturality diagrams are pullbacks.

**Definition A.0.2.** Let  $\mathcal{C}$  be a category and  $f : A \rightarrow B$ . We say that  $f$  is a **strong epimorphism** if it for each commutative diagram of solid arrows

$$\begin{array}{ccc} A & \xrightarrow{f} & B \\ u \downarrow & \exists! d \nearrow & \downarrow v \\ C & \xrightarrow{m} & D \end{array}$$

where  $m$  is a mono, there exists a unique morphism  $d$  such that  $df = u$  and  $md = v$ .

**Proposition A.0.3.** Let  $\mathcal{C}$  be a category with pullbacks and  $f : A \rightarrow B$ . The following are equivalent:

1.  $f$  is a strong epimorphism;
2. Given a factorization  $f = mg$ , if  $m$  is a mono, then it is an iso.

*Proof.* (1  $\implies$  2) Let  $f = mg$  be a factorization with  $m$  mono and consider

the square:

$$\begin{array}{ccc} A & \xrightarrow{f} & B \\ g \downarrow & \searrow d & \downarrow 1_B \\ C & \xrightarrow{m} & B \end{array}$$

By (1) we have  $md = 1_B$ . Furthermore  $mdm = m$  and, since  $m$  is mono, we have that  $dm = 1_C$ . Thus  $m$  is an iso.

(2  $\implies$  1) We want to prove that  $f$  is a strong epi. Consider the pullback

$$\begin{array}{ccccc} A & & & & \\ & \searrow & & \searrow f & \\ & & C \times_D B & \xrightarrow{\text{proj}_2} & B \\ & \swarrow u & \downarrow \text{proj}_1 & & \downarrow v \\ & & C & \xrightarrow{m} & D \end{array}$$

Since  $m$  is a mono, the projection  $\text{proj}_2$  is a mono. Now  $f$  factors through  $\text{proj}_2$ , so by (2)  $\text{proj}_2$  is an iso. We define  $d = \text{proj}_1(\text{proj}_2)^{-1}$ , which is the morphism we are looking for.  $\square$

**Definition A.0.4.** Let  $\mathcal{C}$  be a category and  $f : A \rightarrow B$ ,  $g : A' \rightarrow B$  be morphisms. We say that the pair  $(f, g)$  is a **strongly epimorphic pair** if for each commutative diagram of solid arrows

$$\begin{array}{ccccc} & & B & & \\ & g \nearrow & \downarrow f & \searrow h & \\ A' & & A & & D \\ & g' \searrow & \downarrow f' & \nearrow m & \\ & & C & & \end{array}$$

$\exists! d$

where  $m$  is a mono, there exists a unique morphism  $d$  such that  $f' = df$ ,  $g' = dg$  and  $md = h$ .

**Proposition A.0.5.** Let  $\mathcal{C}$  be a category with pullbacks and  $f : A \rightarrow B$ ,  $g : A' \rightarrow B$  be morphisms. The following are equivalent:

1.  $(f, g)$  is a strong epimorphic pair;
2. Given factorizations  $f = mf'$  and  $g = mg'$ , if  $m$  is a mono, then it is an iso.

*Proof.* By adding the morphism  $g'$  and its conditions in the proof of *Proposition A.0.3*, we obtain the proof of the current proposition.  $\square$

**Proposition A.0.6.** *Let  $\mathcal{C}$  be a category with binary coproducts and  $f : A \rightarrow B$ ,  $g : A' \rightarrow B$  be morphisms. The following are equivalent:*

1.  $(f, g)$  is a strongly epimorphic pair;
2.  $[f, g]$  is a strong epimorphism.

*Proof.* (1  $\implies$  2)  $(f, g)$  is a strongly epimorphic pair. Consider the commutative diagram where  $m$  is a mono

$$\begin{array}{ccc} A + A' & \xrightarrow{[f, g]} & B \\ \downarrow l & & \downarrow h \\ C & \xrightarrow{m} & D \end{array}$$

Composing with the injection  $\iota_1 : A \rightarrow A + A'$  and  $\iota_2 : A' \rightarrow A + A'$ , we have the following commutative diagram


$$\begin{array}{ccccc} & & B & & \\ & g \nearrow & & h \searrow & \\ A' & & A & & D \\ & \nwarrow l_2 & \nwarrow l_1 & & \\ & & C & & \end{array}$$

Since  $(f, g)$  is strongly epimorphic, there exists a unique morphism  $d : B \rightarrow C$  such that  $l_1 = df$ ,  $l_2 = dg$  and  $md = h$ . The first two yield the equalities  $d[f, g]\iota_1 = df = l_1$  and  $[f, g]\iota_2 = dg = l_2$ . So, by the universal property of the coproduct, we have that  $d[f, g] = l$ .

$$\begin{array}{ccc} A & & A \\ \downarrow \iota_1 & & \downarrow \iota_1 \\ A + A' & \xrightarrow{[f, g]} B \xrightarrow{d} C & A + A' \xrightarrow{l} C \\ \uparrow \iota_2 & & \uparrow \iota_2 \\ A' & & A' \end{array}$$

We already know that  $md = h$ , hence  $[f, g]$  is a strong epimorphism.

(2  $\implies$  1)  $[f, g]$  is a strong epimorphism. Consider the following commutative diagram where  $m$  is a mono



Thus we have the commutative diagram


$$\begin{array}{ccc} A + A' & \xrightarrow{[f,g]} & B \\ [f',g'] \downarrow & & \downarrow h \\ C & \xrightarrow{m} & D \end{array}$$

Since  $[f, g]$  is a strong epimorphism, there exists a unique morphism  $d : B \rightarrow C$  such that  $d[f, g] = [f', g']$  and  $md = h$ . Hence  $f' = [f', g']_{\iota_1} = d[f, g]_{\iota_1} = df$  and  $g' = [f', g']_{\iota_2} = d[f, g]_{\iota_2} = dg$ . Thus  $(f, g)$  is a strongly epimorphic pair.  $\square$

**Proposition A.0.7.** *If a regular epi has a kernel pair, then it is the coequalizer of its kernel pair.*

*Proof.* Let  $p$  be a regular epi, and let  $(r_1, r_2)$  be the kernel pair of  $p$ . Since  $p$  is a regular epi, it is the coequalizer of some parallel arrows. Let  $(s_1, s_2)$  be such pair as in the diagram below. We want to prove that  $p$  is the coequalizer of  $(r_1, r_2)$ .


We already know that  $pr_1 = pr_2$ , as  $(r_1, r_2)$  is the kernel pair of  $p$ . It remains to establish the universal property of the coequalizer. Let  $q : B \rightarrow D$  be a morphism such that  $qr_1 = qr_2$ . By the universal property of the kernel pair, there exists a unique morphism  $t : E \rightarrow A$  such that  $r_1t = s_1$  and  $r_2t = s_2$ . We have that  $qs_1 = qr_1t = qr_2t = qs_2$ , so, by the universal property of the coequalizer for the pair  $(s_1, s_2)$ , there exists a unique morphism  $u : C \rightarrow D$  such that  $q = up$ . Thus the universal property of the coequalizer also holds for the pair  $(r_1, r_2)$ .




□

**Proposition A.0.8.** *If a kernel pair has a coequalizer, then it is the kernel pair of its coequalizer.*

*Proof.* Let  $(K, r_1, r_2)$  be a kernel pair of some arrow  $q : A \rightarrow B$ , and assume that such kernel pair has a coequalizer  $e : B \rightarrow Q$ . We want to prove that  $(r_1, r_2)$  is the kernel pair of its coequalizer  $e$ .




We already know that  $er_1 = er_2$ , since  $e$  is the coequalizer of  $(r_1, r_2)$ . It remains to show that the universal property of the kernel pair holds. Consider  $h : X \rightarrow A$  and  $h' : X \rightarrow A'$  morphisms such that  $eh = eh'$ . Since  $(r_1, r_2)$  is the kernel pair of  $q$ , it holds that  $qr_1 = qr_2$ . Thus by the universal property of the coequalizer  $e$ , there exists a unique morphism  $l : Q \rightarrow B$  such that  $q = le$ . Composing with such morphism  $l$  we have that  $leh = leh'$ , hence  $qh = qh'$ . Since  $(r_1, r_2)$  is the kernel pair of  $q$ , by the universal property there exists a unique morphism  $g : X \rightarrow K$  such that  $r_1g = h$  and  $r_2g = h'$ . Hence  $(r_1, r_2)$  is the kernel pair of  $e$ , its coequalizer.




□

**Proposition A.0.9.** *In an exact category  $\mathcal{A}$ , each equivalence relation has a coequalizer and it is the kernel pair of its coequalizer.*

*Proof.* Consider an equivalence relation  $(R, r_1, r_2)$  in  $\mathcal{A}$ . Since  $\mathcal{A}$  is exact,  $(R, r_1, r_2)$  is the kernel pair of some arrow  $q : A \rightarrow B$ .  $\mathcal{A}$  is also regular, so  $q$  admits a factorization  $q = me$ , where  $m$  is a mono and  $e$  is a regular epi, ie the coequalizer of some arrows. Let  $s_1$  and  $s_2$  be such arrows, we have that  $es_1 = es_2$ . Composing with  $m$ , we have that  $qs_1 = mes_1 = mes_2 = qs_2$ . By the universal property of the kernel pair, there exists a unique morphism  $t : C \rightarrow R$  such that  $r_1t = s_1$  and  $r_2t = s_2$ .



Now we want to prove that  $e$  is the coequalizer of  $(r_1, r_2)$ . We have that  $mer_1 = fr_1 = fr_2 = mer_2$ . Thus, since  $m$  is a mono,  $er_1 = er_2$ . It remains to prove the universal property of the coequalizer. Let  $f : A \rightarrow D$  be a morphism such that  $fr_1 = fr_2$ . Composing with  $t$  we have  $fr_1t = fr_2t$ , hence  $fs_1 = fs_2$ . By the universal property of the coequalizer  $e$ , as coequalizer of  $s_1$  and  $s_2$ , there exists a unique morphism  $g : Q \rightarrow D$  such that  $ge = f$ . Hence  $e$  is also the coequalizer of  $r_1$  and  $r_2$ .



Finally, we observe that  $(r_1, r_2)$  is the kernel pair of  $e$ , since by *Proposition A.0.8*, a kernel pair that admits a coequalizer is the kernel pair of its coequalizer.  $\square$

**Proposition A.0.10.** *If a morphism  $f$  admits a factorization  $f = mg$ , with  $m$  mono, then  $f$  and  $g$  share the same kernel pair.*

*Proof.* Let  $(E_q, r_1, r_2)$  be the kernel pair of  $f : A \rightarrow B$  and  $g : A \rightarrow Q$ , such that  $f = mg$  with  $m$  mono. We have that  $fr_1 = fr_2$ , hence  $mgr_1 = mgr_2$ .

Since  $m$  is a mono, we have that  $gr_1 = gr_2$ . Conversely, if  $(\text{Eq}, r_1, r_2)$  is the kernel pair of  $e$ , then we have that  $gr_1 = gr_2$ . Just composing with  $m$  we have that  $mgr_1 = mgr_2$ . Hence  $fr'_1 = fr'_2$ .

$$\begin{array}{ccc} A & \xrightarrow{f} & B \\ & \searrow g \quad \nearrow m & \\ & Q & \end{array}$$

Now we want to prove that if the universal property of the kernel pair holds for  $f$ , then it also holds for  $g$ . Let  $s_1 : X \rightarrow A$  and  $s_2 : X \rightarrow A$  be morphisms such that  $gs_1 = gs_2$ . Just composing with  $m$ , we have that  $mgs_1 = mgs_2$ . Hence  $fs_1 = fs_2$ . By the universal property of the kernel pair of  $f$ , there exists a unique morphism  $k : X \rightarrow \text{Eq}$  such that  $r_1k = s_1$  and  $r_2k = s_2$ . Conversely, consider  $s_1 : X \rightarrow A$  and  $s_2 : X \rightarrow A$  be morphisms such that  $fs_1 = fs_2$ , and suppose that the universal property holds for  $g$ . That is  $mgr_1 = mgr_2$ . Since  $m$  is a mono, we have that  $gr_1 = gr_2$ . By the universal property of the kernel pair of  $g$ , there exists a unique morphism  $k : X \rightarrow \text{Eq}$  such that  $r_1k = s_1$  and  $r_2k = s_2$ .

$$\begin{array}{ccccc} X & & & & \\ & \searrow k & & & \\ & \text{Eq} & \xrightarrow{r_2} & A & \\ & \downarrow r_1 & & \downarrow f & \\ & A & \xrightarrow{f} & B & \\ & & & & \downarrow m \\ & & & & Q \end{array}$$

Curved arrows:  $X \xrightarrow{s_1} A$ ,  $X \xrightarrow{s_2} A$ ,  $X \xrightarrow{g} Q$ ,  $A \xrightarrow{g} Q$ ,  $B \xrightarrow{g} Q$ .

□