

Homework 7 Writeup


Christian Diangco
AMATH 301 B

Problem 1

For each solution, as time increases, the pendulum swings slower and slower until its velocity eventually reaches zero.

Let's say velocity is positive going clockwise. When the initial conditions are $(\pi, 0.1)$, the pendulum starts from the top and starts swinging clockwise with a velocity of 0.1. When the initial conditions are $(\pi, -0.1)$, the pendulum also starts from the top, but starts swinging counter-clockwise with a velocity of -0.1.

When the initial conditions are $(2\pi, -3)$, the pendulum starts from the bottom and starts swinging counter-clockwise with a velocity of -3. When the initial conditions are $(-2\pi, 3)$, the pendulum also starts from the bottom, but starts swinging clockwise with a velocity of 3.




```
theta, v = np.meshgrid(np.linspace(-3*np.pi, 3*np.pi, 25), np.linspace(-3, 3, 25))
g = 9.8
L = 21
o = 0.15
dt = 0.01
tspan = np.arange(0, 50+dt, dt)
dthetadt = lambda theta, v: v
dvdt = lambda theta, v: -g / L * np.sin(theta) - o*v
odefun = lambda t, y: np.array([dthetadt(y[0], y[1]), dvdt(y[0], y[1])])
traj1 = scipy.integrate.solve_ivp(odefun, np.array([tspan[0], tspan[-1]]), np.array([np.pi, 0.1]), t_eval = tspan)
traj2 = scipy.integrate.solve_ivp(odefun, np.array([tspan[0], tspan[-1]]), np.array([np.pi, -0.1]), t_eval = tspan)
traj3 = scipy.integrate.solve_ivp(odefun, np.array([tspan[0], tspan[-1]]), np.array([2*np.pi, -3]), t_eval = tspan)
traj4 = scipy.integrate.solve_ivp(odefun, np.array([tspan[0], tspan[-1]]), np.array([-2*np.pi, 3]), t_eval = tspan)


fig, ax = plt.subplots()
ax.quiver(theta, v, dthetadt(theta, v), dvdt(theta, v))
ax.set_xlabel("$\\theta$", fontsize=20)
ax.set_ylabel("$v$", fontsize=20)
ax.set_title("$\\theta$ vs $v$", fontsize=20)
ax.plot(traj1.y[0,:], traj1.y[1,:])
ax.plot(traj2.y[0,:], traj2.y[1,:])
ax.plot(traj3.y[0,:], traj3.y[1,:])
ax.plot(traj4.y[0,:], traj4.y[1,:])
ax.set_xlim(-3*np.pi, 3*np.pi)
ax.set_ylim(-3, 3)
```

The error **increases** over time. From the plot we can see that with each pass over the equilibrium position, the lines get further and further apart from the RK45 line, which we are assuming to be the exact solution. The only one that doesn't appear to move away is the Leapfrog line, but that could be due to being too zoomed out.

θ vs v



Forward Euler

[illegible]

```

RK_sol = RK.y

fig2, ax2 = plt.subplots()
ax2.quiver(theta, v, dthetadt(theta, v), dvdt(theta, v))
ax2.set_xlabel("$\\theta$", fontsize=20)
ax2.set_ylabel("v", fontsize=20)
ax2.set_title("$\\theta$ vs v", fontsize=20)
ax2.plot(FE_sol[0:], FE_sol[1:], "c", linewidth=2, label="Forward Euler")
ax2.plot(BE_sol[0:], BE_sol[1:], "m", linewidth=2, label="Backward Euler")
ax2.plot(LF_sol[0:], LF_sol[1:], "r", linewidth=2, label="Leapfrog")
ax2.plot(RK_sol[0:], RK_sol[1:], "y:", linewidth=2, label="RK45")
ax2.legend()

```