



**UNIVERSITÀ
DEGLI STUDI
DI TRIESTE**

From Latent to Deep Latent Variable Models


A Study On Causal Effect Inference with CEVAE

Valeria De Stasio, Christian Faccio, Giovanni Lucarelli

June 17, 2025

Objective: estimating causal effects

Estimate how a medical treatment T affects the health Y of a *random* patient.



$$P(Y = y | \text{do}(T = t)) = P(Y = y | T = t)$$

$$\text{ITE} = \mathbb{E}[(Y | \text{do}(T = 1))] - \mathbb{E}[(Y | \text{do}(T = 0))]$$

This is usually the case in a **randomized controlled trial** (RCT), where the treatment is randomly assigned to the patients.


Confounder

In a observational study:

- **no** control over the treatment T assignment
→ there may be a confounder X (e.g. Income) that influences both variables

$$P(Y|\text{do}(t)) \neq P(Y|t)$$


$$P(Y|\text{do}(t), x) = P(Y|t, x)$$



If the confounder is observed even a linear regression can do the job!

Latent confounder

But what if the confounder Z (e.g. Wealth) is **not** observed and the observed X is only a proxy of it?



Idea: estimate the latent variable and condition on it

Vanilla Latent Variable Model


- Generative model:

$$Z \sim \mathcal{N}(0, 1)$$

$$X_j \sim \mathcal{N}(az + b, \text{diag}(\sigma_x^2))$$

$$T \sim \text{Be}(\sigma(cz))$$

$$Y \sim \mathcal{N}(et + fz, \sigma_Y^2)$$



- Guide:** mean-field assumption over the observations
- Problem:** we need to train a new model at test time!

Latent Variable Model (LVM):


- Linear functions
- SVI + mini-SVI for new x
- Small number of parameters
- High interpretability

Deep Latent Variable Model (CEVAE):


- MLP functions
- SVI + Amortized inference
- Large number of parameters
- High flexibility

CEVAE = LVM + MLPs + Amortized Inference
same causal idea, more flexibility and speed at test time.

Deep Latent Variable Model: CEVAE



Synthetic linear dataset



$$Z \sim \mathcal{N}(0, 1)$$

$$a_j \sim \mathcal{U}(-10, 10)$$


$$X_j \sim \mathcal{N}(a_j Z, \sigma_X^2)$$

$$T|Z \sim \text{Be}(\sigma(\beta Z))$$

$$Y|T, Z \sim \mathcal{N}(z + t, \sigma_Y^2)$$

Remark: ITE is constant for all x !

Synthetic non linear dataset



$$Z \sim \mathcal{N}(0, 1)$$

$$a \sim \mathcal{U}(-10, 10)^d$$

$$X_1, \dots, X_d \sim \mathcal{N}(a \tanh(z), \sigma_X^2 \mathbb{I})$$


$$T|Z \sim \text{Be}(\sigma(\beta z))$$

$$Y|T, Z \sim \mathcal{N}(\text{NonLin}(z, t), \sigma_Y^2)$$


$$\text{NonLin}(z, t) = \sin(z) + \frac{1}{2}z + t \left(1 + \frac{1}{2}z\right)$$

Remark: ITE depends on interaction between t and z !


Results (1/2): Linear Dataset




Results (2/2): Non Linear Dataset




Internal Representation: Linear Dataset




Internal Representation: Non Linear Dataset




Experiment 1: Increasing the sample size




Experiment 2: Increasing decorrelation among proxies




Experiment 3: Increasing latent dimension



Experiment 4: latent distribution misspecified



Experiment 5: Increasing the number of proxies



Conclusions

- **Linear synthetic data** – Linear models *benefit* from strong prior assumptions and recover the true effects almost perfectly; CEVAE offers no gain and adds variance.
- **Non-linear synthetic data** – With a tuned latent dimension, CEVAE outperforms linear baselines; if untuned it over-fits and the latent loses causal meaning.
- **Practical takeaway** – Use simple models when you can state strong causal assumptions; switch to CEVAE to relax those assumptions—provided you can afford heavier training and tuning.
- **Next steps** – Test on real datasets, explore more sophisticated variant of CEVAE (e.g. *TEDVAE*, *DCEVAE*, *ICEVAE*)

Thank You!