

## COSMIC REIONIZATIONS AND 21CM OBSERVATIONS

The redshifted 21cm line of neutral hydrogen (HI) is an incisive probe of the high-redshift universe, allowing in principle for creation of full spatial 3D-maps of hydrogen as a function of cosmic time in the redshift range  $z \sim 6 - 200$  (Furlanetto et al., 2006; Pritchard and Loeb, 2012). The HI line is redshifted to frequencies below 200MHz and can therefore be observed by radio telescopes (Tozzi et al., 2000). Beyond maps, this cosmic signature will test fundamental physics from the Dark Ages, the formation of the first compact radiation sources at Cosmic Dawn, and the reionization of neutral hydrogen with galaxy formation during the Epoch of reionization (Barkana, 2016).

First-generation radio telescopes have targeted two summary statistics of the 21cm line: the all-sky average brightness temperature (the "global" signal) from Cosmic Dawn as a function of redshift, and the power spectrum of spatial fluctuations as a function of Fourier wavemodes and redshift. As expected, different physics is probed at different redshifts: During the *Dark Ages*, between recombination and the formation of the first astrophysical objects, the baryonic perturbations remain linear and the physics is simple enough that the 21cm intensity directly can be traced back to cosmological parameters and for example break the degeneracy between the neutrino mass and the CMB optical depth (Pritchard and Pierpaoli, 2008), distinguish inflationary models (Masui and Pen, 2010), and test various physics beyond the standard model including neutrino decay (Chianese et al., 2019) and dark matter annihilation (Koopmans et al., 2021). The *Cosmic Dawn* era probes astrophysics like the star formation rate, X-ray SEDs, and the mass of star-forming halos (e.g. Barkana (2016) and Bye et al. (2022)). Finally, the Epoch of Reionization constrains the ionization history of hydrogen, the mean free path of ionizing photons in Stromgren spheres, the CMB optical depth, and X-ray heating (DeBoer et al., 2017).

Experiments targeting the global signal include the Experiment to Detect the Global EoR Signature (EDGES, Bowman and Rogers, 2010; Monsalve et al., 2017, 2018, 2019), SARAS (Shaped Antenna measurement of the background RADIO Spectrum, Singh et al., 2022), REACH (Radio Experiment for the Analysis of Cosmic Hydrogen, de Lera Acedo et al., 2022), PRIZM (Probing Radio Intensity at high-Z from Marion, Philip et al., 2019), MIST (Mapper of the IGM Spin Temperature, <http://www.physics.mcgill.ca/mist/>), and DAPPER (Dark Ages Polarimeter Pathfinder, Burns et al., 2019). These experiments have only recently reached the necessary sensitivity to observe cosmological signals, but EDGES have put a lower limit on the length of the Epoch of Reionization (Bowman and Rogers, 2010) and has made a claimed detection of the global signal (Bowman et al., 2018). This claim has subsequently been countered by a claim of non-detection (Singh et al., 2022), but if truly of cosmic origin, the measured absorption profile can likely not be explained by standard physics as it suggests cooling of the hydrogen gas faster than what can be achieved with adiabatic cooling due to the expansion of the universe. So-called exotic models—which go beyond  $\Lambda$ CDM—have been proposed as explanations for the EDGES detection, including charged Dark Matter (Muñoz et al., 2018), dark matter-baryon interactions (Barkana et al., 2018; Barkana, 2018), and early Black Holes (Ewall-Wice et al., 2018).

For the power spectrum, upper limits are being placed (The HERA Collaboration et al., 2022; Abdurashidova et al., 2022), which constrain early heating of the IGM, but no claim of the same magnitude as EDGES have been made.

Future plans for the field include improved sensitivity of current telescopes, new earth-based telescopes like the Square Kilometer Array (Gupta, 2022), and pathfinders for space based missions (e.g. LuSEE, Bale et al. (2020)).

## References

- Z. Abdurashidova, J. E. Aguirre, P. Alexander, Z. S. Ali, Y. Balfour, A. P. Beardsley, G. Bernardi, T. S. Billings, J. D. Bowman, R. F. Bradley, P. Bull, J. Burba, S. Carey, C. L. Carilli, C. Cheng, D. R. DeBoer, M. Dexter, E. de Lera Acedo, T. Dibblee-Barkman, J. S. Dillon, J. Ely, A. Ewall-Wice, N. Fagnoni, R. Fritz, S. R. Furlanetto, K. Gale-Sides, B. Glendenning, D. Gorthi, B. Greig, J. Grobbelaar, Z. Halday, B. J. Hazelton, J. N. Hewitt, J. Hickish, D. C. Jacobs, A. Julius, N. S. Kern, J. Kerrigan, P. Kittiwisit, S. A. Kohn, M. Kolopanis, A. Lanman, P. La Plante, T. Lekalake, D. Lewis, A. Liu, D. MacMahon, L. Malan, C. Malgas, M. Maree, Z. E. Martinot, E. Matsetela, A. Mesinger, M. Molewa, M. F. Morales, T. Mosiane, S. G. Murray, A. R. Neben, B. Nikolic, C. D. Nunhokee, A. R. Parsons, N. Patra, R. Pascua, S. Pieterse, J. C. Pober, N. Razavi-Ghods, J. Ringuette, J. Robnett, K. Rosie, P. Sims, S. Singh, C. Smith, A. Syce, N. Thyagarajan, P. K. G. Williams, H. Zheng, and HERA Collaboration. First Results from HERA Phase I: Upper Limits on the Epoch of Reionization 21 cm Power Spectrum. , 925(2):221, Feb. 2022. doi: 10.3847/1538-4357/ac1c78.
- S. D. Bale, J. W. Bonnell, J. Burns, K. Goetz, J. S. Halekas, D. M. Malaspina, B. Page, M. Pulupa, A. R. Poppe, R. J. MacDowall, M. Maksimovic, and A. Zaslavsky. The Lunar Surface Electromagnetics Experiment (LuSEE). In *Lunar Surface Science Workshop*, volume 2241 of *LPI Contributions*, page 5001, May 2020.
- R. Barkana. The rise of the first stars: Supersonic streaming, radiative feedback, and 21-cm cosmology. , 645:1–59, July 2016. doi: 10.1016/j.physrep.2016.06.006.
- R. Barkana. Possible interaction between baryons and dark-matter particles revealed by the first stars. , 555(7694):71–74, Mar. 2018. doi: 10.1038/nature25791.
- R. Barkana, N. J. Outmezguine, D. Redigol, and T. Volansky. Strong constraints on light dark matter interpretation of the EDGES signal. , 98:103005, Nov. 2018. doi: 10.1103/PhysRevD.98.103005.
- J. D. Bowman and A. E. E. Rogers. A lower limit of  $\Delta z \gtrsim 0.06$  for the duration of the reionization epoch. , 468(7325):796–798, Dec. 2010. doi: 10.1038/nature09601.
- J. D. Bowman, A. E. E. Rogers, R. A. Monsalve, T. J. Mozdzen, and N. Mahesh. An absorption profile centred at 78 megahertz in the sky-averaged spectrum. , 555(7694):67–70, Mar. 2018. doi: 10.1038/nature25792.
- J. O. Burns, S. Bale, and R. F. Bradley. Dark Cosmology: Investigating Dark Matter and Exotic Physics using the Redshifted 21-cm Global Signal with the Dark Ages Polarimeter Pathfinder (DAPPER). In *American Astronomical Society Meeting Abstracts #234*, volume 234 of *American Astronomical Society Meeting Abstracts*, page 212.02, June 2019.
- C. H. Bye, S. K. N. Portillo, and A. Fialkov. 21cmVAE: A Very Accurate Emulator of the 21 cm Global Signal. , 930(1):79, May 2022. doi: 10.3847/1538-4357/ac6424.
- M. Chianese, P. Di Bari, K. Farrag, and R. Samanta. Probing relic neutrino radiative decays with 21 cm cosmology. *Physics Letters B*, 790:64–70, Mar. 2019. doi: 10.1016/j.physletb.2018.09.040.
- E. de Lera Acedo, D. I. L. de Villiers, N. Razavi-Ghods, W. Handley, A. Fialkov, A. Magro, D. Anstey, H. T. J. Bevins, R. Chiello, J. Cumner, A. T. Josaitis, I. L. V. Roque, P. H. Sims, K. H. Scheutwinkel, P. Alexander, G. Bernardi, S. Carey, J. Cavillot, W. Croukamp, J. A. Ely,

- T. Gessey-Jones, Q. Gueuning, R. Hills, G. Kulkarni, R. Maiolino, P. D. Meerburg, S. Mittal, J. R. Pritchard, E. Puchwein, A. Saxena, E. Shen, O. Smirnov, M. Spinelli, and K. Zarb-Adami. The REACH radiometer for detecting the 21-cm hydrogen signal from redshift  $z \approx 7.5$ –28. *Nature Astronomy*, 6:984–998, July 2022. doi: 10.1038/s41550-022-01709-9.
- D. R. DeBoer, A. R. Parsons, J. E. Aguirre, P. Alexander, Z. S. Ali, A. P. Beardsley, G. Bernardi, J. D. Bowman, R. F. Bradley, C. L. Carilli, C. Cheng, E. de Lera Acedo, J. S. Dillon, A. Ewall-Wice, G. Fadana, N. Fagnoni, R. Fritz, S. R. Furlanetto, B. Glendenning, B. Greig, J. Grobbelaar, B. J. Hazelton, J. N. Hewitt, J. Hickish, D. C. Jacobs, A. Julius, M. Kariseb, S. A. Kohn, T. Lekalake, A. Liu, A. Loots, D. MacMahon, L. Malan, C. Malgas, M. Maree, Z. Martinot, N. Mathison, E. Matsetela, A. Mesinger, M. F. Morales, A. R. Neben, N. Patra, S. Pieterse, J. C. Pober, N. Razavi-Ghods, J. Ringuette, J. Robnett, K. Rosie, R. Sell, C. Smith, A. Syce, M. Tegmark, N. Thyagarajan, P. K. G. Williams, and H. Zheng. Hydrogen Epoch of Reionization Array (HERA). , 129(974):045001, Apr. 2017. doi: 10.1088/1538-3873/129/974/045001.
- A. Ewall-Wice, T. C. Chang, J. Lazio, O. Doré, M. Seiffert, and R. A. Monsalve. Modeling the Radio Background from the First Black Holes at Cosmic Dawn: Implications for the 21 cm Absorption Amplitude. , 868(1):63, Nov. 2018. doi: 10.3847/1538-4357/aae51d.
- S. R. Furlanetto, S. P. Oh, and F. H. Briggs. Cosmology at low frequencies: The 21 cm transition and the high-redshift Universe. , 433(4-6):181–301, Oct. 2006. doi: 10.1016/j.physrep.2006.08.002.
- Y. Gupta. The Square Kilometre Array: current status and India’s role in this upcoming facility. *Journal of Astrophysics and Astronomy*, 43(2):68, Dec. 2022. doi: 10.1007/s12036-022-09856-1.
- L. V. E. Koopmans, R. Barkana, M. Bentum, G. Bernardi, A.-J. Boonstra, J. Bowman, J. Burns, X. Chen, A. Datta, H. Falcke, A. Fialkov, B. Gehlot, L. Gurvits, V. Jelić, M. Klein-Wolt, J. Lazio, D. Meerburg, G. Mellema, F. Mertens, A. Mesinger, A. Offringa, J. Pritchard, B. Semelin, R. Subrahmanyan, J. Silk, C. Trott, H. Vedantham, L. Verde, S. Zaroubi, and P. Zarka. Peering into the dark (ages) with low-frequency space interferometers. *Experimental Astronomy*, 51(3): 1641–1676, June 2021. doi: 10.1007/s10686-021-09743-7.
- K. W. Masui and U.-L. Pen. Primordial Gravity Wave Fossils and Their Use in Testing Inflation. , 105(16):161302, Oct. 2010. doi: 10.1103/PhysRevLett.105.161302.
- R. A. Monsalve, A. E. E. Rogers, J. D. Bowman, and T. J. Mozdzen. Results from EDGES High-band. I. Constraints on Phenomenological Models for the Global 21 cm Signal. , 847(1):64, Sept. 2017. doi: 10.3847/1538-4357/aa88d1.
- R. A. Monsalve, B. Greig, J. D. Bowman, A. Mesinger, A. E. E. Rogers, T. J. Mozdzen, N. S. Kern, and N. Mahesh. Results from EDGES High-band. II. Constraints on Parameters of Early Galaxies. , 863(1):11, Aug. 2018. doi: 10.3847/1538-4357/aace54.
- R. A. Monsalve, A. Fialkov, J. D. Bowman, A. E. E. Rogers, T. J. Mozdzen, A. Cohen, R. Barkana, and N. Mahesh. Results from EDGES High-Band. III. New Constraints on Parameters of the Early Universe. , 875(1):67, Apr. 2019. doi: 10.3847/1538-4357/ab07be.
- J. B. Muñoz, C. Dvorkin, and A. Loeb. 21-cm Fluctuations from Charged Dark Matter. , 121(12): 121301, Sept. 2018. doi: 10.1103/PhysRevLett.121.121301.

- L. Philip, Z. Abdurashidova, H. C. Chiang, N. Ghazi, A. Gumba, H. M. Heilgendorff, J. M. Jáuregui-García, K. Malepe, C. D. Nunhokee, J. Peterson, J. L. Sievers, V. Simes, and R. Spann. Probing Radio Intensity at High-Z from Marion: 2017 Instrument. *Journal of Astronomical Instrumentation*, 8(2):1950004, Jan. 2019. doi: 10.1142/S2251171719500041.
- J. R. Pritchard and A. Loeb. 21 cm cosmology in the 21st century. *Reports on Progress in Physics*, 75(8):086901, Aug. 2012. doi: 10.1088/0034-4885/75/8/086901.
- J. R. Pritchard and E. Pierpaoli. Constraining massive neutrinos using cosmological 21cm observations. , 78(6):065009, Sept. 2008. doi: 10.1103/PhysRevD.78.065009.
- S. Singh, N. T. Jishnu, R. Subrahmanyam, N. Udaya Shankar, B. S. Girish, A. Raghunathan, R. Somashekar, K. S. Srivani, and M. Sathyanarayana Rao. On the detection of a cosmic dawn signal in the radio background. *Nature Astronomy*, 6:607–617, Feb. 2022. doi: 10.1038/s41550-022-01610-5.
- The HERA Collaboration, Z. Abdurashidova, T. Adams, J. E. Aguirre, P. Alexander, Z. S. Ali, R. Baartman, Y. Balfour, R. Barkana, A. P. Beardsley, G. Bernardi, T. S. Billings, J. D. Bowman, R. F. Bradley, D. Breitman, P. Bull, J. Burba, S. Carey, C. L. Carilli, C. Cheng, S. Choudhuri, D. R. DeBoer, E. de Lera Acedo, M. Dexter, J. S. Dillon, J. Ely, A. Ewall-Wice, N. Fagnoni, A. Filkov, R. Fritz, S. R. Furlanetto, K. Gale-Sides, H. Garsden, B. Glendenning, A. Gorce, D. Gorthi, B. Greig, J. Grobbelaar, Z. Haldy, B. J. Hazelton, S. Heimersheim, J. N. Hewitt, J. Hickish, D. C. Jacobs, A. Julius, N. S. Kern, J. Kerrigan, P. Kittiwisit, S. A. Kohn, M. Kolopanis, A. Lanman, P. La Plante, D. Lewis, A. Liu, A. Loots, Y.-Z. Ma, D. H. E. MacMahon, L. Malan, K. Malgas, C. Malgas, M. Maree, B. Marero, Z. E. Martinot, L. McBride, A. Mesinger, J. Mirocha, M. Molewa, M. F. Morales, T. Mosiane, J. B. Muñoz, S. G. Murray, V. Nagpal, A. R. Neben, B. Nikolic, C. D. Nunhokee, H. Nuwegeld, A. R. Parsons, R. Pascua, N. Patra, S. Pieterse, Y. Qin, N. Razavi-Ghods, J. Robnett, K. Rosie, M. G. Santos, P. Sims, S. Singh, C. Smith, H. Swarts, N. Thyagarajan, M. J. Wilensky, P. K. G. Williams, P. van Wyngaarden, and H. Zheng. Improved Constraints on the 21 cm EoR Power Spectrum and the X-Ray Heating of the IGM with HERA Phase I Observations. *arXiv e-prints*, art. arXiv:2210.04912, Oct. 2022.
- P. Tozzi, P. Madau, A. Meiksin, and M. J. Rees. Radio Signatures of H I at High Redshift: Mapping the End of the “Dark Ages”. , 528(2):597–606, Jan. 2000. doi: 10.1086/308196.