

ATLAS 13 TeV Data Analysis using Particle Flow

Christian Kirfel

Bachelorarbeit in Physik
angefertigt im Physikalischen Institut

vorgelegt der
Mathematisch-Naturwissenschaftlichen Fakultät
der
Rheinischen Friedrich-Wilhelms-Universität
Bonn

November 2016

DRAFT

Ich versichere, dass ich diese Arbeit selbstständig verfasst und keine anderen als die angegebenen Quellen und Hilfsmittel benutzt sowie die Zitate kenntlich gemacht habe.

Bonn,
Datum

.....
Unterschrift

- 1. Gutachter: Prof. Dr. John Smith
- 2. Gutachterin: Prof. Dr. Anne Jones

Acknowledgements

I would like to thank ...

You should probably use `\chapter*` for acknowledgements at the beginning of a thesis and `\chapter` for the end.

DRAFT

DRAFT

Contents

1	Introduction	1
2	Theoretical and experimental basics	3
2.1	The Standard Model of Particle Physics	3
2.2	The LHC and ATLAS	4
2.2.1	The LHC	4
2.2.2	The ATLAS Detector	4
2.3	The Particle Flow Algorithm	4
2.3.1	Track selection	6
2.3.2	Clustering	6
2.3.3	Matching Track to Cluster	6
2.3.4	Cell Subtraction	6
2.4	Calibration in Particle Flow and its difficulties at the time	7
3	Results	9
	Bibliography	11
A	Useful information	13
	List of Figures	15
	List of Tables	17

DRAFT

Introduction

At the end of the introduction it is normal to say briefly what comes in the following chapters.

The lines at the end of this file are used by AUCTeX to specify which is the master \LaTeX file, so that you can compile your thesis directly within emacs.

Theoretical and experimental basics

2.1 The Standard Model of Particle Physics

The Standard Model of Particle Physics summarizes the current knowledge of fundamental particles and their interactions. The model holds at scales of 1 fm and below. Gravity, being the fourth fundamental force is not included because it is negligible for most phenomena at this scale.

The current view is that all matter is made out of three kinds of elementary particle being leptons quarks and mediators. There are six leptons falling into three families according to charge, electron number, muon number and tau number. To each of these leptons there also is an antilepton with all signs reversed. Similiar to that there are six flavors of quarks separated by Strangeness (S), charm (C), Beauty (B), and truth (T). As well as the leptons the quarks fall into three generations. For both kinds of particles the mass rises with the generations and every generation comes as a doublet. The first particle of each lepton doublet is not charged and referred to as a neutrino while the second particle has charge -1. For each quark doublet there is an element with fractional charge $-\frac{1}{3}$ and an element with fractional charge $\frac{2}{3}$. To every of these particles there is an anti particle with opposite charge.

The third kind of particle included in the standard model is the mediator. Mediators are gauge bosons whose exchange allows the particles to interact. There are four kinds of of elemtary interactions of which the strong electromagnetic and weak interaction are included in the model. The fourth interaction is the gravitational interaction. The gauge particles for the strong interaction are the gluons carrying colour charge, the electromagnetic mediator is the photon (γ) and the weak mediators are the W^\pm and Z bosons

Table 2.1: Lepton properties

	symbol	Charge Q	L_e	L_μ	L_τ
First generation{	e	-1	1	0	0
	ν_e	0	1	0	0
Second generation{	μ	-1	0	1	0
	ν_μ	0	0	1	0
Third generation{	τ	-1	0	0	1
	ν_τ	0	0	0	1

Table 2.2: Quark properties

	Symbol	Charge Q	mass	D	U	S	C	B	T
First generation {	d	$-\frac{1}{3}$	4.8MeV	-1	0	0	0	0	0
	u	$\frac{2}{3}$	2.3MeV	0	1	0	0	0	0
Second generation {	s	$-\frac{1}{3}$	95MeV	0	0	-1	0	0	0
	c	$\frac{2}{3}$	1275MeV	0	0	0	1	0	0
Third generation {	b	$-\frac{1}{3}$	4180MeV	0	0	0	0	-1	0
	t	$\frac{2}{3}$	173210MeV	0	0	0	0	0	1

Table 2.3: Mediator properties

Interaction	Theory	Mediator	Charge	Coupling
Strong	QCD	gluons (8)	colour	1
Electromagnetic	QED	photon γ	electric charge	10^{-1}
Weak	GSW	W^{\pm}, Z	weak isospin	20^{-6}

2.2 The LHC and ATLAS

The analysis for this thesis has been performed in the ATLAS collaboration. The ATLAS-Detector is one of the four big experiments at the LHC at Cern. Therefore this chapter gives a brief overview over the LHC and ATLAS focusing on the properties directly relevant for Particle Flow Analysis.

After a description of the ATLAS detector in general I will give a little bit more information about the detector components directly relevant for the explanation of particle flow being the tracking detector and the calorimeter.

2.2.1 The LHC

The Large Hadron Collider ("LHC") at CERN was built to extend the frontiers of modern particle physics by delivering high luminosities and unprecedented high energies.

The LHC is designed to collide bunches of up to 10^{11} protons

2.2.2 The ATLAS Detector

2.2.3 Tracking detectors

2.2.4 Calorimeter

2.3 The Particle Flow Algorithm

Recently only either the Calorimeter or the tracker information was used to reconstruct Jets in ATLAS events. The Particle Flow Algorithm combines tracker and calorimeter information to achieve better resolution especially at lower energies. The main advantages of including the tracker information into reconstruction are listed here:

- For low energy charged particles the momentum resolution of the tracking detector is superior to the calorimeter.




Figure 2.1: Sketch of the LHC ring, the position of the experiments and the surrounding countryside. The four big LHC experiments are indicated. The location of the injection lines and the SPS are also shown.




Figure 2.2: Sketch of the ATLAS detector [1]

- The Tracking detector is able to reconstruct soft particles, which would not pass the noise threshold of the calorimeter.
- The ATLAS tracking detector has a superior angular resolution for single charged particles.
- Low p_T charged particles may be swept out of the cone before reaching the calorimeter by the magnetic field. The tracker information allows to cluster these particles into the jet.

The advantages of Particle Flow have already been shown for Run 1 data in




Figure 2.3: Flowchart of the steps of the Particle Flow algorithm

2.3.1 Track selection

2.3.2 Clustering

The algorithm uses information from the tracker and the calorimeter as input. The information from the calorimeter is given as topological clusters. The construction of these clusters is briefly described in this chapter to give the reader a basic understanding which input information the algorithm uses. Every cluster is being constructed around a so called seed cell. A seed cell is a cell for which the deposited energy exceeds the expected noise by four times the standard deviation. If a seed is found all the neighboring cells which exceed the noise by at least two times the standard deviation are added to the cluster. Lastly all the cells neighboring these clusters are also added.

2.3.3 Matching Track to Cluster

The algorithm tries to match every track to one or more calorimeter clusters. First the algorithm tries to match a single best-match topo-cluster to every selected track. To do so first the distances in $\Delta\phi$ and $\Delta\eta$ from the track are extrapolated to the second layer of the EM calorimeter and then to the topo cluster. Then the topo-clusters get ranked based on the metric:

$$\Delta R' = \sqrt{\left(\frac{\Delta\phi}{\sigma_\phi}\right)^2 + \left(\frac{\Delta\eta}{\sigma_\eta}\right)^2} \quad (2.1)$$

where σ_η and σ_ϕ refer to the angular topo-cluster width, computed from the standard deviation of the displacements of the topo clusters

2.3.4 Cell Subtraction

The last step in the Particle Flow algorithm after matching a set of topo-clusters to a track is the cell-wise subtraction of energy deposits to remove remnants and determine which energy deposits belong to the given particle. If the energy deposited in the set of clusters falls below the expected energy the clusters are simply removed. Otherwise, a cell by cell subtraction is performed.

The first step of the cell subtraction is generating a shower shape from the extrapolated track. Around the extrapolated track rings in η , ϕ space are generated, being just wide enough to independently contain at least one cell from the extrapolated position. Furthermore the rings are restricted to one layer and for each layer have the same radial size. After the generation of rings in each layer the average energy density in each ring is computed and the rings are ranked by energy density in descending order. The layer is not used in any way for this ranking. The subtraction then starts from the ring with highest energy density and proceeds successively to rings in lower order until the next ring's energy exceeds or falls below. If the ring's energy exceeds the energy still to be subtracted the energy in each cell is scaled down by the fraction needed to reach the expected energy then the process halts and the remaining cells are removed as remnants. An example of the process is sketched in *fasofoifjaor*

2.4 Calibration in Particle Flow and its difficulties at the time

Results

I know nothing

Bibliography

- [1] A. Collaboration, *A (Not So) Short Introduction to LaTeX2e*,
URL: <http://atlas.cern/resources/multimedia> (cit. on p. 5).

APPENDIX **A**

Useful information

In the appendix you usually include extra information that should be documented in your thesis, but not interrupt the flow.

List of Figures

2.1	Sketch of the LHC ring, the position of the experiments and the surrounding countryside.	5
2.2	Sketch of the ATLAS detector	5
2.3	Flowchart of the steps of the Particle Flow algorithm	6

List of Tables

2.1	Lepton properties	3
2.2	Quark properties	4
2.3	Mediator properties	4