

# **ATLAS 13 TeV Data Analysis using Particle Flow**

Christian Kirfel

Bachelorarbeit in Physik  
angefertigt im Physikalischen Institut

vorgelegt der  
Mathematisch-Naturwissenschaftlichen Fakultät  
der  
Rheinischen Friedrich-Wilhelms-Universität  
Bonn

November 2016

DRAFT

Ich versichere, dass ich diese Arbeit selbstständig verfasst und keine anderen als die angegebenen Quellen und Hilfsmittel benutzt sowie die Zitate kenntlich gemacht habe.

Bonn, .....  
Datum

.....  
Unterschrift

- 1. Gutachter: Prof. Dr. John Smith
- 2. Gutachterin: Prof. Dr. Anne Jones

# Acknowledgements

---

I would like to thank ...

You should probably use `\chapter*` for acknowledgements at the beginning of a thesis and `\chapter` for the end.

DRAFT

DRAFT

# Contents

---

<b>1</b>	<b>Theoretical and experimental basics</b>	<b>1</b>
1.1	The Standard Model of Particle Physics . . . . .	1
1.2	The LHC and ATLAS . . . . .	2
1.2.1	The LHC . . . . .	2
1.2.2	The ATLAS Detector . . . . .	2
1.3	The Particle Flow Algorithm . . . . .	2
1.4	Calibration in Particle Flow and its difficulties at the time . . . . .	3
<b>2</b>	<b>Results</b>	<b>5</b>
	<b>Bibliography</b>	<b>7</b>
<b>A</b>	<b>Useful information</b>	<b>9</b>
	<b>List of Figures</b>	<b>11</b>
	<b>List of Tables</b>	<b>13</b>

DRAFT

## Theoretical and experimental basics

### 1.1 The Standard Model of Particle Physics

The Standard Model of Particle Physics summarizes the current knowledge of fundamental particles and their interactions. The model holds at scales of 1 fm and below. Gravity, being the fourth fundamental force is not included because it is negligible for most phenomena at this scale. The current view is that all matter is made out of three kinds of elementary particle being leptons quarks and mediators. There are six leptons falling into three families according to charge, electron number, muon number and tau number. To each of these leptons there also is an antilepton with all signs reversed. Similiar to that there are six flavors of quarks separated by Strangeness (S), charm (C), Beauty (B), and truth (T). As well as the leptons the quarks fall into three generations.

Table 1.1: My caption

	symbol	Charge $Q$	$L_e$	$L_\mu$	$L_\tau$
First generation{	$e$	-1	1	0	0
	$\nu_e$	0	1	0	0
Second generation{	$\mu$	-1	0	1	0
	$\nu_\mu$	0	0	1	0
Third generation{	$\tau$	-1	0	0	1
	$\nu_\tau$	0	0	0	1

Table 1.2: My caption

	Symbol	Charge $Q$	D	U	S	C	B	T
First generation {	$d$	$-\frac{1}{3}$	-1	0	0	0	0	0
	$u$	$\frac{2}{3}$	0	1	0	0	0	0
Second generation {	$s$	$-\frac{1}{3}$	0	0	-1	0	0	0
	$c$	$\frac{2}{3}$	0	0	0	1	0	0
Third generation {	$b$	$-\frac{1}{3}$	0	0	0	0	-1	0
	$t$	$\frac{2}{3}$	0	0	0	0	0	1

## 1.2 The LHC and ATLAS

The analysis for this thesis has been performed in the ATLAS collaboration. The ATLAS-Detector is one of the four big experiments at the LHC at Cern. Therefore this chapter gives a brief overview over the LHC and ATLAS focusing on the properties directly relevant for Particle Flow Analysis.

### 1.2.1 The LHC

The Large Hadron Collider ("LHC") at CERN was built to extend the frontiers of modern particle physics by delivering high luminosities and unprecedented high energies.

The LHC is designed to collide bunches of up to  $10^{11}$  protons




Figure 1.1: Sketch of the LHC ring, the position of the experiments and the surrounding countryside. The four big LHC experiments are indicated. The location of the injection lines and the SPS are also shown.

### 1.2.2 The ATLAS Detector

## 1.3 The Particle Flow Algorithm

Recently only either the Calorimeter or the tracker information was used to reconstruct Jets in ATLAS events. The Particle Flow Algorithm combines tracker and calorimeter information to achieve better resolution especially at lower energies. The main advantages of including the tracker information into reconstruction are listed here:

- For low energy charged particles the momentum resolution of the tracking detector is superior to the calorimeter.






Figure 1.2: Sketch of the ATLAS detector

- The Tracking detector is able to reconstruct soft particles, which would not pass the noise threshold of the calorimeter.
- The ATLAS tracking detector has a superior angular resolution for single charged particles.
- Low  $p_T$  charged particles may be swept out of the cone before reaching the calorimeter by the magnetic field. The tracker information allows to cluster these particles into the jet.

## 1.4 Calibration in Particle Flow and its difficulties at the time



# Results

---

I know nothing



# Bibliography

---

- [1] D. Griffiths, *Introducton to Elementary Particles*, 2nd ed., WILEY-VCH, 2008.



## APPENDIX **A**

---

### **Useful information**

---

In the appendix you usually include extra information that should be documented in your thesis, but not interrupt the flow.





# List of Figures

---

- 1.1 Sketch of the LHC ring, the position of the experiments and the surrounding countryside. 2
- 1.2 Sketch of the LHC ring, the position of the experiments and the surrounding countryside. 3



# List of Tables

---

1.1	My caption	1
1.2	My caption	1