

Article 0.11: Scope and Limits of the Spectral Reduction Lemma – A Pedagogical Synthesis (Revised Edition – 40 Problems)

Christian Kithima
Independent Researcher, Kinshasa
christiankithimaomba@gmail.com
WhatsApp: +243995176701
Telegram: +243818136082
ORCID: 0009-0003-3829-8519

GitHub: <https://github.com/christiankithimaomba-cyber/spectral-reduction-framework>

May 2026

Abstract

We clarify the scope and limits of the Spectral Reduction Lemma (Kithima's lemma). The lemma provides a universal method to solve discrete decision problems, but it is not a panacea. We summarise the conditions under which a problem is amenable to the four core strategies (CD, EB, FV, FD) and the advanced strategies (SRH, SRP, SUTL, SHS). We list the forty problems already resolved, explain why some famous open problems (Collatz, Hodge, odd perfect numbers) remain out of reach or require additional breakthroughs, and identify classes of problems that are fundamentally incompatible (e.g., smooth Poincaré in dimension 4, Iliev-Sendov, most Erdős conjectures). Crucially, we emphasise the distinction between **theoretical decidability** (guaranteed by the lemma) and **practical feasibility** (not guaranteed). The lemma is a mathematical certification tool, not an engineering blueprint. This pedagogical synthesis helps researchers assess whether their problem may be solvable by the spectral method and what caveats apply.

Contents

1	Introduction	2
2	The four core strategies: conditions for success	2
3	Advanced strategies: status and conditions	3
3.1	Strategies that have produced complete proofs (accepted)	3
3.2	Strategies still open (conditions not met)	3
3.3	The Hilbert-Pólya conjecture	3
3.4	The Navier-Stokes spectral approach (incomplete)	3
4	Domains successfully covered by the lemma	4
5	What makes a problem amenable to the core strategies?	4
6	Domains that are out of reach (with explanations)	5
6.1	Potentially approachable via open advanced strategies	5
6.2	Problems fundamentally incompatible with the spectral framework	5

7	The crucial nuance: theoretical vs. practical feasibility	6
8	Conclusion	6

1 Introduction

The Spectral Reduction Lemma (Kithima's lemma) and its algorithmic completeness theorem provide a powerful method for solving discrete decision problems via ground state extraction. However, not every problem can be encoded to satisfy the four pillars. Moreover, even for problems that can be encoded, the theoretical guarantee of polynomial-time decidability does **not** imply practical feasibility on current or foreseeable hardware. This article clarifies the scope and limits of the lemma by:

- Summarising the conditions under which a problem is amenable to the four core strategies (CD, EB, FV, FD).
- Listing the **forty problems** already resolved in their definitive order.
- Explaining why advanced strategies (FM, DUST, SFF, Backward Transfer, CTP) remain open, while SRH, SRP, SUTL, SHS are complete.
- Discussing the incomplete spectral approach to Navier-Stokes as an example of a promising but not yet definitive line of research.
- Identifying classes of problems that are fundamentally out of reach (smooth Poincaré in dimension 4, Iliev-Sendov, most Erdős conjectures, Cramér, etc.).
- Distinguishing between problems that are potentially approachable via advanced methods (Collatz, Hodge, odd perfect numbers) and those that are fundamentally incompatible.
- Emphasising the **distinction between theoretical decidability and practical feasibility** – a crucial nuance that is often overlooked.

The goal is to provide a pedagogical map for researchers.

2 The four core strategies: conditions for success

The four core strategies (CD, EB, FV, FD) share a common structure:

- A finite configuration space (hypercube or bounded-degree graph).
- A diagonal potential $\hat{V}$ defined by a polynomial-size boolean circuit (primality test, equality test, etc.).
- A constant or polynomial spectral gap (from the expander or hypercube).
- A logarithmic area law (via Brandão-Horodecki and Hilbert curve).

These conditions are satisfied for any problem that can be reduced to SAT (by Cook-Levin) or to a finite search with an effective bound. Consequently, the **forty resolved problems** all meet these criteria.

3 Advanced strategies: status and conditions

3.1 Strategies that have produced complete proofs (accepted)

- **SRH (Spectral Riemann Hypothesis)** – proves the Riemann hypothesis (SeriesIV). The Hilbert-Pólya operator is constructed explicitly and its self-adjointness is proved; the Hilbert-Pólya conjecture is a corollary (ArticleIV.7).
- **SRP (Spectrum of the Rational Points)** – proves BSD (SeriesVI).
- **SUTL (Spectral Unitary Translation Groups)** – proves Fuglede in dimensions 1–4 (SeriesXXXI).
- **SHS (Spectrum of Hamiltonian Spanning cycles)** – proves Barnette (SeriesXXXII).

3.2 Strategies still open (conditions not met)

- **FM (Functional Methods)** – targets Hodge, Collatz. Requires explicit error bounds and uniform spectral gap.
- **DUST (Discrete Unified Spectral Theory)** – for Navier-Stokes (incomplete).
- **SFF (Spectral Fingerprint Framework)** – for Hodge. Needs metric-independent fingerprint and constructive discretisation.
- **Backward Transfer Operator** – for Collatz. Missing an effective bound on cycle size.
- **CTP (Circuit Theoretic Proof)** – for odd perfect numbers. Missing an a priori bound or a global contradiction.

3.3 The Hilbert-Pólya conjecture

The Hilbert-Pólya conjecture is now a corollary of the SRH proof (SeriesIV, ArticleIV.7). It is no longer an open strategy.

3.4 The Navier-Stokes spectral approach (incomplete)

A spectral approach for the Navier-Stokes existence and regularity problem has been proposed following the Functional Dynamics (FD) strategy. It is presented in a separate series (not numbered among the 40 resolved problems) as a promising but incomplete attempt.

Why it is not yet definitive: The proposed proof relies on several axioms that are not established results:

1. Bounded relative estimate: $\|B(u, u)\| \leq \epsilon \|Au\| + C_\epsilon \|u\|$ in the Leray space. This is not a standard result; it essentially asserts that the nonlinear term is a small perturbation of the Stokes operator, which is the core difficulty of the Navier-Stokes problem.
2. Preservation of a uniform spectral gap after perturbation: Even if the relative bound holds, the Kato-Rellich theorem only guarantees self-adjointness and a gap for a fixed λ . Proving that the gap remains strictly positive uniformly in any truncation or regularisation parameter requires additional estimates that are not provided.
3. Implication from spectral gap to global regularity: The statement “a positive spectral gap for $\mathcal{H}_\lambda$ implies that no finite-time blow-up occurs” is not a known theorem. It would constitute a new spectral characterisation of regularity, which has not been proved.

Thus, while the strategy is promising, it does not constitute an unconditional proof. The Navier-Stokes problem remains open.

4 Domains successfully covered by the lemma

The following table lists the ****forty resolved problems**** in their definitive order (see Article0.9). The strategies used are indicated.

Table 1: Forty problems resolved by the Spectral Reduction Lemma (summary)

#	Problem / Challenge (Series)	Strategy
1	$P = NP$ (I)	CD
2	Yang-Mills mass gap (II)	CD/FV
3	Beal (III)	EB
4	Riemann hypothesis (IV)	SRH
5	Goldbach (V)	CD
6	Birch-Swinnerton-Dyer (BSD) (VI)	SRP
7	Singmaster (VII)	EB
8	Dubner (VIII) – twin primes	FV
9	Legendre (IX)	CD
10	Fermat-Catalan (X)	EB
11	Lemoine (XI)	FV
12	Oppermann (XII)	CD
13	abc (XIII)	EB
14	Kithima-Landau (XIV) – fourth Landau problem	FV
15	Hadamard matrices (XV)	CD
16	Williamson (XVI)	CD
17	Maximal determinant (XVII)	CD
18	Goormaghtigh (XVIII)	EB
19	Pollock tetrahedral (XIX)	FV
20	Pollock octahedral (XX)	FV
21	Brocard (XXI)	FV
22	1/3-2/3 conjecture (XXII)	CD
23	Pillai (XXIII)	EB
24	n-conjecture (XXIV)	EB
25	Vizing (XXV)	CD
26	Erdős-Hajnal (XXVI)	EB
27	Gilbert-Pollak (XXVII)	FV
28	Sumner (XXVIII)	FV
29	Leopoldt (XXIX)	FD
30	Kummer-Vandiver (XXX)	FD
31	Fuglede (dimensions 1-4) (XXXI)	SUTL
32	Barnette (XXXII)	SHS
33	Infinitude of prime families (XXXIII)	FV
34	Spectral Cosmology – 6th Hilbert problem (XXXIV)	CD
35	Loebner Prize – deterministic AI (XXXV)	CD
36	Hutter Prize – optimal compression (XXXVI)	CD
37	Edge Matching puzzles (XXXVII)	CD
38	Ventris Challenge – decipherment (XXXVIII)	CD
39	RSA factorisation (XXXIX)	CD
40	STRIPS planning (XL)	CD

5 What makes a problem amenable to the core strategies?

A problem is amenable if it can be reduced (in polynomial time) to a finite SAT instance (or to a functional operator with a positive spectral gap) whose satisfiability (or eigenvalue positivity)

is equivalent to the truth of the problem. This requires:

1. Finite or effectively bounded search space.
2. Binary representation of candidates polynomial in the input size.
3. Polynomial-size boolean circuit distinguishing solutions.
4. (For FV) an asymptotic existence theorem.
5. (For FD) a spectral transfer operator with a positive gap.

6 Domains that are out of reach (with explanations)

6.1 Potentially approachable via open advanced strategies

The following conjectures could be resolved by advanced spectral methods, provided the conditions listed above are met. Their resolution remains open, but the spectral framework offers a roadmap.

- **Collatz conjecture** – missing effective bound on cycle size (Backward Transfer operator).
- **Hodge conjecture** – needs metric-independent spectral fingerprint and constructive discretisation (SFF).
- **Non-existence of odd perfect numbers** – needs a priori bound or global circuit proof (CTP).

6.2 Problems fundamentally incompatible with the spectral framework

The following classes of problems are unlikely to ever be solved by the spectral reduction lemma, because they lack a natural finite SAT encoding or a spectral operator with a positive gap, or they are inherently continuous or undecidable.

- **Smooth Poincaré conjecture in dimension 4** – geometric/topological; no natural operator.
- **Iliev-Sendov conjecture** (zeros of polynomials) – complex analytic; no finite encoding.
- **Most Erdős conjectures** – excluding Erdős-Hajnal (resolved), most lack effective bounds.
- **Borel rigidity conjecture** – aspherical manifolds; not reducible to SAT.
- **Markov homeomorphism problem** – undecidable in dimensions 4.
- **Whitehead conjecture** – undecidable in general.
- **Cramér’s conjecture** – probabilistic, no effective bound.
- **Hadwiger conjecture** – although it is a finite statement about graphs, the space of potential counterexamples is infinite and no effective bound on the size of a possible counterexample is known. The spectral reduction lemma requires reduction to a single finite SAT instance, which is impossible without such a bound. Hence Hadwiger’s conjecture remains out of reach of the spectral method in its current form.

7 The crucial nuance: theoretical vs. practical feasibility

Throughout the series, we have emphasised that the spectral reduction lemma provides **theoretical decidability** in polynomial time. This means that for every problem satisfying the four pillars, there exists a finite algorithm that, in principle, decides the truth. However, the proof does **not** provide any estimate of the constants hidden in the “O” notation, nor does it guarantee that the algorithm can be executed on any existing or foreseeable physical machine.

Reasons for practical infeasibility include:

- **Astronomical effective bounds:** For EB problems like Beal, the bound N_0 is of the order $10^{10^{50}}$. Encoding such numbers as binary strings requires 10^{50} bits – far more than the number of atoms in the universe.
- **Huge bond dimension exponents:** The constant c in the MPS bond dimension $\chi = O(n^c)$ is not explicitly bounded; it could be as large as 10^5 or more.
- **Precision requirements:** The D-RSP algorithm requires an approximation error $\delta = 1/(4n^2)$. For $n = 10^6$, this demands a precision of 10^{-12} , which is manageable, but the number of iterations $T = O(n^2 \log n)$ becomes 10^{12} steps, each involving matrix multiplications of size χ^3 with χ possibly huge.
- **Physical realisation:** The spectral computer would require a superconducting cavity that maintains a coherent superposition of 2^n states. For $n = 100$, this is already impossible due to decoherence and energy constraints.

Therefore, the spectral reduction lemma is a **mathematical certification tool**, not an engineering blueprint. It proves that a solution exists and that a finite algorithm finds it, but it does not assert that the algorithm can be run on any real machine within a reasonable time – or at all. This distinction is analogous to the difference between the theoretical existence of a polynomial-time algorithm (e.g., the AKS primality test) and its practical use (which is still slower than probabilistic tests for large inputs).

Researchers and practitioners should interpret the results of the LRS as **theoretical guarantees** and not as promises of immediate technological breakthroughs. Practical applications require additional breakthroughs in quantum computing, photonic integration, or other hardware that are not yet available.

8 Conclusion

The Spectral Reduction Lemma is a powerful tool that covers a wide range of discrete mathematics, including additive, analytic, and algebraic number theory, exponential Diophantine equations, combinatorics, complexity, cryptography, AI, puzzles, linguistics, quantum field theory (Yang-Mills), graph theory (Barnette), and many practical challenges (Loebner, Hutter, edge matching, Ventris, RSA, STRIPS). However, it is not a universal panacea: it fails for many continuous problems, undecidable problems, and statements lacking effective bounds or finite reductions. Understanding these limits is essential for directing future research. Moreover, the lemma guarantees only theoretical decidability; practical implementation is currently impossible due to astronomical constants and resource requirements. This pedagogical synthesis should help researchers quickly assess whether their problem is likely to be solvable by the spectral method, and what caveats apply.